

Homebound by COVID19: The Benefits and Consequences of Non-pharmaceutical Intervention Strategies

Buse Eylul Oruc

Georgia Institute of Technology College of Engineering

Arden Baxter

Georgia Institute of Technology College of Engineering

Pinar Keskinocak (✉ pinar@isye.gatech.edu)

Georgia Institute of Technology College of Engineering <https://orcid.org/0000-0003-2686-546X>

John Asplund

Georgia Institute of Technology College of Engineering

Nicoleta Serban

Georgia Institute of Technology College of Engineering

Research article

Keywords: agent-based disease modeling, non-pharmaceutical intervention strategies, COVID19

Posted Date: August 18th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-51320/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Public Health on April 6th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-10725-9>.

Abstract

Background. Recent research has been conducted by various countries and regions on the impact of non-pharmaceutical interventions (NPIs) on reducing the spread of COVID19. This study evaluates the tradeoffs between potential benefits (e.g., reduction in infection spread and deaths) of NPIs for COVID19 and being homebound (i.e., refraining from community/workplace interactions).

Methods. An agent-based simulation model to project the disease spread and estimate the number of homebound people and person-days under multiple scenarios, including combinations of shelter-in-place, voluntary quarantine, and school closure in Georgia from March 1 to September 1, 2020.

Results. Compared to no intervention, under voluntary quarantine, voluntary quarantine with school closure, and shelter-in-place with school closure scenarios 3.43, 19.8, and 200+ homebound adult-days were required to prevent one infection, with the maximum number of adults homebound on a given day in the range of 121K-268K, 522K-567K, 5,377K-5,380K, respectively.

Conclusions. Voluntary quarantine combined with school closure significantly reduced the number of infections and deaths with a considerably smaller number of homebound person-days compared to shelter-in-place.

Background

Recent research and experiences from various communities around the world highlighted the potential benefits of non-pharmaceutical interventions (NPIs) for slowing down the spread of COVID19 and reducing the severe health outcomes [1-3]. NPIs include school closure, reducing public gatherings, social distancing, restricting travel, and voluntary quarantine (entire household staying at home if someone in the household has symptoms) [4-7] and more stringent interventions such as shelter-in-place [8, 9].

People may become “homebound” (i.e., stay home and refrain from interactions in the community/workplace) due to complying with some of the NPIs (even if they do not experience symptoms), showing symptoms, or providing childcare. Hence, despite their benefits, there are also unintended consequences of NPIs, including the impact on the economy, unemployment, household spending, mobility, energy usage, etc. [10-13] and the social impact on caring for the elderly, education of the young, family support, domestic violence, and personal health and wellbeing [14-23].

Some NPIs, such as shelter-in-place, apply to large populations for an extended duration, whereas others, such as voluntary quarantine, impact targeted populations for a limited time. It is important to understand the tradeoffs between the public health benefits and other consequences of NPIs, particularly, as measured by homebound person-days or the size of the homebound population over time. There is sparse research on assessing which interventions have a higher overall impact in reducing societal interactions versus the ability to reduce infection spread and adverse outcomes [8, 9, 24, 25].

This study evaluates the trade-offs between the public health impact measures (e.g., the number of cases, hospitalizations and deaths [26]) and intervention metrics, including number of homebound people and person-days under various NPI scenarios, including variations of shelter-in-place, voluntary quarantine, and school closure. The intervention metrics aim to capture how much an intervention reduces societal activity and interaction, much needed to maintain economic and social life. Such evaluations can assist local and national decision makers in choosing different combinations of targeted interventions over time to reduce infection spread while considering the societal and economic impact.

Methods

Intervention Analysis

The following NPIs, with varying combinations and compliance levels in different scenarios (*Figure 1*), are analyzed in this study and compared to the baseline of no intervention (NI):

1. School Closure (SC) – No peer-group interactions among children or youth (i.e., no K-12 school interactions).
2. Voluntary Quarantine (VQ) – Household members stay home if any member of the household is symptomatic, until the entire household is symptom-free.
3. Shelter-in-Place (SIP) – Household members stay home.

Modeling Case Projection and Estimating Intervention Impact

An agent-based simulation model with heterogeneous population mixing was utilized and adapted, which has been previously applied to project the number of COVID19 infections and severe outcomes under various social distancing strategies [26]. The study period is March 1, 2020- September 1, 2020.

The population in the simulation includes children (ages 0-9), youth (ages 10-19), adults (ages 20-64), and elderly (ages 65+). The simulation monitors the health status (e.g., symptomatic, hospitalized, dead) as well as the homebound status of each household member (for further details see Supplementary Section A, Additional File 1 and Supplementary Table 1, Additional File 1).

- Homebound: For adults and elderly, this status is defined as staying home due to voluntary quarantine, symptoms, shelter-in-place, or *at home childcare*, i.e., providing supervision to a child who is home due to their status (e.g., due to symptoms or school closure). For example, if a child is at home in need of supervision, the status of an adult or elderly member in the household is updated to indicate that they provide supervision, labeled as *at home childcare*. For children and youth, homebound is defined as staying home due to voluntary quarantine, symptoms, or school closure.
- Inactive: For adults and elderly, a status of inactive refers to being inactive from society due to being homebound, hospitalized, or providing hospital care, i.e., caring for a child or youth who became hospitalized. A status of inactive for children and youth is defined as being inactive from society due to being homebound or hospitalized.

Infection Spread Outcome Measures and Intervention Metrics

The infection spread outcome measures reported for the study period include:

- Cumulative deaths: Number of people who died due to COVID19.
- Cumulative infections: Number of people infected (including asymptomatic infections).
- Peak day: The day when the number of new infections was highest.
- Peak infection: The highest number (or percentage) of the population infected on a given day.

A statistical summary of infection spread outcome measures under baseline and intervention scenarios is provided in Supplementary Table 2, Additional File 1.

The infection spread measures are contrasted with the following intervention metrics, which are reported for the study period:

- Homebound or inactive subpopulation: Number of people in a subpopulation (adults/elderly or children/youth) with homebound or inactive status, respectively, on a given day.
- Percentage of days adults homebound or inactive: Average percentage of days an adult has homebound or inactive status, respectively.
- Homebound days: Average number of days a (sub)population has homebound status.
- Homebound or inactive peak day: The day when the number of a (sub)population has homebound or inactive status, respectively, is highest.
- Homebound or inactive peak: The highest number (or percentage) of a (sub)population homebound or inactive, respectively, on a given day.
- Adults absent from work: The number of adults who are absent from work due to an inactive status (further details are provided in Supplementary Section B, Additional File 1).
- Homebound days to prevent an infection: Additional adult homebound days needed to prevent an infection (in Scenario X, relative to Scenario 1), calculated as follows:

$$\frac{\text{Adult Homebound Days in Scenario X} - \text{Adult Homebound Days in Scenario 1}}{\text{Cumulative Infections in Scenario 1} - \text{Cumulative Infections in Scenario X}}$$

- Homebound days to prevent a death: Additional adult homebound days needed to prevent a death (in Scenario X, relative to Scenario 1), calculated as follows:

$$\frac{\text{Adult Homebound Days in Scenario X} - \text{Adult Homebound Days in Scenario 1}}{\text{Cumulative Deaths in Scenario 1} - \text{Cumulative Deaths in Scenario X}}$$

Results

Supplementary Table 2, Additional File 1 presents the infection spread outcome measures, including the population infected or dead and the peak infection.

Figure 2 presents the daily new infections and the homebound adults over time across all scenarios. Under Scenarios 1, 3a, 3b, 3c (non-school closure scenarios), the homebound peak for adults decreased from 267,566 under Scenario 3a to 121,346 under Scenario 3c, and the peak under Scenario 1 was 242,948. Under Scenarios 2, 4a, 4b, 4c, 5a, 5b, 5c (school closure scenarios), the homebound peak for adults was highest under Scenarios 5a, 5b, 5c, due to shelter-in-place, ranging from 5,377,886 to 5,379,960, followed by homebound peak of 584,235 under Scenario 2. Adults absent from work followed a similar pattern as homebound adults across all scenarios (Supplementary Figure 1, Additional File 1).

Higher compliance with voluntary quarantine reduced homebound peak for adults to 566,973, 535,559, 522,775 under Scenarios 4a, 4b, 4c, respectively (*Figure 2*), decreased the peak infection (in Scenarios 3a, 3b, 3c, 4a, 4b, 4c) by at least half, and delayed the peak day by 14-61 days compared to Scenario 1 (Supplementary Table 2, Additional File 1).

Figure 3 presents a comparison of the percentage of the population infected or dead and the percentage of days adults homebound. The percentage of the population infected was 60.09% under Scenario 1 (no intervention) and 51.69% under Scenario 2 (school closure only). The percentage of the population infected reduced to a range of 11.68-44.15% under Scenarios 3a, 3b, 3c (voluntary quarantine) and 4.53-31.07% under Scenarios 4a, 4b, 4c (voluntary quarantine with school closure). The percentage of days adults homebound was 0.72% under Scenario 1 and 7.16% under Scenario 2 (school closure only). The percentage of days adults homebound ranged from 1.36-1.63% and 7.62-7.74% under Scenarios 3a, 3b, 3c (voluntary quarantine) and Scenarios 4a, 4b, 4c (voluntary quarantine with school closure), respectively. Compared to Scenario 2 (school closure only), Scenarios 5a, 5b, 5c (shelter-in-place with school closure) reduced the percentage of the total population infected from 51.69% to 48.11-50.55% but more than doubled the percentage of days adults homebound to a range of 18.92-30.66%. Supplementary Table 3, Additional File 1 provides the percentage of days children, youth, adults, and elderly are homebound across all scenarios.

Figure 4 presents the homebound days to prevent an infection or death. The homebound days to prevent an infection was 87 under Scenario 2 (school closure only) and over 200 under Scenarios 5a, 5b, 5c (shelter-in-place with school closure). The homebound days to prevent an infection was 1.9, 3.7, 4.7 under Scenarios 3c, 3b, 3a (voluntary quarantine), respectively, versus 14.4, 17.8, 27.3 under Scenarios 4c, 4b, 4a (voluntary quarantine with school closure), respectively. The homebound days to prevent a death was 30,650 under Scenario 2 (school closure only) and over 60,420 under Scenarios 5a, 5b, 5c (shelter-in-place with school closure). The homebound days to prevent a death was 500, 928, 1,130 under Scenarios 3c, 3b, 3a (voluntary quarantine), respectively, versus 4,050, 4,840, 7,360 under Scenarios 4c, 4b, 4a (voluntary quarantine with school closure), respectively.

Supplementary Table 4, Additional File 1 presents the homebound and inactive peak percentages for children, youth, adults, elderly, and the total population. Increasing voluntary quarantine compliance,

regardless of school closure, decreased the homebound and inactive peak percentage for adults, elderly, and the total population. Supplementary Figures 2 and 3, Additional File 1 present the homebound peak broken down by statuses for adults and elderly and for children and youth, respectively.

Supplementary Figure 4, Additional File 1 shows the percentage distribution of statuses (at home childcare, voluntary quarantine, symptoms) for the homebound peak for adults. At the homebound peak, among homebound adults: (i) Under Scenarios 2, 4a, 4b, 4c (school closure scenarios without shelter-in-place), 0.33%-27.26% and 72.75%-83.47% were symptomatic or providing at home childcare, respectively. (ii) Under Scenarios 3a, 3b, 3c (non-school closure scenarios), 3.61%-26.59% and 0.89%-4.04% were symptomatic or providing at home childcare, respectively. (iii) Under no intervention, 89.99% and 10.01% were symptomatic or providing at home childcare, respectively.

Supplementary Tables 5-7, Additional File 1 summarize the impact of voluntary quarantine, school closure and shelter-in-place by comparing the percentage difference between a pair of scenarios in terms of the homebound days (for children, youth, adult and elderly populations), cumulative infections, and deaths.

Discussion

Many governments are faced with difficult decisions about when and how quickly to lift social distancing restrictions and reopen their economies; hence, it is crucial to analyze the benefits of NPIs in decreasing the spread of COVID19 versus the economic and social consequences considering the people who become homebound due to illness or due to complying with NPIs. In the state of Georgia, school closure began on March 16th [27] and a shelter-in-place order was enacted from April 3rd to April 30th, and the reopening of the state continued gradually since then [28]. The number of new COVID19 confirmed cases in Georgia have rapidly increased since early June [29].

This study focused on the public health benefits versus the need to refrain from societal or workplace interactions due to NPIs. The COVID19 pandemic led to widespread school closure and shelter-in-place orders in the United States [30, 31]. Despite the potential public health benefits, there were many concerns about the economic impacts of shelter-in-place [10-13] and the disruptive effects of school closures on the education of children and youth [14, 21-23, 32]. This study analyzed and compared several NPI scenarios, including combinations of school closure, voluntary quarantine, and shelter-in-place, with varying compliance levels and durations, as well as baseline scenarios of no intervention (Scenario 1) and school closure only (Scenario 2).

School closure alone had limited impact on reducing the spread of COVID19. Compared to no intervention, school closure only reduced the percentage of the population infected by less than 10% while more than doubling the peak number of adults homebound and causing nearly 450,000 work absences, the majority of which due to the need to provide at home childcare.

Shelter-in-place combined with school closure (Scenarios 5a-5c) temporarily slowed down the infection spread and delayed the peak, but had little impact on the magnitude of the peak and the cumulative number of infections and deaths, which were similar to that observed in the school closure only scenario. However, under Scenarios 5a-5c, the peak number of homebound adults was 9-44 times larger than all other intervention scenarios. Hence, the limited positive public health impact of shelter-in-place came at a very high societal cost.

Under voluntary quarantine (Scenarios 3a, 3b, 3c) the percentage of the population infected was 11.68%-44.15% (compared to 60.09% under no intervention), with the peak number of adults homebound being 267,566 under low, 211,695 under medium, and 121,346 under high compliance. Under voluntary quarantine combined with school closure (Scenarios 4a, 4b, 4c) the percentage of the population infected was 4.53%- 31.06% (compared to 51.69% under school closure only), with the peak number of adults homebound being 566,973 under low, 535,559 under medium, and 522,775 under high compliance. Higher levels of voluntary quarantine compliance decreased the percentage of the population infected and the peak number of adults homebound (or absent from work).

Voluntary quarantine compliance provided the greatest benefits in terms of the reduction in infections and deaths compared to the number of adults homebound. Compared to school closure only, voluntary quarantine combined with school closure yielded up to a 92% decrease in cumulative infections and deaths while homebound days increased by at most 8% for adults, 6% for elderly and 1.5% for the total population. Under voluntary quarantine scenarios, the number of homebound days to prevent an infection or death was 3-154 times lower than that of all other scenarios.

Conclusion

While large-scale interventions such as shelter-in-place temporarily slow down the infection spread, they are highly disruptive to the society and their public health impact is limited unless they are imposed for long durations of time, with high compliance levels, or followed by additional interventions.

Targeted interventions such as voluntary quarantine or voluntary quarantine combined with school closure significantly reduce the infection spread without causing a social and economic disruption as in the case of an extended shelter-in-place.

Strong public messaging should continue about voluntary quarantine, voluntary shelter-in-place (if possible), as well as other practices of physical distancing and the usage of facemasks.

Some of the conclusions of the study may be generalized to other states/countries that have geographic and population characteristics similar to the state of Georgia. The model and analysis would need to be adjusted for other pandemics; for example, COVID19 leads to fewer adverse health outcomes in younger populations and this may explain why school closure have a lesser impact on reducing infection spread.

Limitations

If facemask usage was also considered in the NPI scenarios, the relative reduction in the number of cases and deaths could be higher compared to baseline scenarios. The simulation was populated with data from the state of Georgia and the results presented may not apply to other states or regions which have significantly different population characteristics or density.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request. All data was publicly available.

Competing interests: The authors declare that they have no competing interests.

Funding: The research was supported by the William W. George and by the Virginia C. and Joseph C. Mello endowments at Georgia Tech. This research was supported in part by NSF grant MRI 1828187 and research cyberinfrastructure resources and services provided by the Partnership for an Advanced Computing Environment (PACE) at the Georgia Institute of Technology. The funders played no role in the study design, data collection, analysis, interpretation, or in writing the manuscript.

Authors' contributions: All authors contributed equally to this work.

Acknowledgements: The authors are thankful to state representatives for their support and guidance and for sharing data sources. The authors are also thankful to Georgia Tech students and colleagues Melody Shellman, Hannah Lin, Ethan Channel, Pravara Harati, April Zhuoting Yu, Gabriel Siewert, and Christopher Stone for their contributions and support.

References

1. Hartley DM, Perencevich EN: **Public Health Interventions for COVID-19: Emerging Evidence and Implications for an Evolving Public Health Crisis.** *JAMA* 2020.
2. Cowling BJ, Ali ST, Ng TW, Tsang TK, Li JC, Fong MW, Liao Q, Kwan MY, Lee SL, Chiu SS: **Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study.** *The Lancet Public Health* 2020.
3. Ngonghala CN, Iboi E, Eikenberry S, Scotch M, MacIntyre CR, Bonds MH, Gumel AB: **Mathematical assessment of the impact of non-pharmaceutical interventions on curtailing the 2019 novel Coronavirus.** *Mathematical Biosciences* 2020:108364.
4. Qualls N, Levitt A, Kanade N, Wright-Jegade N, Dopson S, Biggerstaff M, Reed C, Uzicanin A, Group CCMGW: **Community mitigation guidelines to prevent pandemic influenza—United States, 2017.** *MMWR Recommendations and Reports* 2017, **66**(1):1.

5. Tian H, Li Y, Liu Y, Kraemer MU, Chen B, Cai J, Li B, Xu B, Yang Q, Yang P: **Early evaluation of Wuhan City travel restrictions in response to the 2019 novel coronavirus outbreak.** *Medrxiv* 2020.
6. Lasry A, Kidder D, Hast M, Poovey J, Sunshine G, Zviedrite N, Ahmed F, Ethier KA: **Timing of community mitigation and changes in reported COVID-19 and community mobility—four US metropolitan areas, February 26–April 1, 2020.** 2020.
7. Peak CM, Childs LM, Grad YH, Buckee CO: **Comparing nonpharmaceutical interventions for containing emerging epidemics.** *Proceedings of the National Academy of Sciences* 2017, **114**(15):4023-4028.
8. Dave DM, Friedson AI, Matsuzawa K, Sabia JJ: **When do shelter-in-place orders fight covid-19 best? policy heterogeneity across states and adoption time.** In.: National Bureau of Economic Research; 2020.
9. Friedson AI, McNichols D, Sabia JJ, Dave D: **Did california's shelter-in-place order work? early coronavirus-related public health effects.** In.: National Bureau of Economic Research; 2020.
10. Wright AL, Sonin K, Driscoll J, Wilson J: **Poverty and economic dislocation reduce compliance with covid-19 shelter-in-place protocols.** *University of Chicago, Becker Friedman Institute for Economics Working Paper* 2020(2020-40).
11. Baker SR, Farrokhnia RA, Meyer S, Pagel M, Yannelis C: **How does household spending respond to an epidemic? Consumption during the 2020 COVID-19 pandemic.** In.: National Bureau of Economic Research; 2020.
12. Chen S, Igan D, Pierri N, Presbitero AF: **Tracking the Economic Impact of COVID-19 and Mitigation Policies in Europe and the United States.** 2020.
13. Couch KA, Fairlie RW, Xu H: **The Impacts of COVID-19 on Minority Unemployment: First Evidence from April 2020 CPS Microdata.** *Available at SSRN 3604814* 2020.
14. **Adverse consequences of school closures** [h
15. Armitage R, Nellums LB: **COVID-19 and the consequences of isolating the elderly.** *The Lancet Public Health* 2020, **5**(5):e256.
16. Wang G, Zhang Y, Zhao J, Zhang J, Jiang F: **Mitigate the effects of home confinement on children during the COVID-19 outbreak.** *The Lancet* 2020, **395**(10228):945-947.
17. Duan L, Zhu G: **Psychological interventions for people affected by the COVID-19 epidemic.** *The Lancet Psychiatry* 2020, **7**(4):300-302.
18. Pfefferbaum B, North CS: **Mental health and the Covid-19 pandemic.** *New England Journal of Medicine* 2020.
19. Campbell AM: **An increasing risk of family violence during the Covid-19 pandemic: Strengthening community collaborations to save lives.** *Forensic Science International: Reports* 2020:100089.
20. Bradley NL, DiPasquale AM, Dillabough K, Schneider PS: **Health care practitioners' responsibility to address intimate partner violence related to the COVID-19 pandemic.** *CMAJ* 2020.

21. Esposito S, Principi N: **School closure during the coronavirus disease 2019 (COVID-19) pandemic: an effective intervention at the global level?** *JAMA pediatrics* 2020.
22. Sharfstein JM, Morpew CC: **The Urgency and Challenge of Opening K-12 Schools in the Fall of 2020.** *JAMA* 2020.
23. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, Mytton O, Bonell C, Booy R: **School closure and management practices during coronavirus outbreaks including COVID-19: a rapid systematic review.** *The Lancet Child & Adolescent Health* 2020, **4**(5):397-404.
24. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, Whittaker C, Zhu H, Berah T, Eaton JW *et al.*: **Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe.** *Nature* 2020.
25. Hsiang S, Allen D, Annan-Phan S, Bell K, Bolliger I, Chong T, Druckenmiller H, Huang LY, Hultgren A, Krasovich E *et al.*: **The effect of large-scale anti-contagion policies on the COVID-19 pandemic.** *Nature* 2020.
26. Keskinocak P, Oruc Aglar BE, Baxter A, Asplund J, Serban N: **The Impact of Social Distancing on COVID19 Spread: State of Georgia Case Study.** *medRxiv* 2020:2020.2004.2029.20084764.
27. **Kemp orders schools closed through end of school year** [h
28. **Kemp: Some Georgia businesses allowed to reopen April 24; shelter-in-place to end April 30** [h
29. **Georgia Department of Public Health COVID-19 Daily Status Report** [h
30. **Map: Coronavirus and School Closures** [h
31. **See Which States and Cities Have Told Residents to Stay at Home** []
32. Kuhfeld M, Soland J, Tarasawa B, Johnson A, Ruzek E, Liu J: **Projecting the potential impacts of COVID-19 school closures on academic achievement.** In.: Annenberg Institute at Brown University; 2020.

Figures




Figure 1

Scenarios. Description of the intervention scenarios considered in this study.




Figure 2

Homebound adults and daily new infections. Homebound adults and daily new infections over time. Scenarios 2, 4a, 4b, 4c include school closure.




Figure 3

Comparison of adults homebound to infected or dead populations. Percentage of days adults homebound compared to the percentage of the population infected (left figure) and dead (right figure).




Figure 4

Homebound days to prevent an infection or death. Homebound days to prevent an infection (left figure) or a death (right figure).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFigure4.tiff](#)
- [SupplementaryFigure3.tiff](#)
- [SupplementaryFigure2.tiff](#)
- [SupplementaryFigure1.tiff](#)
- [BMCSupplement.docx](#)