

Risk factors for poor health and performance in European broiler production systems.

CURRENT STATUS: UNDER REVISION

BMC Veterinary Research  BMC Series

Tommy A.A. Van Limbergen
PEHESTAT

✉ tommy@pehestat.be *Corresponding Author*
ORCID: <https://orcid.org/0000-0002-0517-0880>

Steven Sarrazin
Universiteit Gent Faculteit Diergeneeskunde

Ilias Chantziaras
Universiteit Gent Faculteit Diergeneeskunde

Jeroen Dewulf
Universiteit Gent Faculteit Diergeneeskunde

Richard Ducatelle
Universiteit Gent Faculteit Diergeneeskunde

Ilias Kyriazakis
Newcastle University School of Natural and Environmental Sciences

Paul McMullin
Poultry Health Services International

Jesús Méndez
Cooperativas Orensanas S.C.G.

Jarkko K. Niemi
Natural Resources Institute Finland

Sotiris Pappasolomontos
Vitatrace Nutrition Ltd.

Piotr Szeleszczuk
Warsaw University of Life Sciences Department of Pathology and Veterinary Diagnostics

Johan Van Erum
GALLUVET

Dominiek Maes
Universiteit Gent Faculteit Diergeneeskunde

DOI:

10.21203/rs.2.14519/v1

SUBJECT AREAS

Small Animal Medicine

KEYWORDS

Broiler health, European broiler production, Housing , Risk-factors, Statistical modelling

Abstract

Background Conventional broilers are currently one of the most efficient protein converters. Although decades of progress in genetic selection and feed formulation have led to high standards of efficient broiler production, still a lot of variability is found between farms and between successive flocks. The aim of this study was to investigate risk- and/or protective factors for poor health and performance in conventional broiler-farms in Europe by developing eight multivariable linear mixed models. Three different models were used to investigate mortality (overall, first week, after first week), three models for performance variables (growth, feed conversion, European production index) and two models were related to slaughterhouse data (i.e. dead on arrival and condemnation rate).

Results Several factors related to management and housing were significantly associated with health and performance of broilers. The following factors were associated with increased mortality: floor quality, neonatal septicemia, ventilation type and other professional activities of the farmer. The factors associated with performance were chick sex, coccidiosis infections, necrotic enteritis, dysbacteriosis, light intensity adaptations, ventilation type, comparing daily flock results with previous flock results by farmer, daily check of feed and water system and type of feed. For dead on arrival three risk factors were identified i.e. daily growth, type of light adaptation and type of drinkers system. For condemnation rate seven risk factors were found, i.e. type of drinking system, daily growth, feed withdrawal time, type of ventilation, house size, septicemia after seven days and type of feed.

Conclusions These results imply that a multifactorial approach is required with adaptations involving both improvements in management, housing, health programs and an increasing level of professionalism of the farmer in order to improve broiler performance and health.

Background

Poultry meat production has grown rapidly during the last 40 years and is expected to continue to grow [01]. The demand for meat is increasing because of population growth, rising income and urbanization, and poultry meat has shown the fastest rise in the last decades. In order to be as sustainable as possible, the ecological footprint of broiler meat production needs to be restricted. This can be done in different ways. One way is by genetic selection of birds with higher feed efficiency. The industry has been very successful and further improvements are made continuously.

However, the gap between the potential of the birds and the actual performance achieved under practical conditions in the field is getting wider.

A large number of infectious and non-infectious risk factors have already been identified to cause decreased performance and increased mortality in conventional broilers [2].

Avian pathogenic *E.coli* (APEC) for example, has been known for decades to cause disease and mortality in broilers [3]. APEC can persist in a dry environment, and dust in poultry houses may contain up to 10^6 colony-forming units of *E. coli* per gram [4]. The most important predisposing factor for APEC infections in broilers however is stress, which can be induced by a series of inappropriate husbandry and management practices [5]. There are only few epidemiological studies taking a more integrated approach with multivariable modeling in order to identify and weigh the different risk factors threatening performance and health of broilers in field conditions. Flock size, stocking density, use of paper underlay for feeding during the 1st week for example have been identified as risk or protective factors for first week mortality [6]. The same study also identified that age of the broiler house, heating system and control system for the air intake could be associated with broiler mortality after seven days. A study from Taiwan investigated risk factors for first week mortality and identified other risk and protective factors such as water cooled ventilation, high flock size and distance from hatchery < 50km [7].

Unfortunately, many experimental studies investigating the factors that can influence broiler production have not taken into account the dynamic nature of these processes, by only focusing on specific measurements taken at a specific point in time (e.g. investigating footpad dermatitis at slaughter), often leading to conflicting results [8,9,10].

More large scale epidemiological studies using an integrated approach under different environmental and epidemiological conditions are necessary for a better understanding of the multiple factors affecting performance and health in conventional broiler production.

The aim of this study was to investigate and quantify risk- and/or protective factors for different health and performance parameters in conventional broiler-farms in different EU member states.

Results

Descriptive results

General farm characteristics

A total of 358 broiler-farms were included in this study. Seventy-eight percent (n=278) of the farms stated that the production of broilers was their only activity on the farm; 80.1% (n=281) was part of a quality assurance scheme and 52.8% (n=186) were part of an integrated system with some degree of

standardized management procedures.

Housing

An overview of the descriptive results for subcategories “Floor quality”, “Ventilation” and “Heating” is provided in table 1. The average broiler farm contained three broiler houses of 24 years old with each of them housing approx. 24 000 broilers using an average density of 17.5 day old chicks per square meter. Almost half (n=160) of the mechanically ventilated houses also contained specific ventilators meant for the recirculation of air inside the broiler house. Natural ventilation was still present in 7.3% (n=26) of broiler farms. A cooling system of some type was present in 71.5% (n=256) of all broiler farms for the prevention of heat stress in broilers. The most common type of cooling system was based on the creation of a fog in the neighbourhood of the air inlets of the broiler house, *i.e.* 78.8% (n=201), of which the position of the fog system was located inside the house in 79.6% (n=160) of the cases. Pad cooling was present in 20.7% (n=53) of farms that had a cooling system.

Table 1: Descriptive results of the housing of the broiler farms included in the study.

Subcategory	Situation on farm	Number of farms ⁴	% of tot
Floor quality ¹	Smooth impervious	132	37.7
	Fair condition	46	13.1
	Cracked to some degree	165	45.9
	Compacted earth	15	4.2
Ventilation ²	Roof ventilation	104	29.1
	Cross ventilation	74	20.7
	Roof x Tunnel ventilation	72	20.2
	Tunnel	40	11.2
	Cross x Tunnel ventilation	37	10.3
	Natural ventilation	26	7.3
	Other	5	1.4
Heating system ³	Direct	194	54.3
	Indirect	164	45.7

¹Floor quality was scored into four categories from “smooth impervious” (best situation) to “compacted earth” (worst situation). ²The type of ventilation system used to refresh the air inside the broiler house. ³The heating system was classified into two categories: direct heating (CO₂ is produced inside the broiler house) and indirect heating (no CO₂ is produced inside the broiler house). ⁴ Number of farms that have this situation on farm, a total of 358 broiler farms participated in this study.

Feed and water supply

Two types of feed systems were used in broiler farms: pan feeders (n=268, *i.e.*74.9%) and feeding troughs (n=90, *i.e.*25.1%). Pan feeders had an average length of 1.97 cm per broiler and with feeding troughs an average length of 2.74 cm was available per broiler (based upon the number of broilers

placed in the broiler house). The type of feed provided to broilers was a mixed feed compound in 199 farms (*i.e.*55.6%), while the other farms used a concentrated feed and added other ingredient (*e.g.*, whole wheat (150 farms, *i.e.* 94.3%), maize (2 farms, *i.e.* 1.3%) or a combination of whole wheat and maize (7 farm, *i.e.*4.4%). The main sources of drinking water were municipal water (n=158 farms, *i.e.*44.0%) and by using a ground well (n=188 farms, *i.e.* 52.4%), for ground wells the average depth was 91.9 meter. About half of the farms (n=177, *i.e.* 49.4%) used water disinfection protocols to sanitize the drinking water, the most commonly used chemicals for disinfection were chlorine dioxide (n=90, *i.e.* 51.1%) and peroxides (n=39, *i.e.* 22.2%). The drinking system of the investigated broiler farms was on average 16 years old. Over 89% of all broiler farms (*i.e.* 319 farms) used nipple drinkers with 13.3 chicks per drinking nipple. A minority of the farms was equipped with cup drinkers (n=20, *i.e.* 5.6%) or round drinkers (n=19 farms, *i.e.* 4.8%).

Treatments and diseases

An overview of the responses regarding the main health problems in the investigated broiler farms is provided in table 2. The majority of investigated broiler farms used a standard anti-coccidial treatment in the feed (346 farms, *i.e.* 96.6%). Vaccination against coccidiosis was performed in all flocks in only 6 farms (*i.e.* 1.7% of farms). Alternating vaccinated against coccidiosis and flocks treated with anti-coccidials was done in 42 farms (11.8% of broiler farms). The most frequently used anti-coccidials in starter feed were narasin-nicarbazine (n=197, 54.9%), nicarbazine (n=58, 16.2%), monensin-sodium (n=39, 11.0%), narasin (n= 26, 7.2%) and salinomycin (n= 23, 6.3%). For grower feed the most commonly used anti-coccidials were monensin-sodium (n= 151, 42.1%), narasin-nicarbazine (n=121, 33.7%), salinomycin (n=38, 10.7%) and narasin (n= 31, 8.6%). In finisher feed the main anti-coccidials were narasin (n=219, 61.1%), monensin-sodium (n=55, 15.5%), salinomycin (n=43, 12.0%) and narasin-nicarbazine (n=24, 6.6%). When a withdrawal period was required for the anti-coccidial, the feed was on average two days free of anti-coccidial before loading (thinning included). Thinning of the flock before complete emptying of the broiler house was done in 70.1% of all broiler farms (n=251). About half of these broiler farms (n=125, 49.8%) restarted with in feed-anti-coccidials after thinning.

Table 2. Number and % of farms with health problems in 2016.

Health problems	No problem		Mild problem		Clinical problem	
	N ¹	%	N ¹	%	N ¹	%
Coccidiosis	232	64.9	94	26.3	32	8.8
Septicemia before 7 days of age	149	41.6	57	15.9	152	42.4
Septicemia after 7 days of age	154	43.0	61	17.0	143	39.9
Dysbacteriosis	217	60.6	94	26.2	47	13.1
Necrotic enteritis	274	76.4	68	18.9	17	4.7
Wet litter syndrome	145	40.4	190	53.1	23	6.5
High mortality	284	79.2	56	15.7	18	5.1
Bad flock uniformity	189	52.7	160	44.7	9	2.6

¹Number of farms, in total 358 farms participated in the study.

Specific broiler information

Approximately 56.9% of all questioned broiler farmers (n=204) knew the age of the breeder parent stock from which their day old chicks originated. The initial bodyweight of day old chicks was known by 83.1% (n=297) of all broiler farmers and 95% (n=282) of them used such information to adapt the production management e.g. by using a higher set temperature in case of a low initial bodyweight. The transport time between the departure of day old chicks from the hatchery and the arrival in the broiler house was less than four hours in 81.6% of all broiler farms (n=292), while 2.8% of all farmers (n=10) received day old chicks after more than eight hours. The majority of broiler farms (n=255) always received day old chicks as hatched (71.3%), *i.e.* with no sex differentiation.

Table 3. Descriptive results of the production management of the broiler farms included in the study.

Subcategory	Situation on farm	Number of farms ⁵	% of
Date of placement ¹	Complete area of the house ³	206	57.5
	Floor temperature measured ⁴	263	73.6
Light intensity	Adapted during production cycle	245	68.5
Litter material	wood shavings	102	28.5
	cut straw	90	25.1
	peat	46	12.8
	rice hulls	39	10.9
	complete straw	31	8.7
Daily data registration	water intake	314	87.7
	feed intake	146	40.8
	bodyweight	97	27.1
Light during catching ²	Adaptation of light	349	97.5
	- Decrease of light intensity	181	51.8
	- Use of red lights	105	30.1
	- Use of blue lights	63	18.1

¹Date when day old chicks are placed into the broiler house; ²Light that is used when broilers are caught for transport to slaughterhouse; ³Day old chicks have access to the full floor area of the broiler house; ⁴Floor temperature is measured when day old chicks are placed into the house; ⁵Number of farms that have this situation on farm.

Production management and performances

An overview of different aspects of the production management of the farms included in the study was provided in table 3. Farmers tended to start preheating their broiler houses on average 34.7 hours before placement with an average set temperature of 33.2°C. During the first three days after placement, a lighting schedule of an average of 68.2 hours of light (*i.e.* 94.2% of the time) was used in the broiler house. During the rest of the production period, an average of 5.9 hours of darkness per 24 hours was provided to the broilers.

At 15.5% of broiler farms (n=55) extra drinkers were provided to the day old chicks and these were removed after 4.9 days on average. Almost all broiler farms (n=339, 94.6%) provided extra feeders for day old chicks. The majority of broiler farmers (n=273, 76.3%) checked crop fill of chicks during the first 24 hours after arrival on the farm. The average feed withdrawal time before loading was 7.3 hours, while water was withdrawn only very shortly before loading, *i.e.* on average 0.5 hours before loading. A number of parameters can be registered automatically on a daily basis in broiler farms, farmers were questioned about the registration of average body weight, average feed intake and average water intake (see table 3). The majority of farmers (n=348, 97.2%) compared the recorded information with data from previous batches or general schemes. The flow rate of the drinking system was checked daily by 53.9% of all farmers (n=193), while 42.2% of farmers (n=151) did not check water flow rate every day but checked flow rates as soon as abnormal fluctuations in the daily water intake were observed. Drinking nipples and feeding system were checked daily by 94.1% of the farmers (n=337). Almost all broiler farmers (n=352, 98.3%) stated that when abnormal birds (*e.g.* runt or lame birds) were observed during daily inspection, these animals are culled. Forty percent of the questioned farmers (n=143) stated that they received no information at all from the slaughterhouse regarding health and/or welfare parameters of their broilers.

A summary of the specific health and performance parameters for broiler farms is provided in table 4. These values are based on 2309 flocks from 358 broiler farms.

Table 4. Performance and health parameters based on 2309 flocks from 358 broiler farms in 2016 originating from 7 EU member states.

Parameter	Average	Median	SD ¹	Minimum	Maximum
Average age at slaughter (days)	41.33	41	3.62	30.26	58.0
Average weight at slaughter (kg)	2.47	2.50	0.43	1.66	3.31
Overall mortality (%)	3.82	3.70	1.40	1.00	14.8
First week mortality (%)	0.94	0.90	0.51	0.03	3.29
Dead on arrival (%)	0.20	0.14	0.37	0.01	4.60
Condemnation rate (%)	1.23	1.00	0.92	0.05	6.73
Feed conversion rate	1.74	1.70	0.17	1.23	2.06
Daily gain (gram/day)	59.79	60.62	5.89	41.19	72.9
EPI ²	338.41	345.85	53.07	183.93	432.1

¹Standard deviation; ²European Production Index, calculated by multiplying average bodyweight with livability, dividing this result by the product of FCR and average age. This result multiplied with “100”, provides the EPI. Livability is defined as the percentage of the total number of broilers at placement that reaches slaughter-age.

Risk factor analysis

Eight linear mixed models were fitted based on these health and performance parameters. A summary of the significant risk factors is provided in table 5.

The first linear model included overall mortality as the dependent variable. The following three independent variables were identified as potential risk factors: mortality rate after 7 days; floor quality and the occurrence of neonatal septicemia on the farm. The higher the mortality rate after 7 days, the higher the overall mortality was in the investigated flocks. Poor floor quality was a significant risk factor for overall mortality in broilers as farms with lower quality floors with cracks, in their broiler houses appeared to have significantly higher overall mortality rates. The third risk factor for overall mortality was the occurrence of problems with neonatal septicemia.

Four significant risk factors were found for first week mortality: floor quality, ventilation type, presence of other professional activities of the farmer and the occurrence of problems with neonatal septicemia. The risk was higher in farms with cracked floors compared to floors without cracks, in farms using tunnel or roof ventilation, in farms in which the farmer had multiple professional activities and in farms experiencing neonatal septicemia in combination with tunnel ventilated broiler houses.

For mortality after 7 days, only one significant risk factor was found *i.e.* poor floor quality (cracks are present in the floor in which pathogens might survive from previous flocks) of the broiler house.

Table 5. Multivariable linear mixed models related to performance and health parameters in 2309 flocks from 358 broiler farms in 7 EU member states.

Model	Dependent variable	Independent variable ¹	p-value
1	Overall mortality rate	Mortality rate after 7 days	<0.001
		Floor quality ²	<0.001
		Neonatal septicemia	<0.001
2	First week mortality rate	Floor quality	0.004
		Ventilation type ³	<0.001
		No other professional activities by farmer	<0.001
		Neonatal septicemia ⁴	0.014
3	Mortality rate after seven days	Floor quality	0.046
4	Dead on arrival	Light management during catching ⁵	0.003
		Type of drinking system	<0.001
		Daily growth (g/d)	<0.001
5	Condemnation rate	Type of drinking system	0.001
		Daily growth (g/d)	<0.001
		Feed withdrawal time	<0.001
		Number of birds in the broilerhouse	0.002
		Presence of recirculation vents	0.039
		Septicemia after seven days	0.012
		Type of feed	<0.001
Ventilation type ^{3,6}	0.057		
6	Feed conversion rate	Daily growth (g/d)	<0.001
		Light intensity adaptations	0.013
		Ventilation type ³	0.001
		Necrotic enteritis problems	0.014
		Daily check of drink water flow	0.002
7	Daily growth	Feed conversion rate	<0.001
		Coccidiosis problems	0.004
8	European production index	Sex of day old chicks ⁷	< 0.001
		Dysbacteriosis problems	0.002
		Evaluation of daily registered results	<0.001
		Daily inspection of feed and water system	0.029
		Type of drinking system ⁸	0.006
Type of feed ⁹	0.024		

¹Only the statistically significant risk factors in the final models are presented; ²The reference used for floor quality was a floor in perfect conditions without cracks; ³The reference used for type of ventilation was roof ventilation; ⁴This was only the case in tunnel-ventilated broiler houses; ⁵The reference for light adaptation was dimming the light intensity; ⁶Interpreted as a trend, as it was not significant (p >0.05); ⁷If only male chicks were housed, a higher EPI was present; ⁸The reference used for drinking system was the nipple drinking system; ⁹Only the case when dysbacteriosis was absent in the flock.

The analysis for farm specific risk factors for high number of dead on arrival (DOA) to slaughterhouse identified three significant variables: the type of light adaptation when broilers are caught, the drinking system and the daily growth during the cycle. The use of red lights for catching broilers

increased the risk for dead on arrival in the slaughterhouse in comparison with decreased light intensity. The other methods of light adaptation had no significant effect. Other significant risk factors were not using nipple drinkers system and higher daily growth.

Seven significant risk factors and one borderline non-significant factor could be associated with condemnation rate: drinking system, daily growth, feed withdrawal time, ventilation system, average house size, the presence of recirculation vents, problems with septicemia after the first seven days and the type of feed that was used during the grower phase. The usage of cup drinkers was associated with a lower condemnation rate in comparison with nipple drinking systems, while no significant effects were noticed for the other types of drinkers. A significant positive association was found between increasing daily growth and increasing condemnation rate at slaughterhouse. Sufficient feed withdrawal time was associated with lower condemnation rates. Condemnation rates further increased with higher capacity of a broiler house (*i.e.* the number of birds that can be housed in the same airspace), in case no recirculator vents (which are used to mix up the air in a broiler house) were present in the broiler house, when there were septicemia problems after seven days of age and in case of concentrate + wheat was used compared to the use of complete compound feed. Although not statistically significant, there was a trend that condemnation rate was lower in cross ventilated broiler houses in compare to roof ventilated houses.

Feed conversion rate (FCR) was significantly associated with five variables: daily growth (DG), the possibility to change light intensity, type of ventilation, the occurrence of necrotic enteritis (NE) and daily check of drinking water flow rate. A clear significant negative association was found between DG and FCR. Feed conversion rate was significantly higher in broiler houses that could not adapt the light intensity of the broiler house. A clear effect of the type of ventilation system was found on FCR. Roof ventilated broiler houses appeared to have the best FCR, followed by cross ventilated and cross x tunnel ventilated broiler houses. FCR was significantly lower in mechanically ventilated broiler houses compared to natural ventilated broiler houses. The absence of problems associated with NE was associated with a significantly better FCR. FCR was significantly better in farms that did not perform a daily check of the flow rate of the drinking system, but systematically checked the flow rate of the drinking system in case abnormal fluctuation of the water intake occurred.

For DG only FCR and the occurrence of problems with coccidiosis infections were risk factors. A negative association was found between FCR and DG. Farms that stated to have no problems with coccidiosis during the production period were found to have a significantly higher DG.

Six variables significantly influenced European Production Index (EPI): Sex of day old chicks (male chicks had a higher EPI), the occurrence of problems with dysbacteriosis, if farmers compared their daily registered results with results from previous batches, if farmers checked the drinkers and feeders on a daily basis, the type of drinkers and the type of feed. The EPI was significantly higher in

broiler farms that only housed male broilers compared to farms that housed only female broiler or a mix of both genders (as hatched). Farms that only housed female broilers had a significantly lower EPI. Farms that encountered no problems with dysbacteriosis had a significantly higher EPI compared to farms that had problems with dysbacteriosis. Farmers that did not compare their recorded information (e.g. bodyweight, feed- and water intake) with data from previous batches or general schemes had a significant lower EPI in comparison with farmers that compared flock results. Farmers who stated that they did not check the feeders and drinkers frequently had a significant lower EPI compared to farmers who did this on a daily basis. Farms with nipple or cup drinkers had a better EPI compared to farms with round drinkers. There was no difference between farms with nipple and cup drinkers. Broiler farms that had no problems of dysbacteriosis and used a concentrate with whole wheat, had a significant lower EPI compared to farms without problems of dysbacteriosis that used a complete compound.

Discussion

This study identified different management, housing and health factors as risk or protective factors for broiler performance and health on conventional broiler farms in seven EU member states.

A full randomization in the selection of the farms could not be established, a selection-bias is therefore likely to be present to some extent. Indeed, the method in which a questionnaire is presented has been identified to have an effect on data-quality [11]. For this reason, data collection by farm visits was encouraged, allowing a personal contact between the interviewer and the responsible of the poultry farm. Personal contacts were present in the vast majority of the collected data, but some data from two out of seven countries were unfortunately collected via conventional mail or telephone interviews instead of farm visits. Interviewers were asked to assure that the collected data was genuine in case no farm visits were possible or allowed, by cross-checking the replies to the questionnaires with data that was already recorded in these farms for other purposes, in order to limit the potential loss of data-quality. This questionnaire was designed to collect details about farm housing and management practices in a standardized way in all participating farms. The possible effects of interviewer bias on data quality were limited as much as possible by reducing the number of people responsible for the collection of data to an absolute minimum of maximum one person per country and also by training the interviewers to follow a specific standard procedure when performing the farm visit. The relatively high number of broiler farms that was included in this investigation increases the representativeness of the study population. The farm characteristics in the seven participating member states were in accordance with previous publications [12,13].

Three different models were constructed to investigate mortality rate in broilers, *i.e.* overall mortality, first week mortality and mortality after seven days. This classification was used because previous research about mortality rate in average broiler production was shown to follow typical patterns with a peak in mortality at day three and four after placement, a stable low mortality rate during the rest of the production cycle with a slight increased mortality rate during the sixth week of broiler production [14,15,16,17]. For this reason a distinction was made between mortality rate before and after seven days of placement [6]. It was impossible to distinguish the effect of culling in the current dataset. Culling programs might vary between different broiler producers and culling is used to remove birds that are about to succumb [15,18]. A proper culling program has beneficial effects on the FCR and flock uniformity [18].

In the current study, all types of mortality were associated with poor floor quality. Poor floor quality included the presence of cracks in the floor, which might be associated with mortality because of a higher pathogen load, due to a less efficient cleaning and disinfection of the broiler house [19].

Neonatal septicaemia was positively associated with both a higher overall and first week mortality. Avian pathogenic *E.coli* (APEC) has been known for decades to cause disease and mortality in broilers [3]. APEC can persist in a dry environment, and dust in poultry houses may contain up to 10^6 colony-forming units of *E. coli* per gram [4]. The most important predisposing factor for APEC infections in broilers is stress, which can be induced by a series of inappropriate husbandry practices [5].

First week mortality was also positively associated with the use of tunnel or roof ventilation in broiler houses. The use of tunnel ventilation appeared to have a higher association with first week mortality compared to the use of roof ventilation. This result is confirmed by other studies, who found a higher first week mortality in broiler houses with negative pressure ventilation compared to naturally ventilated houses [6,7].

When the farmer had other professional activities besides broiler production, the first week mortality appeared to be higher compared to farmers for whom broiler production was their sole professional activity. A possible explanation for this finding is that the farmers had less time available for the preparation of a broiler house and the management during the first week after placement. These things are crucial to provide the birds with an optimal start of the production cycle [6,7,16,17]. The combination of broiler production with other professional might lead to a less strict follow up of broilers, which might lead to a higher first week mortality.

Although several risk factors were identified to be related with the different types of mortality, previous studies identified additional risk factors. A Norwegian study by Heier et al. [6] identified flock size, stocking density, use of paper underlay for feeding during the first week as risk- or protective factors for first week mortality. The same study also identified age of the broiler house, heating system and control system for the air intake as risk factors for broiler mortality after seven days. A

study from Taiwan investigated risk factors for first week mortality and identified other risk and protective factors such as water cooled ventilation, high flock size and distance from hatchery < 50km [7]. The difference between the current study and the other studies that were mentioned is most likely due to different regional and climatic conditions.

The model that was designed to investigate risk factors for DOA identified three significant risk or protective factors. The use of red lights during catching appeared to result in a higher number of DOA compared to other types of light management, *i.e.* dimmed lights or blue lights. This was in accordance with the results of Prayitno et al. [20], who investigated the effects of color of lighting, *i.e.* blue, green, red or white light, on the behaviour and production of broilers. Prayitno et al. [20] found that broilers were more active in red light and showed greater floor-pecking, wing-stretching and aggression in the red light when compared to blue or green light. Birds kept in green or blue light were calmer, which is beneficial to reduce stress when broilers are caught. The use of nipple drinkers was found to be associated with a less DOA. However, this type of drinkers system was present in almost 90% of the broiler houses and has become the new standard in broiler houses. Houldcroft et al. [21] indicated that when choice is offered, broilers tend to prefer bell drinkers or troughs over nipple drinkers in order to express the stereotypic “scoop” action while obtaining water. Nipple drinkers have the important advantage that less water is spilled. Wet litter can cause hock burn and pododermatitis [21]. An important aspect in nipple drinkers is the height of the drinking lines, which can be adapted according to the size of the broilers and was shown to have a significant impact on broiler performance [21,22,23]. Unfortunately, the latter could not be investigated in the current study design. It is important to state that the conditions during transportation and in the waiting area of the slaughterhouses were not included in this analysis although they may have an impact on the dependent variable: “dead on arrival”.

A high DG was associated with higher DOA, which is in accordance with previous studies [24,25]. Other studies have clearly identified transport of broilers from farm to slaughterhouse as a major risk factor for DOA [26,27, 28]. In the current study, however, transport details after loading the broilers were not collected.

Condemnation rate in this study included all reasons for condemnation of carcasses in the slaughterhouse. It was not possible to obtain sufficient data from slaughterhouse reports which allowed a more detailed approach in investigating the different reasons for condemnation of broiler carcasses. Nevertheless our findings can provide an indirect indications for the reasons why carcasses were rejected. The presence of cup drinkers in the broiler house was linked to a higher condemnation rate compared to nipple drinking systems. Cup drinkers are known to result in more water spillage than nipple drinkers [21]. Wet litter is known to induce skin lesions in broilers at the plantar surface of the feet (“footpad dermatitis”), the caudal aspect of the intertarsal joint (“hockburn”) and over the

sternum (“breast blisters”) [21,29]. Increasing DG was associated with higher condemnation rates in our current study. Production of heavy broilers indeed results in a higher condemnation rate compared to the production of standard or light types of broilers [30]. The most common reasons for condemnation in that study were: “not identified”; “emaciation”; “congestion” and “infected skin lesions” [30]. Feed withdrawal time was negatively associated with condemnation rate in broilers. This might be due to contamination of carcasses by the content of crop and digestive tract that is still present when the withdrawal period is too short [31]. Withdrawal of feed for 8 to 12h before slaughter has the lowest risk for carcass contamination and has minimal losses for carcass weight [31,32]. Flock size was also previously identified as a risk factor for condemnation rate in broilers [30]. There is no clear explanation for this finding. The use of recirculation ventilators was found to be negatively associated with condemnation rate. The main beneficial aspect of such ventilators is that the air quality in the broiler house is more uniform and that a higher level of oxygen is available at the height of the broilers [33]. There was a trend for a lower condemnation rate in cross ventilated houses. A possible explanation might be the slightly lower air velocity in this type of ventilation. High air velocity (125m/min) has indeed been identified to have a negative effect on condemnation rates [34]. When septicemia occurred after 7 days of age, there was a significantly higher condemnation rate. APEC can cause aerocolitis, polyserositis, septicaemia and other mainly extraintestinal diseases in broilers, turkeys and other avian species [35]. APEC are found in the intestinal microflora of healthy birds and most of the diseases associated with them are secondary to environmental and host predisposing factors [35]. The use of a concentrated feed with whole wheat added to it appeared to be associated with a lower condemnation rate. Whole wheat enhances feed efficiency because of a better utilization of nutrients in commercial feeds [36]. The use of whole wheat leads to higher relative weights of gizzard and pancreas [37]. A direct relation between whole wheat feed and condemnation rate is unclear, but the positive effects on gizzard and pancreas might indirectly result in a lower condemnation rate by a lower presence of runts.

FCR was lower in farms that were able to adapt light intensity in the broiler house. It is known that light intensity can influence the activity level of growing broilers [38]. Farms that can adapt light intensity in the broiler house may stimulate or slow down the feed consumption of broilers and thereby improve performance parameters such as FCR. Mechanical ventilation was also found to be associated with a better FCR compared to natural ventilation. This is most likely due to a better air quality in general in mechanically ventilated houses as the growing chick has a high oxygen requirement in order to sustain rapid growth and optimal feed efficiency [39]. FCR was significantly higher in flocks that had necrotic enteritis problems. The latter was also shown in literature before by multiple authors [40,41,42]. In flocks where farmers checked the flow rate of the drinking system in case of abnormal fluctuations a tendency to a better FCR was seen. Maiorka et al. [43] showed that

the lack of water produced the same effect as the lack of feed, both causing a higher number of villi per area with reduction in villus size, when compared with feed and water ad libitum treatments. The model for DG in broilers clearly showed that the presence of coccidiosis problems was a risk factor. Coccidiosis has been shown by many authors to negatively influence performance [44,45]. As expected the sex of the day old chicks appeared to have a significant effect on the EPI. The EPI was significantly higher in flocks that contained only male birds, when compared to as hatched flocks. When only female birds were housed, the EPI was lower than compared with as hatched flocks. This sex-difference in performance was also shown by previous [46,47]. Farms that had recurrent problems of dysbacteriosis were found to have lower EPI. This is mainly due to the negative impact of dysbacteriosis on DG and FCR [45,48]. Both parameters are included in the formula that is used to calculate EPI, and thereby have a direct impact on EPI. The type of drinkers was also directly influencing the EPI, the effect of this factor on EPI was mainly due to its effect on FCR (see above). In our study the EPI was also influenced by the farmer's stockmanship, *i.e.* comparison of daily results with results of previous cycles and the daily check of the feed- and water system. It is well known that the level of expertise varies significantly between farmers and this may directly impact production results [49,50]. This further emphasizes the importance of farmer-education and an integrated approach of farm guidance e.g. by specialised poultry veterinarians and nutritionists [51,52].

Conclusion

This study showed that broiler health and performance are strongly influenced by many factors directly related to the broiler farm's management and housing. Also the impact of health problems caused by septicemia, coccidiosis and dysbacteriosis was shown to have a major influence on broiler performance. These health problems are known to have a multifactorial origin in which the presence of the primary pathogen on its own does not necessarily causes clinical problems and the impact of the primary pathogen on broiler production becomes higher when circumstances are in their favor. In this study a number of risk and protective factors for broiler health and performance were identified. For housing this mainly related to floor quality, ventilation, light management at catching and light intensity throughout the cycle. Management practices such as the level of professionalism, feed withdrawal time and light adaptations during the productive period also had an impact on broiler performance.

The results of this study imply that an multifactorial approach is necessary in order to improve health and performance in broiler production. This involves adaptations of management, housing and health

programs and increasing the level of professionalism of the farmer.

Methods

Farms

This observational study was performed in seven broiler producing EU member states: Belgium, Cyprus, Finland, Greece, Poland, Spain and the United Kingdom as a part of the EU-funded project PROHEALTH. In each EU member state, approx. 50 conventional broiler farms were included. Farms were selected by the locally assigned members of the PROHEALTH consortium to provide a representative population sample. It was agreed between the participating investigators within each member state that all participating farms in a member state needed to be as representative for the broiler-industry of that specific member state as possible, preferably by using a randomized sample of the local broiler-industry. Broiler farms that were producing in free-range or bio-label conditions were excluded. Upon request of the participating poultry companies and as agreed by the PROHEALTH consortium, member state codes were used throughout the analysis and the results.

Collection of Farm Data

All interviews were performed between February and August 2016. In Belgium, Cyprus, Greece, Spain and the United Kingdom, data were collected during farm-visits, in Poland, data were collected through conventional mail and telephone interviews, and in Finland through on-line survey and telephone interviews and from company databases. A protocol for the interviewers was developed by the first author. To ensure inter-farm comparability and to reduce interviewer bias as much as possible, all interviews were performed by maximum one person in each country. The interviewers involved in this investigation were instructed about the protocol of the interview during a PROHEALTH consortium workshop prior to the start of the farm-visits to standardize the approach in the different countries. An example of the questionnaire is shown in the supplementary data.

Two types of data were collected from each broiler farm, *i.e.* potential risk factors for poor health and production were collected using a questionnaire, and data on performance and health were collected from at least six subsequent flocks, *i.e.* at least over one year (including potential seasonal effects). The questionnaire contained a total of 67 questions, including both multiple choice and open-ended questions. The questions were classified into six categories, *i.e.* general information, housing, feed and water supply, treatments and diseases, broiler details and management. The questionnaire was tested on 10 broiler farms prior to its use in this project.

Performance and health parameters included “average age” (in days), “average weight” (in kg), “overall mortality” (expressed as percentage), “first week mortality” (expressed as percentage), “dead on arrival” to slaughterhouse (DOA, expressed as percentage), “condemnation rate” in slaughterhouse (expressed as percentage), “feed conversion rate” (FCR, calculated by dividing total amount of feed consumed by the birds (in kg), by total live weight of birds at slaughter (in kg)), “daily

weight gain” (DG, calculated by dividing the average live weight at slaughter by the average age of the birds at slaughter in days) and European production index (EPI), calculated by multiplying average bodyweight with livability, dividing this result by the product of FCR and average age. This result multiplied with “100”, provides the EPI. Livability is defined as the percentage of the total number of broilers at placement that reaches slaughter-age. Mortality in broilers in the current investigation was the sum of the number of dead birds that were found during daily inspection and the number of birds that were culled. Health problems that had occurred in one or more flocks during the twelve months before questioning were recorded. They included: coccidiosis, septicemia, dysbacteriosis, necrotic enteritis, wet litter syndrome, high mortality and poor flock uniformity.

Data and Statistical Analysis

Eight linear mixed models were developed. The following factors were used as dependent variables: overall mortality, first week mortality, mortality after seven days, DOA, condemnation rate, FCR, DG and EPI. If necessary, transformation of the dependent variables was considered to obtain a normally distributed outcome variable [53]. A random effect for country was included to correct for clustering of farms within a member state. Initially, univariable linear mixed regression models between the dependent variables and each predictor variable were examined. Furthermore, for continuous predictor variables, the assumption of linearity was examined by the Loess curves between each individual predictor variable and the dependent variables and by the scaled residuals of the univariable models. If necessary, transformation of the predictor variables was considered. Then, the independent variables were used to build a multivariable linear regression model by a manual stepwise forward model building procedure. The predictor variables ‘age’ and ‘stocking density’ were maintained in the model to correct for possible confounding. Variance inflation factors were examined to check for multicollinearity when adding predictors to the models and in case of multicollinearity, the biologically most relevant predictor was retained in the final model. Statistical significance during this step was assessed at $P < 0.05$. The estimates of the significant predictor variables are presented with their corresponding 95% confidence interval. Finally, all biologically plausible two-way interactions were tested and removed when non-significant ($P > 0.05$). To check whether the assumptions of normality and homogeneity of variance had been fulfilled, scaled residuals were examined. Only the results of the final models are shown. Statistical analysis were performed in SAS 9.4 (SAS Institute Inc., Cary, NC, USA).

Abbreviations

APEC: Avian pathogenic *E.coli*

DG: Daily Growth

DOA: Dead On Arrival in slaughterhouse

EPI: European Production Index

FCR: Feed Conversion Rate

NE: Necrotic Enteritis

Declarations

Ethics approval and consent to participate

No ethical approval was required for this observational study according to European and Belgian legislation involving experimental animals. No animal experiments were performed during this investigation.

A written informed consent was obtained from all farmers that participated in this study and farmers were aware that their (anonymized) data was used in the scope of this meta-analysis.

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by the European Union's Seventh framework Programme for research, technological development and demonstration under grant agreement no: 613574.

-

Authors' contributions

Conceived and designed the study: TVL, IK, PM, JM, JN, SP, PS and DM. Performed the data collection: TVL, PM, JM, JN, SP and PS. Analyzed the data: TVL and SS. Wrote and revised the manuscript: TVL, RD and DM. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank all participating broiler-farmers, integrators and farm-veterinarians for their kind cooperation, and the PROHEALTH consortium for their help with this study.

References

1. FAO - Food and Agriculture Organisation of the United Nations (2011) World Livestock 2011 - Livestock in food security. Rome, FAO.
2. Jones P.J., Niemi J., Christensen J-P., Tranter R.B., Bennett R.M.. A review of the financial impact of production diseases in poultry production systems. *Animal Production Science* <https://doi.org/10.1071/AN18281>.
3. Dziva F. and M.P. Stevens. 2008. Colibacillosis in poultry: unravelling the molecular basis of virulence of avian pathogenic *Escherichia coli* in their natural hosts. *Avian Pathology* 37:355-366.
4. Harry, E.G. 1964. The survival of *Escherichia coli* in the dust of poultry houses. *The Veterinary Record*, 76: 466-470.
5. Barnes, H.J. and Gross, W.B. 1997. "Colibacillosis". In *Diseases of Poultry*, 10th edn,

Edited by: Calnek, B.W., Barnes, H.J., Beard, C.W., McDougald, L.R. and Saif, Y.M.
131-141. Ames, IA: Iowa State University Press.

6. Heier B.T., Høgasen H.R. and J.Jarp. 2002. Factors associated with mortality in Norwegian broiler flocks. *Preventive Veterinary Medicine* 53:147-158.
7. Chou C.C., Jiang D.D. and Hung Y.P. 2010. Risk factors for cumulative mortality in broiler chicken flocks in the first week of life in Taiwan. *British Poultry Science* 45:573-577.
8. Kapperud, G., Skjerve, E., Vik, L., Hauge, K., Lysaker, A., Aalmen, I., and Potter, M. (1993). Epidemiological investigation of risk factors for *Campylobacter* colonization in Norwegian broiler flocks. *Epidemiology and Infection*, 111(2), 245-256.
9. Rose, N., Beaudeau, F., Drouin, P., Toux, J.Y., Rose, V., and Colin, P. 1999. Risk factors for *Salmonella enterica* subsp. *enterica* contamination in French broiler-chicken flocks at the end of the rearing period, *Preventive Veterinary Medicine* 39:265-277.
10. Martrenchar, A., Boilletot, E., Huonnic, D., and Pol, F., Risk factors for foot-pad dermatitis in chicken and turkey broilers in France, *Preventive Veterinary Medicine* 52:213-226
11. Bowling, A. 2005. Mode of questionnaire administration can have serious effects on data quality. *J. Public Health (Oxf)* 3:281-291.
12. Borck Høg, B., H. Rosenquist, A. I. V. Sorensen, L. S. Larsen, J. Osek, K. Wieczorek, P. Kusyk, M. Cerda-Cuellar, R. Dolz, S. Urdaneta, B. David, M. Hofshagen, J. A. Wagenaar, N. Bolder, F. Jorgensen, N. Williams, Y. Merga, and T. Humphrey. 2011. Questionnaire survey among broiler producers in six European countries. 7th Framework EU-project Camcon Deliverable 1.1.2, Tech. Univ. Denmark, Soborg Denmark.

13. Van Limbergen, T., Dewulf, J., Klinkenberg, M., Ducatelle, R., Gelaude, P., Méndez, J., Heinola, K., Papasolomontos, S., Szeleszczuk, P., and D. Maes. On behalf of the PROHEALTH consortium. 2018. Scoring biosecurity in European conventional broiler production. *Sci.* 97:74-83.
14. Xin, H., I. L. Berry, T. L. Barton, and G. T. Tabler. 1994. Feed and water consumption, growth, and mortality of male broilers. *J. Poult. Sci.* 73:610-616.
15. Tabler, G. T., and I. L. Berry. 2001. Applied Broiler Research Unit Report: Ten-year summary of broiler production results. Ark. Farm Bureau Young Farmers and Ranchers Conference, Hot Springs, AR. Aug 3-4.
16. Morrow C. 2008. Management as a cause of disease in poultry. M. Patisson, P. F. McMullin, J. M. Bradburry, and D. J. Alexander, ed. Pages 536–542 in *Poultry Diseases*. 6th ed. Saunders Elsevier, Beijing, China.
17. 2010. Ross environmental management in the broiler house. Aviagen Ltd., Newbridge, Midlothian, Scotland.
18. Watkins, S. E. 2003. Animal welfare audits: What to expect and how to be prepared. *Avian Advice* 5(4):6-8.
19. Tablante, N.L., Myint, M.S., Johnson, Y.J., Rhodes, K., Colby, M., Hohenhaus, G. 2002. A survey of biosecurity practices as risk factors affecting broiler performance on the Delmarva Peninsula. *Avian Dis.* 46:730-734.
20. Prayitno D.S., Phillips C.J. and Omed H. 1997. The effects of color of lighting on the behavior and production of meat chickens. *Poultry Science* 76:452-457.
21. Houldcroft E., Smith C., Mrowicki R., Headland L., Grieveson S., Jones T.A. and M.S. Dawkins. 2008. Welfare implications of nipple drinkers for broiler chickens. *Animal Welfare* 17:1-10.
22. McMasters J.D., Harris G.C. and T.L. Goodwin. 1971. Effects of Nipple and Trough

- Watering Systems on Broiler Performance. *Poultry Science* 50: 432-435.
23. Lott B.D., May J.D., Simmons J.D. and S.L. Branton. 2001. The effect of nipple height on broiler performance. *Poultry Science* 80: 408-410.
 24. Julian R.J. 1998. Rapid growth problems: ascites and skeletal deformities in broilers. *Poultry Science* 77:1773-1780.
 25. Nijdam E., Arens P., Lambooij E., Decuypere E. and J.A. Stegeman. Factors influencing bruises and mortality of broilers during catching, transport, and lairage. *Poultry Science* 83:1610-1615.
 26. Warris P.D., Bevis E.A., Brown S.N. and J.E. Edwards. 1992. Longer journeys to processing plants are associated with higher mortality in broiler chickens. *British Poultry Science* 33:201-206.
 27. Nijdam E., Zailan A.R., van Eck J.H., Decuypere E. and J.A. Stegeman. Pathological features in dead on arrival broilers with special reference to heart disorders. *Poultry Science* 85:1303-1308.
 28. Vecerek V., Grbalova S., Voslarova E., Janackova B. and M. Malena. 2006. Effects of Travel Distance and the Season of the Year on Death Rates of Broilers Transported to Poultry Processing Plants. *Poultry Science* 85:1881-1884.
 29. Martland M.F. 1985. Ulcerative dermatitis dm broiler chickens: The effects of wet litter. *Avian Pathology* 14:353-364.
 30. Lupo C., Chauvin C., Balaine L., Petetin I., Péraste J., Colin P. and S. Le Bouquin. Postmortem condemnations of processed broiler chickens in western France. *Veterinary Record* 162:709-713.
 31. Northcutt J.K., Savage S.I. and L.R. Vest. 1997. Relationship Between Feed Withdrawal and Viscera Condition of Broilers. *Poultry Science* 76:410-414.
 32. Lyon, C. E., C. M. Papa, and R. L. Wilson, Jr., 1991. Effect of feed withdrawal on

- yields, muscle pH, and texture of broiler breast meat. *Poultry Sci.* 70:1020-1025.
33. Boon C.R. and V.A. Battams. 1988. Air mixing fans in a broiler building —Their use and efficiency. *Journal of Agricultural Engineering Research* 39:137-147.
 34. Lott B.D., Simmons J.D. and May J.D. 1998. Air velocity and high temperature effects on broiler performance. *Poultry Science* 77:391-393.
 35. Dho-Moulin M. and J.M. Fairbrother. 1999. Avian pathogenic *Escherichia coli* (APEC). *Vet. Res.* 30:299-316.
 36. Plavnik I., Macovsky B. and Sklan D. 2002. Effect of feeding whole wheat on performance of broiler chickens. *Animal Feed Science and Technology.* 96:229-236.
 37. Gabriel I., Mallet S., Leconte M., Travel A. and Lalles J.P. 2008. Effects of whole wheat feeding on the development of the digestive tract of broiler chickens. *Animal Feed Science and Technology.* 142:144-162.
 38. Downs K.M., Lien R.J., Hess J.B., Bilgili S.F. and Dozier W.A. III. 2006. The Effects of Photoperiod Length, Light Intensity, and Feed Energy on Growth Responses and Meat Yield of Broilers. *The Journal of Applied Poultry Research* 15:406-416.
 39. Decuypere E., Buyse J. and Buys N. 2000. Ascites in broiler chickens: exogenous and endogenous structural and functional causal factors. *World's Poultry Science Journal* 56:367-377.
 40. Hofacre C.L., Beacorn T., Collett S., and G. Mathis. 2003. Using Competitive Exclusion, Mannan-Oligosaccharide and Other Intestinal Products to Control Necrotic Enteritis. *J.Appl.Poult.Res.* 12:60-64.
 41. Skinner J.T., Bauer S., Young V., Pauling G. and Wilson J. 2010. An Economic Analysis of the Impact of Subclinical (Mild) Necrotic Enteritis in Broiler Chickens. *Avian Diseases* 54:1237-1240.
 42. Timbermont L., Haesebrouck F., Ducatelle R., and F. Van Immerseel. 2011. Necrotic

- enteritis in broilers: an updated review on the pathogenesis. *Avian Pathology* 40:341-347.
43. Maiorka, A., E. Santin, F. Dahlke, I. Boleli, R. Furlan and M. Macari. 2003. Post hatching water and feed deprivation affect the gastrointestinal tract and intestinal mucosa development of broiler chicks. *J. Appl. Poult. Res.* 12: 483-494.
 44. Chapman H.D., Cherry T.E., Danforth H.D., Richards G., Shirley M.W. and R.B. William. 2002. Sustainable coccidiosis control in poultry production: the role of live vaccines. *International Journal for Parasitology* 32:617-629.
 45. Yegani M., and D.R. Korver. 2008. Factors Affecting Intestinal Health in Poultry. *Poultry Science* 87:2052-2063.
 46. Jackson S., Summers J.D., and S. Leeson. 1982. Effect of Dietary Protein and Energy on Broiler Performance and Production Costs. *Poultry Science* 61:2232-2240.
 47. Lilly K.G.S., Gehring C.K., Beaman K.R., Turk P.J., Sperow M., and J.S. Moritz. 2011. Examining the relationships between pellet quality, broiler performance, and bird sex. *Journal of Applied Poultry Research* 20:231-239.
 48. Williams R.B. 2005. Intercurrent coccidiosis and necrotic enteritis of chickens: rational, integrated disease management by maintenance of gut integrity. *Avian Pathology* 34:159-180.
 49. Devettere, R.G.R.G. 2009. Practical decision making in health care ethics: Cases and concepts Georgetown University, Washington D.C.
 50. Scholten, M.C.Th., de Boer, I.J.M., Gremmen, B., Lokhorst, C. 2013. Livestock Farming with Care: towards sustainable production of animal-source food- *Wageningen Journal of Life Sciences* 66:3-5.
 51. Frost, A. R., Schofield, C. P., Beulah, S. A., Mottram, T. T., Lines, J. A. and Wathes, C. M. 1997 A review of livestock monitoring and the need for integrated systems.

Computers and Electronics in Agriculture 17, 139-159.

52. Wathes, C. M., Kristensen, H. H., Aerts, J.-M. and Berckmans, D. 2005 Is precision livestock farming an engineer's daydream or nightmare, an animal's friend or foe, and a farmer's panacea or pitfall? In Second European Conference on Precision Livestock Farming (ed. C. S. Cox), pp. 33-46. Uppsala, Sweden: Wageningen Academic Press, The Netherlands.
53. Dohoo, I., W. Martin, and H. Stryhn. 2009. Veterinary epidemiological research. Charlottetown, PE, Canada.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

Supplementary material.XLSX