

# Prevalence and Risk Factors of Central Obesity Among Adults With Normal Bmi in Shaanxi, China: A Cross-sectional Study

Qiang Li (✉ [tjlq@mail.xjtu.edu.cn](mailto:tjlq@mail.xjtu.edu.cn))

Xi'an Jiaotong University <https://orcid.org/0000-0002-1335-6909>

Wen-yu Feng

Xi'an Jiaotong University

Xiang-dong Li

Xi'an Jiaotong University

Juan Li

Xi'an Jiaotong University

Yuan Shen

Xi'an Jiaotong University

Ling-xia Zeng

Xi'an Jiaotong University

Shao-nong Dang

Xi'an Jiaotong University

---

## Research article

**Keywords:** Central obesity, Body mass index, Prevalence, Risk factor, Metabolic disease

**Posted Date:** May 27th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-533401/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** Defined by normal body mass index (BMI) and higher waist-to-height ratio, normal weight with central obesity (NWCO) has been recognised as a risk factor for cardio-metabolic diseases. The study aims to estimate the prevalence of NWCO and to examine the relationship between NWCO and cardiovascular disease risk factors in adults of Shaanxi Province.

**Methods:** A population-based cross-sectional survey was conducted among residents who were aged 18–80 years and had been living in Zhenba County, Shaanxi Province, for over six months in 2018. The multistage stratified cluster sampling method was used to enroll the study participants. Six towns (districts) in Zhenba County, Shaanxi Province, were randomly selected, and then six villages (communities) were randomly selected from each town (district). All residents in the villages (communities) were face-to-face interviewed. Descriptive data analysis and 95% confidence intervals (CIs) of prevalence/frequency were conducted. Logistic regression analyses were used to detect the corresponding factors associated with central obesity.

**Results:** A total of 2312 participants (936 men and 1376 women) were analyzed. The prevalence of NWCO was 58.3%. NWCO was significantly associated with hypertension and dyslipidemia. Compared with normal weight non-central obesity (NWNCO), the adjusted odds ratios (ORs) for hypertension were 1.47, (95% CI 1.10–1.98) in men and 1.55(1.14–2.10) in women, and the corresponding ones for dyslipidemia were 2.71 (1.77–4.13) in men and 1.84 (1.29–2.61) in women. Female sex, age over 58 years, and lower education level were also significantly predictors of abdominal obesity. In males, comparison with no cigarette smoking, daily cigarette smoking was less likely to lead to normal weight central obesity.

**Conclusion:** In people with normal BMI, the prevalence of central obesity was 58.3%. Female, age over 58 years, hypertension and dyslipidemia indicate a higher risk. Body mass index alone as a measure of obesity is not enough to assess health risks. Central obesity index should be used together for clinical assessment.

## Background

Obesity is a medical condition characterised by an abnormal fat accumulation, which may have adverse effects on health, thereby reducing the life expectancy[1]. Excess body weight is the sixth most important risk factor contributing to the overall burden of disease worldwide[2]. In 2016, more than 1.9 billion adults aged 18 years and older were overweight. Of these over 650 million adults were obese[3]. Overweight and obesity have become a major public health problem in China[4, 5]. A recent study showed that abdominal obesity is associated with insulin resistance and higher risks of metabolic syndrome and cardiovascular diseases in Asian females, whereas general obesity is not[6]. Central obesity has been recognised as an independent risk factor for cardio-metabolic diseases and a better predictor of cardiovascular risk than overall obesity[7, 8]. Defined by normal body mass index (BMI) and higher waist-to-height ratio, normal

weight with central obesity (NWCO) has been recognised as a risk factor for cardio-metabolic diseases[9]. The most common form of obesity is usually assessed by BMI[10].

The main limitation of BMI is that it cannot differentiate body fat from lean mass, and central from peripheral fat. Therefore, athletes with enhanced body muscle mass may be misclassified as obese when using only BMI to diagnose obesity, whereas people with low lean mass but high body fat content may still have a normal BMI[11]. By contrast, proxy measures of anthropometric indicators such as waist circumference (WC), waist-to-hip ratio (WHR) and waist-to-height ratio (WHtR) are used to assess central fat distribution[12]. WHtR is a better predictor than BMI and WC for diabetes, dyslipidemia, hypertension, and cardiovascular disease(CVD)[13]. WHtR may be a simpler and better predictor of early health risks[14–17].

The study aims to estimate the prevalence of normal weight central obesity classified using BMI and WHtR and to examine the relationship between normal weight with central obesity and CVD risk factors in adults in Shaanxi, China.

## Methods

### Subjects

A population-based cross-sectional survey was conducted among residents who were 18–80 years old and had been living in Zhenba County, Shaanxi Province, for over six months in 2018. The multistage stratified cluster sampling method was used to enroll the study participants. Six towns (districts) in Zhenba County, Shaanxi Province, were randomly selected, and then six villages (communities) were randomly selected from each town (district). All residents in the villages (communities) were invited to join this survey and 3781 subjects aged over 18 years were recruited. A total of 2312 normal BMI participants were chosen for the study.

### Data Collection

Data were obtained by face-to-face interviews composed of three parts: questionnaire investigation (included socio-demographic characteristics and health-related information), body measurements (such as height, weight, WC and blood pressure), and laboratory measurements (such as serum cholesterol and triglyceride). Demographic variables including sex, age, level of education, cigarette smoking, coffee consumption, tea intake and alcohol drinking were assessed by self-reporting on the use of any tobacco product or alcohol drinking in the past years. All investigators were trained by research assistants to collect the data through the same questionnaire instruction.

### Measurements

Height and weight were measured in increments of 0.1 cm and 0.1 kg, respectively, by trained staff. BMI was calculated as the weight (kg) divided by the squared height ( $\text{m}^2$ ). WC was measured to the nearest 0.1 cm at the umbilical level in a standing position. WHtR was calculated as WC divided by height. Omron automated blood pressure machine was used to determine the subjects' blood pressure on the right arm, after at least 5 minutes of seated rest. Blood pressure was measured two times with intervals of at least one minute. The average value was used for data analysis. Fasting venous blood samples were drawn from the study subjects to measure serum levels of high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), triglycerides, and blood glucose. The samples were stored in a cooler at 4°C for transportation to the Department of Laboratory Medicine, The First People's Hospital of Zhenba County, and measured within 24 h of being drawn.

## Definition Of Related Indicators

BMI was categorised in accordance with the WHO[18] as underweight ( $< 18.5 \text{ kg/m}^2$ ), normal ( $18.5\text{--}24.9 \text{ kg/m}^2$ ), overweight ( $25.0\text{--}29.9 \text{ kg/m}^2$ ) and obese ( $> 30.0 \text{ kg/m}^2$ ). Central obesity was defined according to the WHO criteria[19, 20]: WHtR  $< 0.5$  (non-central obesity) and  $\geq 0.5$  (central obesity). Normal-weight central obesity was defined as central obesity in participants with normal weight (by BMI). Underweight and overweight subjects ( $\text{BMI} < 18.5$  and  $\text{BMI} \geq 25.0 \text{ kg/m}^2$ ) were excluded from the study.

Self-reported hypertension and/or abnormal blood pressure (systolic  $\geq 140\text{mmHg}$  or diastolic  $\geq 90\text{mmHg}$ ) were regarded as hypertension [21]. The patient who had a history of hypertension and was currently taking medication for hypertension was also defined as having hypertension. Self-reported diabetes mellitus (DM) and/or a fasting serum glucose level  $\geq 7.0\text{mmol/L}$  were regarded as DM[22]. The patient who had a history of diabetes and was currently taking medication for diabetes was also defined as having diabetes. According to the criteria of the "Chinese Guidelines on Prevention and Treatment of Dyslipidemia in Adults", hyperlipidemia diagnosed by a physician and/or abnormal blood lipids ( $\text{TC} \geq 6.2\text{mmol/L}$  or  $\text{TG} \geq 2.3\text{mmol/L}$  or  $\text{HDL-C} < 1.0\text{mmol/L}$ ) were defined as hyperlipidemia[23].

## Statistical Methods

Data were expressed as mean values  $\pm$  standard deviations (SD) for continuous variables. Counts (frequencies = n) and proportions (%) were reported for categorical variables. Chi-squared test and Fisher's exact probability method were used to compare categorical variables. The significant variables were included in the binary logistic regression and adjusted for confounding factors. Analysis was carried out at a 95% confidence level. Data were analyzed by the complex sampling function of SPSS 23.0, and p value of  $< 0.05$  was considered statistically significant.

## Results

In the study, we analyzed a total of 2312 participants (936 men and 1376 women). The mean age of the participants was 58.2 (SD  $\pm$  13.1) years. The majority of the participants were elder people and females, and 25.9% received senior middle school education or higher. Of the 2312 participants classified as normal weight using BMI, the prevalence of central obesity was 58.3% by WHtR. The characteristics of the two groups classified by WHtR (NWN0, NWCO) are shown in Table 1. Among people with normal body mass index, the average age of NWCO group and NWN0 group was 60.5 years and 55.1 years, respectively. The older the age, the higher the prevalence of central obesity. The prevalence in NWCO was 50.3% in men and 63.7% in women, respectively. The prevalence of hypertension, dyslipidemia, and diabetes was 31.5%, 19.0% and 7.1% in NWCO and 20.4%, 9.2% and 5.1% in NWN0, respectively. Central obesity was significantly associated with hypertension and dyslipidemia, but not significant for DM among participants with normal weight, measured by BMI. Education, cigarette smoking and coffee consumption were significantly associated with central obesity using WHtR. There was no significant correlation between central obesity and tea intake or alcohol drinking (Table 1).

Table 1  
Univariate analysis of associated factors of normal weight central obesity

	NWCO	NWNO	$t/\chi^2$	$P$
Age	60.46±12.05	55.09±13.88	-8.971	< 0.001
Sex				
Male	471(35.0)	465(48.2)	40.780	< 0.001
Female	876(65.0)	500(51.8)		
Hypertension				
Yes	424(31.5)	197(20.4)	35.025	< 0.001
No	923(68.5)	768(79.6)		
Dyslipidemia				
Yes	256(19.0)	89 (9.2)	42.379	< 0.001
No	1091(81.0)	876(90.8)		
Diabetes				
Yes	96(7.1)	49 (5.1)	4.016	0.045
No	1251(92.9)	916(94.9)		
Tea intake				
Yes	725(53.8)	529(54.8)	0.224	0.636
No	622(46.2)	436(45.2)		
Coffee consumption				
Yes	26 (1.9)	32 (3.3)	4.415	0.036
No	1321(98.1)	933(96.7)		
Alcohol drinking				
Yes	266(19.7)	203(21.0)	0.577	0.447
No	1081(80.3)	762(79.0)		
Cigarette smoking				
None	1092(81.1)	677(70.2)	50.871	< 0.001
Few times	43 (3.2)	24 (2.5)		
Sometimes	10 (0.7)	4 (0.4)		

	NWCO	NWNO	<i>t/χ<sup>2</sup></i>	<i>P</i>
Daily	202(15.0)	260(26.9)		
Education				
Primary school and below	584(43.4)	298(30.9)	47.496	< 0.001
Junior middle school	461(34.2)	371(38.4)		
Senior middle school	201(14.9)	210(21.8)		
College	84(6.2)	59(6.1)		
High than college	17(1.3)	27(2.8)		
Data are expressed as mean ± standard deviations (SD) or n(%)				
NWNO: normal weight non-central obesity; NWCO: normal weight central obesity				

However, after adjusting for confounding factors (diabetes, alcohol drinking, tea intake and coffee consumption), only female, age over 58 years, cigarette smoking, education, hypertension and dyslipidemia were significant associated with central obesity among the study participants. Female participants were 1.66 times more likely to be central obese than their male counterparts (Table 2).

Table 2  
Logistic regression analysis of associated factors of normal weight central obesity

Variables	$\beta$	Wald $\chi^2$	P	OR(95%CL)
Age $\geq$ 58	0.52	26.39	< 0.001	1.69(1.38,2.06)
Sex = Female	0.46	15.43	< 0.001	1.59(1.26,2.00)
Hypertension	0.43	15.71	< 0.001	1.53(1.24,1.89)
Dyslipidemia	0.79	32.48	< 0.001	2.19(1.68,2.88)
Cigarette smoking		27.65	< 0.001	
None				
Few times	0.22	0.61	0.433	1.24(0.72,2.13)
Sometimes	0.44	0.52	0.473	1.56(0.46,5.24)
Daily	-0.67	23.10	< 0.001	0.51(0.39,0.67)
Education		10.37	0.035	
Primary school and below				
Junior middle school	-0.18	2.91	0.088	0.83(0.68,1.03)
Senior middle school	-0.34	6.31	0.012	0.71(0.55,0.93)
College	0.03	0.02	0.884	1.03(0.70,1.52)
High than college	-0.68	4.09	0.043	0.51(0.26,0.98)
Adjusted for diabetes, tea intake, alcohol drinking and coffee consumption				
OR, odds ratio; 95% CI, 95% confidence interval				

Logistic regression analysis was conducted to calculate the ORs (95% CIs) for hypertension, dyslipidemia and other factors in each gender (Table 3 for men and Table 4 for women). When compared with NWN0, the adjusted ORs for hypertension (adjusted OR 1.47, 95% CI 1.10–1.98 in men; 1.55, 1.14–2.10 in women) and dyslipidemia (2.71, 1.77–4.13 in men; 1.84, 1.29–2.61 in women) had statistically significant association in NWCO. In comparison with no cigarette smoking, daily cigarette smoking was associated with lower risk of NWCO. Adjusted OR for NWCO was 0.52 in males, 95% CI 0.40–0.68,  $p < 0.001$ . Among females there was no statistically significant association with NWCO for either cigarette smoking or not.



Table 3  
Logistic regression analysis of associated factors of normal weight central obesity in males

Variables	$\beta$	Wald $\chi^2$	<i>P</i>	OR(95%CL)
Age $\geq$ 58	0.58	13.13	< 0.001	1.78(1.30,2.44)
Hypertension	0.39	6.56	0.010	1.47(1.10,1.98)
Dyslipidemia	1.00	21.25	< 0.001	2.71(1.77,4.13)
Cigarette smoking		21.70	< 0.001	
None				
Few times	0.19	0.33	0.564	1.21(0.64,2.29)
Sometimes	0.24	0.15	0.704	1.28(0.36,4.49)
Daily	-0.64	18.28	< 0.001	0.53(0.39,0.71)
Education		9.84	0.043	
Primary school and below				
Junior middle school	-0.02	0.01	0.915	0.98(0.70,1.37)
Senior middle school	-0.15	0.58	0.448	0.86(0.58,1.27)
College	0.76	6.42	0.011	2.14(1.19,3.86)
High than college	-0.24	0.25	0.619	0.79(0.30,2.04)
Adjusted for diabetes, tea intake, alcohol drinking, and coffee consumption OR, odds ratio; 95% CI, 95% confidence interval				

Table 4  
Logistic regression analysis of associated factors of normal weight central obesity in females

Variables	$\beta$	Wald $\chi^2$	P	OR(95%CL)
Age $\geq$ 58	0.49	12.47	< 0.001	1.62(1.24,2.13)
Hypertension	0.44	7.87	0.005	1.55(1.14,2.10)
Dyslipidemia	0.61	11.37	0.001	1.84(1.29,2.61)
Education		11.64	0.020	
Primary school and below				
Junior middle school	-0.28	3.83	0.050	0.76(0.58,1.00)
Senior middle school	-0.45	5.86	0.015	0.64(0.44,0.92)
College and above	-0.59	4.59	0.032	0.56(0.33,0.95)
High than college	-1.09	5.13	0.023	0.34(0.13,0.86)
Cigarette smoking		4.08	0.253	
Adjusted for diabetes, tea intake, alcohol drinking and coffee consumption				
OR, odds ratio; 95% CI, 95% confidence interval				

## Discussion

In our study, the prevalence of NWCO defined using a combination of BMI and WHtR was 58.3%. As people get older, the risk of central obesity increases gradually. The prevalence of NWCO was higher in women than in men (63.7% vs. 50.3%). Normal weight with central obesity was associated with CVD risks such as hypertension and dyslipidemia in the present study, regardless of sex.

The higher prevalence of normal-weight central obesity found in this study is suggestive of the need to include anthropometric indices in the measurement of excessive body weight, other than BMI alone, as BMI alone might result in misclassifications and underestimation of at-risk individuals. This being the case, it is important to conduct screening for NWCO using a combination of BMI and WHtR[24]. Such individuals would normally need to be offered the appropriate health education and prompt intervention to manage and/or prevent the development of cardio-metabolic complications. At the same time, it also provides a new perspective for prevention of metabolic diseases: Pay attention to your WHtR even when you have a normal BMI.

Only female sex, age over 58 years, cigarette smoking, education, hyperlipidemia and hypertension were associated with central obesity. The association of central obesity with age and sex has been documented by several studies[25]. The higher likelihood of being obese among females can be linked to

women's lower engagement in physical activity as well as the physiological changes which occur during their reproductive years[26, 27]. Besides, the distribution of fat in the body also changes after middle age: Body fat is more likely to accumulate in the abdomen[28]. This could also be a plausible reason for the higher prevalence of obesity among older participants. An inverse association was found between cigarette smoking and central obesity. In comparison with no cigarette smoking, daily cigarette smoking had an inverse association with the risk of NWCO. This is consistent with previous studies[29, 30]. In addition, the cumulative effect of cigarette smoking on health is detrimental, irrespective of its effect on weight loss. The majority of the study participants had a low level of education. There was significant correlation between central obesity and educational background. Central obesity has been shown to be associated with low education and socioeconomic status, which are characterised by poor health behaviour[31, 32].

Hypertension and hyperlipidemia had a positive association with central obesity, which is consistent with previous studies[33]. People with normal BMI but increased WC have a higher risk of developing metabolic diseases. Moreover, a study has also shown that measures of central obesity are more strongly associated with total and cardiovascular disease death than BMI [34]. Central obesity was found to be strongly related to certain metabolic risk factors among non-obese subjects[35]. In our study, although the prevalence of diabetes was higher in obese patients with normal BMI, no significant correlation between central obesity and hyperglycemia was detected. Our study demonstrates that normal-weight central obesity is associated with hypertension and hyperlipidemia. A previous study noted that central obesity among normal-weight individuals has been shown to be associated with a greater cardiovascular risk and mortality than is found among normal weight individuals without central obesity[34]. Thus, blood glucose, blood lipid and blood pressure should be early detected among adults with normal BMI in order to reduce the prevalence of hypertension, diabetes and hyperlipidemia. Measuring waist circumference should be recommended as a simple and efficient tool for screening central obesity and related metabolic risks even in non-obese individuals. This highlights the need to include measures of central obesity in all clinical assessments in this study setting. Measures aimed at reducing obesity should be prioritised to curb the growing prevalence of chronic diseases, which impacts greatly the already over-burdened healthcare system.

The study still has some limitations. First, the cross-sectional nature of this study and the use of self-reported lifestyle behaviour inevitably produced recall bias. Second, the participants were recruited from Zhenba County in Shaanxi Province, so the conclusion cannot represent the situation in other regions of China. Thus, further studies, including prospective studies, will be needed to establish causality. Last, the sample size of this study was limited and more confounding variables could not be adjusted.

## Conclusion

The prevalence of central obesity is high in adults of normal weight. Sex, age, education and cigarette smoking were closely related to central obesity among study participants. The study suggests that using

BMI alone as a measure of obesity is not enough to assess health risks. In clinical practice, WHtR for measuring central obesity should be considered.

## **Abbreviations**

BMI

body mass index; WHtR:waist-to-height ratio; CI:confidence interval; CVD:cardiovascular disease; NWN0:normal weight non-central obesity; NWCO:normal weight central obesity; SBP:systolic blood pressure; DBP:diastolic blood pressure; SD:standard deviation; WC:waist circumference; WHO:World Health Organization.

## **Declarations**

### **Acknowledgements**

We thank all study participants and staff in each of the study sites and communities including Zhenba Provincial Center for Disease Prevention and Control.

### **Authors' contributions**

WYF contributed to the study design and conducted the statistical analysis, manuscript writing. QL was responsible for careful review of the protocol, design, methodology and provided continuous scientific advice for the study. LXZ, SND, YS assisted with the interpretation of the data. WYF, XDL conceived the article. QL, LXZ, SND, YS, XDL, and JL were involved in the manuscript review. All authors read and approved the final manuscript.

### **Availability of data and materials**

The datasets used and/or analyzed during the current study available from the corresponding author on reasonable request.

### **Ethics approval and consent to participate**

We obtained written informed consents of all the participants. The study was conducted by the Declaration of Helsinki, and the protocol was approved by the Ethics Committee of Xi'an Jiaotong University.

### **Consent for publication**

Not applicable.

### **Competing interests**

The authors declare that they have no competing interests.

## Funding

This research is supported by National Key R&D Program of China (Grand No: 2017YFC0907200, 2017YFC0907201, 2017YFC0907202, 2017YFC0907203, 2017YFC0907204 and 2017YFC0907205)

## Author details

<sup>1</sup>Department of Epidemiology and Biostatistics, School of Public Health, Xi'an

Jiaotong University, No. 76 West Yanta Road, Xi'an 710061, Shaanxi, China.

## References

1. Haslam DW, James WP. Obesity. *Lancet*. 2005; 366(9492):1197-1209.
2. McMillen IC, Rattanatrak L, Duffield JA, Morrison JL, MacLaughlin SM, Gentili S, Muhlhausler BS. The early origins of later obesity: pathways and mechanisms. *Adv Exp Med Biol*. 2009; 646(71-81).
3. World Health Organization. Obesity and overweight. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight> Accessed 10 Feb 2021
4. Jaacks LM, Gordon-Larsen P, Mayer-Davis EJ, Adair LS, Popkin B. Age, period and cohort effects on adult body mass index and overweight from 1991 to 2009 in China: the China Health and Nutrition Survey. *Int J Epidemiol*. 2013; 42(3):828-837.
5. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of All-Cause Mortality With Overweight and Obesity Using Standard Body Mass Index Categories A Systematic Review and Meta-analysis. *Jama- J Am Med Assoc*. 2013; 309(1):71-82.
6. Goh VHH, Hart WG. Excess fat in the abdomen but not general obesity is associated with poorer metabolic and cardiovascular health in premenopausal and postmenopausal Asian women. *Maturitas*. 2018; 107(33-38).
7. Cameron AJ, Magliano DJ, Shaw JE, Zimmet PZ, Carstensen B, Alberti KG, Tuomilehto J, Barr EL, Pauvaday VK, Kowlessur S *et al*. The influence of hip circumference on the relationship between abdominal obesity and mortality. *Int J Epidemiol*. 2012; 41(2):484-494.
8. Piche ME, Poirier P, Lemieux I, Despres JP. Overview of Epidemiology and Contribution of Obesity and Body Fat Distribution to Cardiovascular Disease: An Update. *Prog Cardiovasc Dis*. 2018; 61(2):103-113.
9. Thaikruea L, Thammasarot J. Prevalence of normal weight central obesity among Thai healthcare providers and their association with CVD risk: a cross-sectional study. *Sci Rep*. 2016; 6(37100).
10. Adab P, Pallan M, Whincup PH. Is BMI the best measure of obesity? *BMJ*. 2018; k1274.
11. Romero-Corral A, Somers VK, Sierra-Johnson J, Korenfeld Y, Boarin S, Korinek J, Jensen MD, Parati G, Lopez-Jimenez F. Normal weight obesity: a risk factor for cardiometabolic dysregulation and cardiovascular mortality. *Eur Heart J*. 2010; 31(6):737-746.

12. Wiltink J, Michal M, Wild PS, Zwiener I, Blettner M, Munzel T, Schulz A, Kirschner Y, Beutel ME. Associations between depression and different measures of obesity (BMI, WC, WHtR, WHR). *BMC Psychiatry*. 2013; 13(223).
13. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obes Rev*. 2012; 13(3):275-286.
14. Hsieh SD, Muto T. Metabolic syndrome in Japanese men and women with special reference to the anthropometric criteria for the assessment of obesity: Proposal to use the waist-to-height ratio. *Preventive Medicine*. 2006; 42(2):135-139.
15. Tatsumi Y, Watanabe M, Kokubo Y, Nishimura K, Higashiyama A, Okamura T, Okayama A, Miyamoto Y. Effect of Age on the Association Between Waist-to-Height Ratio and Incidence of Cardiovascular Disease: The Suita Study. *J Epidemiol*. 2013; 23(5):351-359.
16. Sahakyan KR, Somers VK, Rodriguez-Escudero JP, Hodge DO, Carter RE, Sochor O, Coutinho T, Jensen MD, Roger VL, Singh P *et al*. Normal-Weight Central Obesity: Implications for Total and Cardiovascular Mortality. *Annals of Internal Medicine*. 2015; 163(11):827.
17. Ashwell M, Gibson S. Waist-to-height ratio as an indicator of 'early health risk': simpler and more predictive than using a 'matrix' based on BMI and waist circumference. *BMJ Open*. 2016; 6(3):e010159.
18. World Health Organization. Obesity and overweight. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight> Accessed 2.10 2021
19. Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev*. 2010; 23(2):247-269.
20. Wakabayashi I. Necessity of Both Waist Circumference and Waist-to-Height Ratio for Better Evaluation of Central Obesity. *Metab Syndr Relat D*. 2013; 11(3):189-194.
21. Liu LS, Writing Group of Chinese Guidelines for the Management of H. [2010 Chinese guidelines for the management of hypertension]. *Zhonghua Xin Xue Guan Bing Za Zhi*. 2011; 39(7):579-615.
22. Weng J, Ji L, Jia W, Lu J, Zhou Z, Zou D, Zhu D, Chen L, Chen L, Guo L *et al*. Standards of care for type 2 diabetes in China. *Diabetes Metab Res Rev*. 2016; 32(5):442-458.
23. Joint committee for guideline r. 2016 Chinese guidelines for the management of dyslipidemia in adults. *J Geriatr Cardiol*. 2018; 15(1):1-29.
24. Mirzaei M, Khajeh M. Comparison of anthropometric indices (body mass index, waist circumference, waist to hip ratio and waist to height ratio) in predicting risk of type II diabetes in the population of Yazd, Iran. *Diabetes Metab Syndr*. 2018; 12(5):677-682.
25. Ford ES, Maynard LM, Li C. Trends in Mean Waist Circumference and Abdominal Obesity Among US Adults, 1999-2012. *JAMA*. 2014; 312(11):1151.
26. Sumithran P, Houlihan C, Shub A, Churilov L, Pritchard N, Price S, Ekinici E, Proietto J, Permezel M. How common is substantial weight gain after pregnancy? *Obesity Research & Clinical Practice*.

2018; 12(2):139-145.

27. Gunderson EP. Childbearing and Obesity in Women: Weight Before, During, and After Pregnancy. *Obstetrics and Gynecology Clinics of North America*. 2009; 36(2):317-332.
28. Tchernof A, Despres JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev*. 2013; 93(1):359-404.
29. Courtemanche C, Tchernis R, Ukert B. The effect of smoking on obesity: Evidence from a randomized trial. *J Health Econ*. 2018; 57(31-44).
30. Mackay DF, Gray L, Pell JP. Impact of smoking and smoking cessation on overweight and obesity: Scotland-wide, cross-sectional study on 40,036 participants. *Bmc Public Health*. 2013; 13(
31. Camoes M, Lopes C, Oliveira A, Santos AC, Barros H. Overall and central obesity incidence in an urban Portuguese population. *Preventive Medicine*. 2010; 50(1-2):50-55.
32. Du TT, Sun XX, Yin P, Huo R, Ni CC, Yu XF. Increasing trends in central obesity among Chinese adults with normal body mass index, 1993-2009. *Bmc Public Health*. 2013; 13(
33. Zhou X, Ji L, Ran X, Su B, Ji Q, Pan C, Weng J, Ma C, Hao C, Zhang D *et al*. Prevalence of Obesity and Its Influence on Achievement of Cardiometabolic Therapeutic Goals in Chinese Type 2 Diabetes Patients: An Analysis of the Nationwide, Cross-Sectional 3B Study. *PLoS One*. 2016; 11(1):e0144179.
34. Sahakyan KR, Somers VK, Rodriguez-Escudero JP, Hodge DO, Carter RE, Sochor O, Coutinho T, Jensen MD, Roger VL, Singh P *et al*. Normal-Weight Central Obesity: Implications for Total and Cardiovascular Mortality. *Ann Intern Med*. 2015; 163(11):827-835.
35. Lukács A, Horváth E, Máté Z, Szabó A, Virág K, Papp M, Sándor J, Ádány R, Paulik E. Abdominal obesity increases metabolic risk factors in non-obese adults: a Hungarian cross-sectional study. *BMC Public Health*. 2019; 19(1):