

Prevalence of people at risk of developing type 2 diabetes mellitus and the involvement of community pharmacies in a national screening campaign: a pioneer action in Brazil

Cassyano Januário Correr

Universidade Federal do Parana

Wendel Coura-Vital (✉ wendelcoura@ufop.edu.br)

Universidade Federal de Ouro Preto <https://orcid.org/0000-0002-1434-7676>

Josélia Cintya Quintao Pena Frade

Conselho Federal de Farmácia

Renata Cristina Rezende Macedo do Nascimento

Universidade Federal de Ouro Preto

Lúbia Guaima Nascimento

Universidade Federal de Ouro Preto

Eliete Bachrany Pinheiro

Conselho Federal de Farmácia

Wesley Magno Ferreira

Conselho Federal de Farmácia

Janice Sepúlveda Reis

Instituto de Ensino e Pesquisa da Santa Casa de Belo Horizonte

Karla Fabiana Santana de Melo

Hospital das Clínicas da Universidade de São Paulo

Roberto Pontarolo

Universidade Federal do Paraná

Mônica Soares Amaral Lenzi

Conselho Federal de Farmácia

José Vanilton de Almeida

Conselho Federal de Farmácia

Hermelinda Cordeiro Pedrosa

Sociedade Brasileira de Diabetes

Walter da Silva Jorge João

Conselho Federal de Farmácia

Research

Keywords: Diabetes Mellitus, Risk Factors, Chronic Disease, prevalence

Posted Date: September 14th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-57968/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 8th, 2020. See the published version at <https://doi.org/10.1186/s13098-020-00593-5>.

Abstract

Background: Brazil is one of top 10 countries with the highest number of people with diabetes mellitus (DM), affecting 16.8 million peoples. It is estimated that 7.7 million people (20–79 years) in the country have not yet been diagnosed, representing an under-diagnosis rate of 46.0%. Herein we aimed to screen people for high blood glucose or risk for developing type 2 DM (T2DM) through community pharmacies in Brazil.

Methods: A cross-sectional study was carried out in November 2018, involving 977 pharmacists from 345 municipalities in Brazil. The study evaluated people between 20 and 79 years old without a previous diagnosis of DM. Glycemia was considered high when its value was ≥ 100 mg/dL fasting and ≥ 140 mg/dL in a casual feeding state. The FINDRISC (Finnish Diabetes Risk Score) was used to estimate the risk for developing T2DM. The prevalence of high blood glucose was estimated and the associated factors were obtained using Poisson's multivariate analysis with robust variance.

Results: During the national screening campaign, 17,580 people were tested with the majority of the consultations (78.2%) being carried out in private pharmacies. The population was composed mainly of women (59.5%) and people aged between 20 and 45 years (47.9%). The frequency of participants with high blood glucose was 18.4% (95% CI 17.9–19.0). Considering the FINDRISC, 22.7% of people had a high or very high risk for T2DM. The risk factors associated with high blood glucose were: Body Mass Index > 25 kg/m², abdominal circumference > 94 cm for men and > 80 cm for women; education level below 15 years of study, no daily intake of vegetables and fruits; previous diagnosis of arterial hypertension; history of high blood glucose and family history of DM.

Conclusions: This is the largest screening study that evaluated the frequency of high blood glucose and its associated factors in a population without a previous diagnosis ever performed in community pharmacies in Brazil. These results may help to improve public health policies and reinforce the role of pharmacists in screening and education actions aimed at this undiagnosed population in a continent-size country such as Brazil.

Background

The epidemiological transition is a major challenge for health systems and societies in the Americas [1]. Population aging, urbanization, and changes in lifestyle have impacted the prevalence of chronic noncommunicable diseases (NCD), with emphasis on hypertension, diabetes mellitus, and dyslipidemia [1, 2].

Brazil is one of the top 10 countries in the world with the highest number of people with DM and currently is the 5th in the ranking [3]. International estimates indicate a growing trend in the prevalence of DM in the country, with an expected increase ranging from 6% to 7.8% in 2030 [4]. Despite the lack of nationwide studies on the prevalence of DM, results from the Vigitel 2017 household survey showed a prevalence of 7.7% in the 27 Brazilian capitals, with 7.1% for men and 8.1% for women [5]. The 2019 9th IDF Diabetes

Atlas estimated that in the Brazilian population between 20 and 79 years old there are approximately 16.8 million people with diabetes in Brazil, and 7.7 million undiagnosed DM, representing an under-diagnosis proportion of 46.0% [3].

Several studies have pointed out impressive data on pre-diabetes in Brazil. The National Health Survey, conducted from 2014-2015, which used glycated hemoglobin (HbA1c) as a diagnostic tool in a sample over 8,500 people, identified a prevalence of pre-diabetes ranging from 6.8% to 16.9%, depending on the criteria used [6]. The ELSA longitudinal study, involving Brazilian adults, reported pre-diabetes prevalence ranging from 20-59% of the sample recruited among university professional staff [7]. In addition to the diagnostic challenges, control rates of DM in Brazil remain unsatisfactory. The last largest nationwide study analyzing glucose control was conducted in 2006 and showed that 75% of 6.671 individuals with T2DM and 90% of those with type 1 DM (T1DM) assisted in private and public services by either specialists or no-specialists presented HbA1c higher than 7% [8]. Other studies have shown similar or even worse results among T1DM [9, 10].

In Brazil, the treatment of diabetes micro and macrovascular complications represents the main component of the disease expenses, with 48.2% of the cost attributed to medicines to treat these complications [11, 12]. Therefore, early diagnosis of T2DM is important to prevent chronic complications associated with the disease and to reduce the costs associated with health care delivery.

The Finnish Diabetes Risk Score (FINDRISC) is a simple, non-invasive and easy-to-use screening instrument used to predict the risk of an adult developing DM2 in 10 years [13]. This instrument has been adopted worldwide to lead T2DM prevention programs in primary health care [14-17] and has been adapted by the Brazilian Diabetes Society to be applied during national awareness campaigns in Brazil [18]. Studies conducted in several countries have shown that pharmacies are successful partners in programs for screening and diagnosing diabetes and in the identification of risk factors for cardiovascular diseases [19-23]. In Brazil, pharmacies are establishments that provide pharmaceutical services as well as individual and collective health guidance, including assessment of capillary blood glucose, blood pressure measurement, among other procedures. Pharmacists are higher education health professionals with responsibilities for carrying out clinical activities, including screening, education, and monitoring of patients [24, 25]. In this context, the present study aimed to identify, among people without a previous diagnosis of DM, the frequency of high blood glucose in the Brazilian population and the associated factors.

Methods

Study design and population

A cross-sectional study was carried out in November 2018, involving 977 pharmacists from 345 municipalities in Brazil, to assess the frequency of high blood glucose in individuals without a previous diagnosis of DM.

Brazil is the largest country in South America, with a territorial area of 8,510,820.623 Km², divided into five geographic regions (North, South, Midwest, Southeast, and Northeast) and subdivided into 27 federative units, including the Federal District. The estimated national population, for the year 2018, was 210,867,954 inhabitants [26]. To calculate the ideal sample size, there were considered the Brazilian population aged between 20 and 79 years (141,802,185 people; IBGE 2017); an estimated prevalence of pre-diabetes or DM of 23% [27]; 95% confidence level; and 1% estimate accuracy. The estimated sample for the country (11,750) was further stratified taking into account the population aged 20 to 79 years in each federative unit.

People aged between 20 and 79 years old who were attended at any of the selected pharmacies, during the study period, were invited by pharmacists to participate in the study. Participants could be in their feeding routine (unknown diet) or fasting (8 hours without any caloric intake). The exclusion criteria were: participants' records presenting blood glucose data collected with glucometers without ISO 15197: 2013 certification (ISO, 2013); previous diagnosis of DM; any data collected with non-standard anthropometric tape; age <20 years or > 79 years; lack of data on weight (measured on a scale), height, Body Mass Index (BMI) and or abdominal circumference (AC); capillary blood glucose (CBG) lower than 70 mg/dL without confirmation or CBG higher than 300 mg/dL without confirmation.

Selection of pharmacies and training of pharmacists

Participated in the study volunteer pharmacists from community pharmacies (private and public), located in different municipalities throughout the national territory. Community pharmacy is a type of health care facility that provides pharmaceutical services or with a given mission around medicines [24, 28].

Pharmacies were selected in a non-probabilistic manner, by fulfilling the following criteria: (i) ability to collect and send data by electronic means; (ii) agreeing to use their inputs (glucometer, blood glucose test strips, scale, anthropometric tape, office supplies); (iii) be located in a municipality that has a public primary health care service and that was covered by the Mobile Emergency Care Service; (iv) be regularly enrolled in the Regional Pharmacy Council of their jurisdiction, with full-time pharmaceutical coverage; (v) have a patient care room allowing visual and sound privacy (pharmaceutical care room).

The recruitment of the the study population was carried out by broadcasting media campaigns (television, radio, Federal Pharmacy Board website [29], and a website developed exclusively for the campaign [30]) and locally, by the participating pharmacies. Standardized disclosure materials were provided to participating pharmacies.

Testing protocol and data collection

During November 2018, people who attended any of the selected pharmacies were invited by the pharmacist to participate in the study, being informed about the objectives of the project and the inclusion criteria. Those who agreed to participate signed the Free and Informed Consent Form, answered a questionnaire with personal information, clinical data and the FINDRISC [13], translated and cross-culturally adapted to Brazilian Portuguese [31]. The FINDRISC analyses eight clinical characteristics: age,

BMI, AC, daily physical activity, eating habits, blood pressure, use of antihypertensive medication, history of hyperglycemia, and family history of diabetes. The final score corresponds to the sum of the scores attributed to each question, ranging from 0 to 26, and the individual risk of developing T2DM was stratified into five categories, ranging from low to very high risk [15]. Subsequently, anthropometric measurements and CBG test were performed. Height was obtained with a tape measure fixed on the wall. Weight was obtained by a digital or analog scale registered by the National Health Surveillance Agency (Anvisa), available at the time of the evaluation. BMI was obtained by dividing the weight (in kilograms) by the square of height (in meters); AC was measured using a professional anthropometric tape, following the recommendations of the Brazilian Association for the Study of Obesity and Metabolic Syndrome [32]; CBG tests were performed using glucometers registered by Anvisa, with ISO 15197: 2013 (ISO, 2013) certification, such as Accu Check Performa® (Roche), Accu Check Performa Connect® (Roche), Accu Check Guide® (Roche), One Touch Select Plus Flex® (J&J), Contour Plus® (Bayer), G-Tech Free Lite (GTech), Freestyle Freedom Lite® (Abbott), and Freestyle Optium Neo® (Abbott). The results were interpreted according to the recommendations of the Brazilian Diabetes Society, considering the patient's feeding status at the time of the CBG test [18]. CBG levels less than 100 mg/dL were considered normal when fasting or less than 140 mg/dL, in a casual feeding state.

The interpretation of the CBG result and the subsequent orientation to the participant were performed by the pharmacist. Participants with results suggestive of pre-diabetes or DM were referred to the local public or private health system, with a proper filled in declaration form [25], containing the result of the parameters evaluated during the screening procedures.

Data analysis

After filling in the survey instruments, the data were entered into an online platform (SurveyMonkey) and later exported for analysis using the Stata Version 14.0 software.

A difference observed in the degrees of risk defined by FINDRISC and CBG levels were evaluated by chi-square test.

The Poisson multivariate model, with robust variance, was used to identify the risk factors associated with high blood glucose and the estimates were obtained using the prevalence ratio (PR) with 95% CI. Initially, a univariate analysis was performed and the variables that showed a statistical association in this analysis ($p < 0.25$) were analyzed in a multivariate model. Variables with more than two categories were transformed into dummy variables. For the construction of the final model, a complete model was started, containing all the variables, and the successive disposal of the variables was carried out until only those with significance level remained in the model ($p < 0.05$).

Results

The final sample screened was 20,171 participants, of which 17,580 (87.2%) were eligible for the study (Figure 1).

Characteristics of the assessed population

Of the 17,580 participants included, most were women (59.5%) and the predominant age group was less than 45 years old (47.9%). The mean and median age were 46 ± 15.7 years, with the first and third quartiles being 33 and 59, respectively. The predominant skin color was white (47.3%), followed by brown (38.0%). In terms of education, 77.7% of the participants had more than eight years of study, 18.0% reported 8 to 10 years of study, 29.1% indicated 11 to 14 years, and 30.6% informed 15 or more years of study. Regarding the practice of physical activities, 68.3% reported not practicing any activity and, concerning eating habits, 43.0% did not eat vegetables or fruits daily. It was observed that 30.8% of the sampled population self-reported hypertension or used antihypertensive medication. As for blood glucose rates, 15.8% reported having high glucose levels events in the past and 58% informed a family history of T1DM or T2DM (Table 1).

Table 1
Baseline characteristics of the population included in the screening of
high blood glucose in Brazil, 2018

Variables	n	%
<i>Gender</i>		
Male	7,112	40.5
Female	10,468	59.5
<i>Age (years)</i>		
20-45	8,412	47.9
45-54	3,374	19.2
55-64	3,188	18.1
65-79	2,606	14.8
<i>Abdominal circumference (cm)</i>		
Male < 94	2,916	41.0
94-102	2,099	29.5
> 102	2,097	29.5
Female < 80	2,222	21.2
80-88	2,184	20.9
> 88	6,062	57.9
<i>Color</i>		
White	8,318	47.3
Brown	6,679	38.0
Black	2,253	12.8
Yellow	192	1.1
Indigenous	52	0.3
Uninformed	86	0.5
<i>Educational level (years)</i>		
Illiterate	814	4.6
1 -3	173	1.0
4 -7	2,936	16.7

Variables	n	%
8 -10	3,166	18.0
11 -14	5,108	29.1
≥ 15	5,383	30.6
<i>Practice of physical activity</i>		
Yes	5,575	31.7
No	12,005	68.3
<i>Consumption of vegetables and/or fruits</i>		
Every day	10,024	57.0
Do not eat every day	7,556	43.0
<i>Diagnosis or use of medicines for hypertension</i>		
Yes	5,413	30.8
No	12,167	69.2
<i>High blood glucose in the past</i>		
Yes	2,780	15.8
No	14,800	84.2
<i>Family member with type 1 or 2 diabetes mellitus</i>		
No	7,377	42.0
Yes (grandparents, uncles, cousins)	3,741	21.3
Yes: parents. siblings or children	6,462	36.7

Risk of developing DM2 in the next 10 years

According to the FINDRISC, 22.1% (n = 3,873) were at low risk for developing T2DM (1 in 100 develops the disease); 35.1% (n = 6,169) slightly moderate risk (1 in 25 develops the disease); 20.1% (n = 3,523) moderate risk (1 in 6 develops the disease); 19.6% (n = 3,436) high risk (1 in 3 develops the disease) and 3.1% (n = 555) very high risk (1 in 2 develops the disease) (Table 2).

Table 2

Risk for developing T2DM (FINDRISC), according to capillary blood glucose levels, regardless of dietary status, Brazil, 2018.

FINDRISC	Blood glucose (mg/dL)*		Total n (%)
	Normal n (%)	High n (%)	
Low risk	3,565 (24.9)	308 (9.5)	3,873 (22.1)
Slightly moderate risk	5,275 (36.8)	894 (27.6)	6,169 (35.1)
Moderate risk	2,828 (19.8)	695 (21.4)	3,523 (20.1)
High risk	2,350 (16.4)	1,086 (33.5)	3,436 (19.6)
Higher risk	297 (2.1)	258 (8.0)	555 (3.1)

* Glycemia considered normal: fasting < 100 mg/dL or casual < 140 mg/dL. Results excluding 24 patients with blood glucose results < 70 mg/dL.

Among the participants with low or moderate risk, 24.9% and 36.8%, respectively, had normal CBG. Participants who presented high or very high FINDRISC were mainly those who had high CBG levels, with percentages much higher than those found in individuals with normal CBG, corresponding to 33.5% and 8.0% respectively.

Frequency of high capillary blood glucose levels

The frequency of high CBG levels (fasting CBG \geq 100mg/dL or in random condition \geq 140 mg/dL) among the individuals enrolled in this present screening was 18.4% (95% CI 17.9–19.0). When the frequency by geographic region was assessed, the Midwest presented the highest frequency (24.6%; 95% CI: 22.4–27.0), followed by the North (22.5%; 95% CI 20.4 –24.8) and Northeast (19.8%; 95% CI: 18.8–20.8) regions (Figure 2).

Risk factors for high capillary blood glucose

Preliminary selection of variables ($p < 0.25$), through univariate analysis, is described in Table 3. According to multivariate analysis (Table 4), the BMI of 25-30 kg/m² (PR 1.1) and BMI > 30 kg/m² (PR 1.3), when compared with BMI < 25 kg/m²; AC for men of 94-102 cm or for women of 80-88 cm (PR 1.2) as well as AC for men > 102 cm or for women > 88 cm (PR 1.3) compared to AC < 94 cm for men or < 80 cm for women, increased the risk of high CBG. People with less education had a higher frequency of high CBG, compared to people with 15 or more years of education (illiterate PR 1.5). People who did not eat fruits or vegetables every day (PR 1.1), reported being hypertensive or use medicines for high blood pressure (PR 1.1), and those informing a high glycemia rate in the past (PR 1.9) and having first-degree

relatives with either T1DM or T2DM (PR 1.3) were associated with a higher prevalence of high CBG. The PR values obtained in the multivariate model were adjusted for the participants' age and sex.

Table 3

Univariate analysis of population characteristics associated with high blood glucose levels, Brazil, 2018

Variables	Blood glucose (mg/dL)*		PR (CI95%)	p-value
	Normal (n/%)	High (n/%)		
<i>Gender</i>				
Male	8,720 (60.9)	1,734 (53.5)		
Female	5,595 (39.1)	1,507 (46.5)	1.3 (1.2–1.4)	0.01
<i>Age (years)</i>				
20–45	7,486 (52.3)	913 (29.2)		
45–54	2,655 (18.5)	716 (22.1)	1.9 (1.8–2.1)	0.01
55–64	2,357 (16.5)	827 (25.5)	2.4 (2.2–2.6)	0.01
65–79	1,817 (12.7)	785 (24.2)	2.8 (2.6–3.0)	0.01
<i>Body Mass Index</i>				
<25	5,220 (35.5)	782 (24.1)		
25–30	5,595 (39.1)	1,326 (40.9)	1.5 (1.4–1.6)	0.01
>30	3,500 (24.5)	1,133 (35.0)	1.9 (1.7–2.0)	0.01
<i>Abdominal circumference (cm)</i>				
♂ < 94 or ♀ < 80	4,491 (31.4)	635 (19.6)		
♂ 94–102 or ♀ 80–88	3,518 (24.6)	760 (23.4)	1.4 (1.3–1.6)	0.01
♂ > 102 or ♀ > 88	6,306 (44.0)	1,846 (57.0)	1.8 (1.7–2.0)	0.01
<i>Color</i>				
White	6,821 (47.6)	1,489 (45.9)		
Brown	5,471 (38.2)	1,200 (37.0)	1.0 (0.9–1.1)	0.91
Black	156 (1.1)	36 (1.1)	1.0 (0.8–1.4)	0.77
Yellow	1,754 (12.3)	491 (15.2)	1.2 (1.1–1.3)	0.01
Indigenous	44 (0.3)	8 (0.3)	0.8 (0.4–1.6)	0.64
Uninformed	69 (0.5)	17 (0.5)	1.1 (0.7–1.7)	0.65
<i>Educational level (years)</i>				

* Glycemia considered normal. fasting < 100 mg/dL or casual < 140 mg/dL. Results excluding 24 patients with blood glucose results < 70 mg/dL. ♂ - Male ; ♀ - Female

Variables	Blood glucose (mg/dL)*		PR (CI95%)	p-value
	Normal (n/%)	High (n/%)		
≥ 15	4,644 (32.4)	734 (22.7)		
11 -14	4,347 (30.4)	752 (23.2)	1.1 (1.0-1.2)	0.10
8 -10	2,498 (17.4)	665 (20.5)	1.5 (1.4-1.7)	0.01
4 -7	2,143 (15.0)	791 (24.4)	2.3 (2.1-2.6)	0.01
1 -3	123 (0.9)	49 (1.5)	2.5(1.8-3.5)	0.01
Illiterate	560 (3.9)	250 (7.7)	2.8 (2.4-3.3)	0.01
<i>Practice of physical activity</i>				
Yes	4,608 (32.2)	964 (29.7)		
No	9,707 (67.8)	2,277 (70.3)	1.1 (1.0-1.2)	0.01
<i>Consumption of vegetables and/or fruits</i>				
Every day	8,216 (57.4)	1,796 (55.4)		
Do not eat every day	6,099 (42.6)	1,445 (44.6)	1.1 (1.0-1.1)	0.04
<i>Diagnosis or use of medicines for hypertension</i>				
No	10,362 (72.4)	1,787 (55.1)		
Yes	3,953 (27.6)	1,454 (44.9)	1.8 (1.7-1.9)	0.01
<i>High blood glucose in the past</i>				
No	12,533 (87.5)	2,243 (69.2)		
Yes	1,782 (12.5)	998 (30.8)	2.4 (2.2-2.5)	0.01
<i>Family member with type 1 or 2 diabetes mellitus</i>				
No	6,159 (43.0)	1,204 (37.1)		
Yes (grandparents, uncles, cousins)	3,228 (22.5)	505 (15.6)	0.8 (0.7-0.9)	0.01
Yes: (parents, siblings or children)	4,928 (34.4)	1,532 (47.3)	1.4 (1.3-1.5)	0.01
* Glycemia considered normal. fasting < 100 mg/dL or casual < 140 mg/dL. Results excluding 24 patients with blood glucose results < 70 mg/dL. ♂ - Male ; ♀ - Female				

Table 4

Multivariate analysis of characteristics associated with high blood glucose, Brazil, 2018

Variables	PR (CI95%) crude	PR (CI95%) adjusted*
<i>Body Mass Index</i>		
<25	-	-
25–30	1.5 (1.4–1.6)	1.1 (1.1–1.2)
>30	1.9 (1.7–2.0)	1.3 (1.2–1.4)
<i>Abdominal circumference (cm)</i>		
♂ < 94 or ♀ < 80	-	-
♂ 94–102 or ♀ 80–88	1.4 (1.3–1.6)	1.2 (1.1–1.3)
♂ > 102 or ♀ > 88	1.8 (1.7–2.0)	1.3 (1.2–1.5)
<i>Educational level (years)</i>		
≥ 15	-	-
8 –10	1.5 (1.4–1.7)	1.3 (1.1–1.4)
4 –7	2.3 (2.1–2.6)	1.4 (1.3–1.5)
1 –3	2.5 (1.8–3.5)	1.5 (1.2–1.9)
Illiterate	2.8 (2.4–3.3)	1.5 (1.3–1.7)
<i>Consumption of vegetables and/or fruits</i>		
Every day	-	-
Do not eat every day	1.1 (1.0–1.1)	1.1 (1.1–1.2)
<i>Diagnosis or use of medicines for hypertension</i>		
No	-	-
Yes	1.8 (1.7–1.9)	1.1 (1.1–1.2)
<i>High blood glucose in the past</i>		
No	-	-
Yes	2.4 (2.2–2.5)	1.9 (1.8–2.0)
<i>Family member with type 1 or 2 diabetes mellitus</i>		
No	-	-
Yes (parents, siblings or children)	1.4 (1.3–1.5)	1.3 (1.2–1.4)
♂ - Male ; ♀ - Female. *Final model adjusted for age and sex.		

Discussion

This is the largest study ever conducted in pharmacies in all regions of Brazil that assessed the frequency of high blood glucose levels and the risk of developing diabetes. Performing NCD screening is important because it can identify asymptomatic people at high risk or with initial signs and symptoms of a disease, allowing for timely diagnosis and treatment, which is very important for a disease such as diabetes. According to the WHO, a substantial fraction of the disease burden and mortality due to NCD are related to a small set of risk factors, among which stand out smoking, inadequate food consumption, physical inactivity, and excessive consumption of alcoholic beverages [33]. Early detection of people at risk of developing T2DM is very important, since changes in lifestyle, including physical activity and proper diet, can reduce the incidence of the disease by approximately 58% [14, 18]. In addition, its early detection and intervention in patients with T2DM can minimize complications from other comorbidities such as non-alcoholic fatty liver disease, with insulin resistance being a common pathophysiological mechanism that mutually interferes in the progression of both diseases [34, 35].

It was found that the frequency of high CBG levels was higher in the Midwest, North, and Northeast regions. These findings can be explained by the heterogeneous profile of the Brazilian population, related to lifestyle and prevalence of obesity which has been increasing. In Brazil, overweight (BMI > 25 kg/m²) and obesity rates are spreading in all age groups, both sexes, and all income levels, with the obesity rates being more frequent in the population with lower family income [36]. Indeed, according to data from Vigitel 2018 [37], the prevalence of obesity (BMI > 30 kg/m²) in adults is 19.8%, with the highest frequencies observed in capitals of the Midwest, North, and Northeast regions, with emphasis on the capital cities like Cuiabá (23.0% CI 95% 20.5-25.4), Manaus (23.0% CI 95% 19.7-26.3), and Recife (21.9% CI 95% 19.3- 24.4).

Regarding the consumption of fruits and vegetables, a similar distribution was found, with the lowest percentages observed in the North and Northeast regions [37]. The Vigitel survey also pointed out that consumption of fruits and vegetables is more frequent among women and, despite the increase in the consumption of these foods observed in the last ten years, only 23.1% of Brazilians consume the five daily portions at least five times a week, as recommended by the WHO [37]. Generally, diets have a high sugar intake, acting as the main source of advanced glycation endproducts (AGEs). These compounds are toxic and are associated with the pathogenesis of diet-related diseases, such as diabetes, insulin resistance, among others [38]. The AGEs, due to hyperglycemia and insulin resistance present in people with diabetes, are an extra source of reactive oxygen species (ROS) which overload the cellular antioxidant machinery, leading the cell to a state of oxidative stress [39]. Thus, natural compounds with antioxidant properties, such as citrus fruits, tomatoes, green tea, can prevent the harmful effects of AGEs, providing benefits in the control of T2DM. [38, 40]. In turn, ROS mediate chronic inflammatory processes that are directly related to the micro and macrovascular complications of diabetes, which are diabetic retinopathy, neuropathy and nephropathy [41, 42]. Inflammatory processes can also be triggered by diets with low intake of vitamin D, a fact corroborated by study that show an inverse correlation between blood levels of vitamin D and C-reactive protein. This relationship being more pronounced in patients with

inflammatory diseases, such as diabetes, compared to patients with non-inflammatory diseases [43]. In addition, it has been speculated that vitamin D supplementation may contribute to reducing the risk of developing diabetes in patients at risk of developing the disease [44].

Similar to the results reported by Passos et al. (2005), it was verified a positive relationship between physical inactivity and high CBG [45]. Sedentary lifestyle has been associated with insulin resistance in individuals without a diagnosis of diabetes, regardless of obesity [46, 47]. Reporting of previous arterial hypertension was also associated with high CBG. It is known that hypertension has a prevalence up to three times higher in people with DM when compared to those who do not have the disease, being the major determinant of atherosclerotic cardiovascular diseases (ASCVD) in this population [48, 49].

The percentage of people at high and very high risk of developing T2DM in Brazil was similar to the screening performed in pharmacies in Spain (23.5% had a high risk of diabetes -FINDRISC > 14 points) [50] and smaller than the risk reported by the screening performed in Thailand, in seven pharmacies, which showed a prevalence of 48% of high risk for DM [51]. The frequency of high CBG in Brazil was higher than the prevalence found in other countries such as Spain, Switzerland, and Thailand [50-52], indicating the need for multidisciplinary monitoring of these people.

T2DM is characterized by a long asymptomatic period before the diagnosis [14]. The early diagnosis of T2DM contributes to the reduction of ASCVD, prevention of microvascular complications (retinopathy, neuropathy, kidney disease), and premature mortality [14, 47, 53, 54]. People with T2DM have a two to four times greater risk of developing coronary heart disease, when compared to the general population, and approximately 8% of people with pre-diabetes develop retinopathy [47]. Moreover, more recent data have shown high percentages of peripheral neuropathy among people with obesity and pre-DM (29% versus 11% with normal glucose), and ranges from 11 to 34% among individuals with pre-DM [55]. Thus, early interventions to prevent or delay progression to T2DM represent an important benefit not only for increasing life expectancy and quality of life, but also reducing costs related to the management of the disease and its complications [15].

Success in the treatment of DM depends on the concomitant implementation of three categories of interventions: educational, self-monitoring, and pharmacological strategies. Whenever possible, it is recommended that care for people with diabetes should include a health care professional (HCP) interdisciplinary team composed of professionals with the proper qualification and practical experience in health education activities [56]. Pharmacists working in community pharmacies can contribute not only with HCP screening and education, but also with the monitoring of patients with DM, in collaboration with other HCP thus helping to improve the treatment and disease control.

A systematic review and meta-analysis assessing the impact of pharmaceutical care for people with diabetes, in outpatient services, revealed a 1.1% reduction in HbA1c compared to standard care (95% CI 0.88-1.27) [57]. In Brazil, a 12-months study enrolling patients with DM, followed in a pharmacy, showed a reduction of 2.2% in HbA1c compared to 0.3% in the control group [58]. These studies suggest that the health care intervention by the pharmacist in patients with DM can improve the HbA1c results acting as

an “additional” effect to the standard care provided to the patient. Other parameters such as blood pressure, LDL cholesterol, triglycerides, BMI, and coronary risk were also sensitive to pharmaceutical intervention in this group of patients, as reported by Pousinho et al. [59].

Barcelo et al.[60] found that most of the direct costs of treating DM in Latin America are related to the treatment of complications. Additionally,the cost of a patient without chronic complications with HbA1c > 10% is 2.4 times higher when compared to a better controlled patient, with HbA1c <8%. If the patient develops chronic complications, the resultant cost is 34 times higher [61]. Therefore, population-based DM screening actions can generate cost-effective results [62].

The campaign to track suspected cases of DM in Brazil, in 2018, reached a population larger than the number initially estimated, reinforcing the role of pharmacies in health promotion and disease prevention activities. As in other countries [51, 52], pharmacies in Brazil are health establishments that provide pharmaceutical services, individual and collective health guidance, handling and/or dispensing medications, and evaluation of CBG , blood pressure and weight measurements, among other procedures [24, 25]. Thus, pharmacies are favorable environments for carrying out clinical activities, including screening, education, and monitoring of people with NCD, particularly diabetes [63].

Strengths and limitations

This is the first Brazilian T2DM screening study in pharmacies,using CBG test. The sample screened in this study was higher than expected; thus, the approach presented herein reinforces the strategic importance of the health services provided by community pharmacies for screening actions across the country. Another strength of the study was the opportunity to expand the application of the FINDRISC tool, which has been already translated and validated for the Brazilian Portuguese and is strongly recommended by the Brazilian Diabetes Society in its yearly detection campaigns. The completion of the questionnaire by health professionals, as performed herein, is expected to have increased the reliability of the data [14].

This study has some limitations. Because it is a cross-sectional study, a causal relationship cannot be established. Another possible limitation was the population screened since it was a convenience sample and most participants had 11 or more years of formal education plus the fact that the majority of the pharmacies were private. Therefore, it is possible that the more vulnerable and least assisted people and also frequently less educated were under evaluated. In Brazil, 20.6% of the inhabitants have an educational level of less than 3 years of formal education [64] and only 5.6% of the evaluated sample had this characteristic. Thus, although the frequency reported herein is alarming, it may be underestimated.

Conclusions

This is the largest study ever conducted in Brazil to assess the frequency of high blood glucose levels and the risk for the development of T2DM , performed by pharmacists,. The profile of the people who

were found to have high CBG indicates that strategies involving health education measures, encouragement to healthy eating, physical activity and weight loss could help to reduce the prevalence of T2DM in the country, especially in the most underserved regions of the country, North, Northeast and Midwest where the increase in obesity is an important concern. Therefore, pharmacists working in community pharmacies may contribute to screening of NCD, such as T2DM, and perform health education, in the early referral of suspected cases to the health service for diagnostic confirmation, as well as in the monitoring of the patient with DM, in collaboration with the other health professionals and produce better results in the disease control.

Public health actions directed to the population with high CBG can enable the early diagnosis of T2DM, and contributes to the reduction of ASCVD, prevention of microvascular complications and avoid premature mortality. Worth to quote the partnership between the Brazilian Diabetes Society and the Federal Council of Pharmacists which represents an important step towards the liaison of pharmacists and other HCP involved in screening and care of people with diabetes in Brazil.

Abbreviations

ASCVD: Atherosclerotic Cardiovascular Diseases; BMI: Body Mass Index; CBG: Capilar Blood Glucose; CI: Confidence Interval; DM: Diabetes Mellitus; FINDRISC: Finnish Diabetes Risk Score; HCP: Health Care Professional; NCD: Non-Communicable Disease; PR: Prevalence ratio; T1DM: Type 1 Diabetes Mellitus; T2DM: Type 2 Diabetes Mellitus

Declarations

Acknowledgements

The authors acknowledge the support from pharmacists, the Brazilian Federal Council of Pharmacy and the Brazilian Diabetes Society during the activities of the Blue Diabetes November campaign.

Authors' contributions

Conception and study design: C.J.C., W.C.V., K.F.S.M., J.C.Q.PF., J.V.A., M.S.A.L., E.B.P., J.S.R., J.P., H.C.P., W.S.J.J. Analysis and interpretation of data: C.J.C., W.C.V., J.C.Q.PF., L.G.N., R.C.R.M.N., M.S.A.L., E.B.P., W.M.F. Drafting of the manuscript; CJS; WCV; RCRMN; JCQPF; LGN; WMF; HCP; WSJJ. All authors read and approved the final manuscript.

Funding

There was no funding for this study.

Availability of data and materials

The datasets used during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The research was approved by the Research Ethics Committees of the Federal University of Parana and the Federal University of Ouro Preto under number 95218118.4.0000.0102 / 2018. All interviews were conducted by the pharmacists preceded by clarifying the objectives for the participant and signing the informed consent form.

Consent for publication

A written consent to publish the information and data of the participants was obtained.

Competing interests

There are no competing interests to declare.

References

- [1] Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2095-128.
- [2] Arredondo A, Aviles R. Costs and epidemiological changes of chronic diseases: implications and challenges for health systems. *PloS one*. 2015;10:e0118611.
- [3] International Diabetes Federation. IDF Diabetes Atlas Ninth edition, accessed 16 jun 2020 in <https://www.diabetesatlas.org/en/>. 2019.
- [4] Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes research and clinical practice*. 2010;87:4-14.
- [5] Brasil. VIGITEL Brasil 2017 Vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Accessed in 08 april 2018 in https://bvsms.saude.gov.br/bvs/publicacoes/vigitel_brasil_2017_vigilancia_fatores_riscos.pdf. 2017.
- [6] Malta DC, Duncan BB, Schmidt MI, Machado IE, Silva AG, Bernal RTI, et al. Prevalence of diabetes mellitus as determined by glycated hemoglobin in the Brazilian adult population, National Health Survey. *REV BRAS EPIDEMIOL*. 2019;22: E190006.
- [7] Schmidt MI, Hoffmann JF, de Fatima Sander Diniz M, Lotufo PA, Griep RH, Bensenor IM, et al. High prevalence of diabetes and intermediate hyperglycemia - The Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Diabetology & metabolic syndrome*. 2014;6:123.

- [8] Mendes AB, Fittipaldi JA, Neves RC, Chacra AR, Moreira ED, Jr. Prevalence and correlates of inadequate glycaemic control: results from a nationwide survey in 6,671 adults with diabetes in Brazil. *Acta diabetologica*. 2010;47:137-45.
- [9] Gomes MB, Coral M, Cobas RA, Dib SA, Canani LH, Nery M, et al. Prevalence of adults with type 1 diabetes who meet the goals of care in daily clinical practice: a nationwide multicenter study in Brazil. *Diabetes research and clinical practice*. 2012;97:63-70.
- [10] Gomes MB, Gianella D, Faria M, Tambascia M, Fonseca RM, Rea R, et al. Prevalence of Type 2 diabetic patients within the targets of care guidelines in daily clinical practice: a multi-center study in Brazil. *The review of diabetic studies : RDS*. 2006;3:82-7.
- [11] Saraiva JFK, Hissa MN, Felício JS, Cavalcanti CAJ, Saraiva GL, Piha T, et al. Diabetes Mellitus in Brazil: clinical characteristics, standard treatment choice and cost. *J Bras Econ Saúde*. 2016;8:80-90.
- [12] Bahia LR, Araujo DV, Schaan BD, Dib SA, Negrato CA, Leao MP, et al. The costs of type 2 diabetes mellitus outpatient care in the Brazilian public health system. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2011;14:S137-40.
- [13] Lindstrom J, Tuomilehto J. The diabetes risk score: a practical tool to predict type 2 diabetes risk. *Diabetes care*. 2003;26:725-31.
- [14] Meijnikman AS, De Block CE, Verrijken A, Mertens I, Corthouts B, Van Gaal LF. Screening for type 2 diabetes mellitus in overweight and obese subjects made easy by the FINDRISC score. *Journal of diabetes and its complications*. 2016;30:1043-9.
- [15] Meijnikman AS, De Block CEM, Verrijken A, Mertens I, Van Gaal LF. Predicting type 2 diabetes mellitus: a comparison between the FINDRISC score and the metabolic syndrome. *Diabetology & metabolic syndrome*. 2018;10:12.
- [16] Makrilakis K, Liatis S, Grammatikou S, Perrea D, Stathi C, Tsiligros P, et al. Validation of the Finnish diabetes risk score (FINDRISC) questionnaire for screening for undiagnosed type 2 diabetes, dysglycaemia and the metabolic syndrome in Greece. *Diabetes & metabolism*. 2011;37:144-51.
- [17] Stiglic G, Fijacko N, Stozer A, Sheikh A, Pajnkihar M. Validation of the Finnish Diabetes Risk Score (FINDRISC) questionnaire for undiagnosed type 2 diabetes screening in the Slovenian working population. *Diabetes research and clinical practice*. 2016;120:194-7.
- [18] Sociedade Brasileira de Diabetes. Diretrizes da Sociedade Brasileira de Diabetes 2017-2018. 2017.
- [19] Krass I, Armour CL, Mitchell B, Brilliant M, Dienaar R, Hughes J, et al. The Pharmacy Diabetes Care Program: assessment of a community pharmacy diabetes service model in Australia. *Diabetic medicine : a journal of the British Diabetic Association*. 2007;24:677-83.

- [20] Fikri-Benbrahim N, Martinez-Martinez F, Saez-Benito L, Luque BS, Corpas JP, Moullin JC, et al. Assessment of a screening protocol for type 2 diabetes in community pharmacy. The DiabNow Study. *Diabetes research and clinical practice*. 2015;108:e49-52.
- [21] Hersberger KE, Botomino A, Mancini M, Bruppacher R. Sequential screening for diabetes—evaluation of a campaign in Swiss community pharmacies. *Pharmacy world & science : PWS*. 2006;28:171-9.
- [22] Papastergiou J, Folkins C, Li W. Community pharmacy-based A1c screening: a Canadian model for diabetes care. *The International journal of pharmacy practice*. 2016;24:189-95.
- [23] Willis A, Rivers P, Gray LJ, Davies M, Khunti K. The effectiveness of screening for diabetes and cardiovascular disease risk factors in a community pharmacy setting. *PloS one*. 2014;9:e91157.
- [24] Brasil. Lei no. 13.021, de 08 de agosto de 2014. Dispõe sobre o exercício e a fiscalização das atividades farmacêuticas. *Diário Oficial da União, Poder Executivo, Brasília, DF, 11 ago. 2014a. Seção 1, p. 1, Edição Extra. 2014.*
- [25] Anvisa. Resolução RDC no 44, de 17 de agosto de 2009. Dispõe sobre Boas Práticas Farmacêuticas para o controle sanitário do funcionamento, da dispensação e da comercialização de produtos e da prestação de serviços farmacêuticos em farmácias e drogarias e dá outras providências. 2009.
- [26] IBGE. Projection of the Brazilian population accessed in 15 jan 2017 in <https://www.ibge.gov.br/apps/populacao/projecao/>. 2017.
- [27] International Diabetes Federation. IDF Diabetes Atlas Ninth edition, accessed 14 jan 2018 in <https://www.diabetes.org.br/profissionais/images/2018/poster-atlas-idf-2017.pdf>. 2017.
- [28] World Health Organization. The legal and regulatory framework for community pharmacies in the WHO European Region. 100p. 2019.
- [29] Conselho Federal de Farmácia. November blue diabetes campaign <https://www.cff.org.br/pagina.php?id=826> Accessed 10 Jun 2020. 2018.
- [30] Conselho Federal de Farmácia. November blue diabetes campaign official website <http://www.diabetes.cff.org.br/> Accessed 10 Jun 2020. 2018.
- [31] Barim EM, McLellan KCP, Ribeiro RS, Carvalho JAM, Lindström J, Tuomilehto J, et al. Translation and cultural adaptation into Brazilian Portuguese of the Finnish Diabetes Risk Score (FINDRISC) and reliability assessment. *REV BRAS EPIDEMIOL*. 2020;23.
- [32] ABESO. Associação Brasileira para o Estudo da Obesidade e da Síndrome Metabólica Diretrizes brasileiras de obesidade Accessed in 10 jan 2018 in <https://abeso.org.br/wp-content/uploads/2019/12/Diretrizes-Download-Diretrizes-Brasileiras-de-Obesidade-2016.pdf>. 2016.

- [33] World Health Organization. Data Global status report on noncommunicable diseases 2014. Switzerland: WHO Library Cataloguing-in-Publication Data. Available in: accessed 16 jun 2020 in https://apps.who.int/iris/bitstream/handle/10665/148114/9789241564854_eng.pdf;jsessionid=7164ABD7E36ECA51992A1C50FA8FCF6B?sequence=1. 2014.
- [34] Rhee EJ. Nonalcoholic Fatty Liver Disease and Diabetes: An Epidemiological Perspective. *Endocrinology and metabolism*. 2019;34:226-33.
- [35] Stefan N, Haring HU, Cusi K. Non-alcoholic fatty liver disease: causes, diagnosis, cardiometabolic consequences, and treatment strategies. *The lancet Diabetes & endocrinology*. 2019;7:313-24.
- [36] Dias PC, Henriques P, Anjos LA, Burlandy L. Obesity and public policies: the Brazilian government's definitions and strategies. *Cad Saúde Pública* 2017;33:1-12.
- [37] Brasil. VIGITEL Brasil 2018 Vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Accessed in april 2019 in <https://portalarquivos2.saude.gov.br/images/pdf/2019/julho/25/vigitel-brasil-2018.pdf>. 2018.
- [38] Cepas V, Collino M, Mayo JC, Sainz RM. Redox Signaling and Advanced Glycation Endproducts (AGEs) in Diet-Related Diseases. *Antioxidants*. 2020;9.
- [39] Nowotny K, Jung T, Hohn A, Weber D, Grune T. Advanced glycation end products and oxidative stress in type 2 diabetes mellitus. *Biomolecules*. 2015;5:194-222.
- [40] Kowluru RA, Kanwar M. Effects of curcumin on retinal oxidative stress and inflammation in diabetes. *Nutrition & metabolism*. 2007;4:8.
- [41] Kaul K, Hodgkinson A, Tarr JM, Kohner EM, Chibber R. Is inflammation a common retinal-renal-nerve pathogenic link in diabetes? *Current diabetes reviews*. 2010;6:294-303.
- [42] Herder C, Roden M, Ziegler D. Novel Insights into Sensorimotor and Cardiovascular Autonomic Neuropathy from Recent-Onset Diabetes and Population-Based Cohorts. *Trends in endocrinology and metabolism: TEM*. 2019;30:286-98.
- [43] Kruit A, Zanen P. The association between vitamin D and C-reactive protein levels in patients with inflammatory and non-inflammatory diseases. *Clinical biochemistry*. 2016;49:534-7.
- [44] Pittas AG, Jorde R, Kawahara T, Dawson-Hughes B. Vitamin D supplementation for prevention of type 2 diabetes mellitus. To D or not to D? *The Journal of clinical endocrinology and metabolism*. 2020.
- [45] Passos VM, Barreto SM, Diniz LM, Lima-Costa MF. Type 2 diabetes: prevalence and associated factors in a Brazilian community—the Bambui health and aging study. *Sao Paulo medical journal = Revista paulista de medicina*. 2005;123:66-71.

- [46] Mayer-Davis EJ, D'Agostino R, Jr., Karter AJ, Haffner SM, Rewers MJ, Saad M, et al. Intensity and amount of physical activity in relation to insulin sensitivity: the Insulin Resistance Atherosclerosis Study. *Jama*. 1998;279:669-74.
- [47] Alberti KG, Zimmet P, Shaw J. International Diabetes Federation: a consensus on Type 2 diabetes prevention. *Diabetic medicine : a journal of the British Diabetic Association*. 2007;24:451-63.
- [48] Kirpichnikov D, Sowers JR. Diabetes mellitus and diabetes-associated vascular disease. *Trends in endocrinology and metabolism: TEM*. 2001;12:225-30.
- [49] Sowers JR, Epstein M, Frohlich ED. Diabetes, hypertension, and cardiovascular disease: an update. *Hypertension*. 2001;37:1053-9.
- [50] Fornos-Perez JA, Andres-Rodriguez NF, Andres-Iglesias JC, Luna-Cano R, Garcia-Soidan J, Lorenzo-Veiga B, et al. Detection of people at risk of diabetes in community pharmacies of Pontevedra (Spain) (DEDIPO). *Endocrinologia y nutricion : organo de la Sociedad Espanola de Endocrinologia y Nutricion*. 2016;63:387-96.
- [51] Dhipayom T, Fuangchan A, Tunpichart S, Chaiyakunapruk N. Opportunistic screening and health promotion for type 2 diabetes: an expanding public health role for the community pharmacist. *Journal of public health*. 2013;35:262-9.
- [52] Bovet P, Hirsiger P, Emery F, De Bernardini J, Rossier C, Trebeljahr J, et al. Impact and cost of a 2-week community-based screening and awareness program for diabetes and cardiovascular risk factors in a Swiss canton. *Diabetes, metabolic syndrome and obesity : targets and therapy*. 2011;4:213-23.
- [53] Chaturvedi N. The burden of diabetes and its complications: trends and implications for intervention. *Diabetes research and clinical practice*. 2007;76 Suppl 1:S3-12.
- [54] Saaristo T, Peltonen M, Keinanen-Kiukaanniemi S, Vanhala M, Saltevo J, Niskanen L, et al. National type 2 diabetes prevention programme in Finland: FIN-D2D. *International journal of circumpolar health*. 2007;66:101-12.
- [55] Grisold A, Callaghan BC, Feldman EL. Mediators of diabetic neuropathy: is hyperglycemia the only culprit? *Current opinion in endocrinology, diabetes, and obesity*. 2017;24:103-11.
- [56] Sociedade Brasileira de Diabetes. Diretrizes Sociedade Brasileira de Diabetes 2019-2020 Accessed in jun 2020 in <https://www.diabetes.org.br/profissionais/images/DIRETRIZES-COMPLETA-2019-2020.pdf>. 2019.
- [57] Fazel MT, Bagalagel A, Lee JK, Martin JR, Slack MK. Impact of Diabetes Care by Pharmacists as Part of Health Care Team in Ambulatory Settings: A Systematic Review and Meta-analysis. *The Annals of pharmacotherapy*. 2017;51:890-907.

- [58] Correr CJ, Melchior AC, Fernandez-Llimos F, Pontarolo R. Effects of a pharmacotherapy follow-up in community pharmacies on type 2 diabetes patients in Brazil. *International journal of clinical pharmacy*. 2011;33:273-80.
- [59] Pousinho S, Morgado M, Falcao A, Alves G. Pharmacist Interventions in the Management of Type 2 Diabetes Mellitus: A Systematic Review of Randomized Controlled Trials. *Journal of managed care & specialty pharmacy*. 2016;22:493-515.
- [60] Barcelo A, Arredondo A, Gordillo-Tobar A, Segovia J, Qiang A. The cost of diabetes in Latin America and the Caribbean in 2015: Evidence for decision and policy makers. *Journal of global health*. 2017;7:020410.
- [61] Menzin J, Langley-Hawthorne C, Friedman M, Boulanger L, Cavanaugh R. Potential short-term economic benefits of improved glycemic control: a managed care perspective. *Diabetes care*. 2001;24:51-5.
- [62] Georg AE, Duncan BB, Toscano CM, Schmidt MI, Mengue S, Duarte C, et al. Economic analysis of a screening program for diabetes mellitus in Brazil. *Rev Saúde Pública*. 2005;39:452-60.
- [63] Conselho Federal de Farmácia. Resolução nº 585, de 29 de Agosto de 2013. *Diário Oficial da União*. 2013.
- [64] IBGE. Brazil in summary, education, years of study. <https://brasilemsintese.ibge.gov.br/educacao/anos-de-estudo.html> Accessed in 01 of september of 2020 2020.

Figures




Figure 1

Population included in the screening of high blood glucose in Brazil, 2018.




Figure 2

Prevalence of high capillary blood glucose, according to Brazilian region, 2018