

Outcomes of children with hepatoblastoma who underwent liver resection at a tertiary hospital in China: A retrospective analysis

Jiahao Li

Guangzhou Women and Children's Medical Center

Huixian Li

Guangzhou Women and Children's Medical Center

Huiying Wu

Guangzhou Women and Children's Medical Center

Huilin Niu

Guangzhou Women and Children's Medical Center

Haibo Li

Guangzhou Women and Children's Medical Center

Jing Pan

Guangzhou Women and Children's Medical Center

Jiliang Yang

Guangzhou Women and Children's Medical Center

Tianbao Tan

Guangzhou Women and Children's Medical Center

Chao Hu

Guangzhou Women and Children's Medical Center

Tao Xu

Guangzhou Women and Children's Medical Center

Xiaohong Zhang

Guangzhou Women and Children's Medical Center

Manna Zheng

Guangzhou Women and Children's Medical Center

Kuanrong Li

Guangzhou Women and Children's Medical Center

Yan Zou

Guangzhou Women and Children's Medical Center

Tianyou Yang (✉ mdtianyouyang@hotmail.com)


Department of Pediatric Surgery, Guangzhou Women and Children's Medical Center, Affiliated Women and Children's Medical Center of Guangzhou Medical University, Guangzhou, China. <https://orcid.org/0000-0001-9140-3806>

Research article

Keywords: hepatoblastoma; surgery; children; liver tumour

Posted Date: January 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.15344/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Pediatrics on May 9th, 2020. See the published version at <https://doi.org/10.1186/s12887-020-02059-z>.

Abstract

Objective: To report the outcomes of hepatoblastoma resected in our institution. **Methods:** We diagnosed 135 children with hepatoblastoma at our institution between January 2010 and December 2017. Patients who underwent liver resection were included for analysis. However, patients who abandoned treatment after diagnosis were excluded from analysis, but their clinical characteristics were provided in the supplementary material. **Results:** Forty-two patients abandoned treatment, whereas 93 patients underwent liver resection and were included for statistical analysis. Thirty-six, 23, 3, and 31 patients had PRETEXT stages II, III, IV, and unspecified tumours, respectively. Seven patients had ruptured tumour; 9 had lung metastasis (one patient had portal vein thrombosis concurrently). Sixteen patients underwent primary liver resection; 22, 25, and 30 patients received cisplatin-based neoadjuvant chemotherapy and delayed surgery, preoperative transarterial chemoembolization (TACE) and delayed surgery, and a combination of cisplatin-based neoadjuvant chemotherapy, TACE, and delayed surgery, respectively. Forty patients had both PRETEXT and POST-TEXT information available for analysis. Twelve patients were down-staged after preoperative treatment, including 2, 8, and 2 patients from stages IV to III, III to II, and II to I, respectively. Ten patients with unspecified PRETEXT stage were confirmed to have POST-TEXT stages II (n=8) and I (n=2) tumours. Seven tumours were associated with positive surgical margins, and 12 patients had microvascular involvement. During a median follow-up period of 30.5 months, 84 patients survived without relapse, 9 experienced tumour recurrence, and 4 died. The 2-year event-free survival (EFS) and overall survival (OS) rates were $89.4 \pm 3.4\%$, and $95.2 \pm 2.4\%$, respectively; they were significantly better among patients without metastasis (no metastasis vs metastasis: EFS, $93.5 \pm 3.7\%$ vs $46.7 \pm 19.0\%$, adjusted $p = 0.004$. OS, $97.6 \pm 2.4\%$ vs $61.0 \pm 18.1\%$, adjusted $p = 0.003$), and similar among patients treated with different preoperative strategies (chemotherapy only vs TACE only vs Both: EFS, $94.7 \pm 5.1\%$ vs $91.7 \pm 5.6\%$ vs $85.6 \pm 6.7\%$, $p = 0.542$. OS, $94.1 \pm 5.7\%$ vs $95.7 \pm 4.3\%$ vs $96.7 \pm 3.3\%$, $p = 0.845$). **Conclusion:** The OS for patients with hepatoblastoma who underwent liver resection was satisfactory. Neoadjuvant chemotherapy and TACE seemed to have a similar effect on OS. However, the abandonment of treatment by patients with hepatoblastoma was common, and may have biased our results.

Background

Hepatoblastoma is the most common childhood liver malignancy, and has a prevalence of 1 per 1 000 000 population(1, 2). The incidence of hepatoblastoma has increased in the past two decades, and this upward trend has been correlated with an increasing survival rate among premature and low-birth-weight infants(3). Hepatoblastoma usually affects children younger than 3 years, and presents as a large abdominal mass. Some patients may present with sudden abdominal pain and haemorrhagic shock in the scenario of tumour rupture. A combination of elevated α -fetoprotein protein (AFP) level and radiographically identified hepatic mass suffices for the clinical diagnosis of hepatoblastoma in children with ages between 6 months and 3 years. However, biopsy, preferably via ultrasound-guided core needle biopsy is recommended for patients of all age groups(4, 5).

The treatment of hepatoblastoma is multidisciplinary; a combination of platinum-based chemotherapy and complete surgical removal is the mainstay of treatment. Cisplatin-based chemotherapy and surgical resection provide standard-risk patients with a 5-year overall survival (OS) of more than 90%(6, 7). Primary

hepatic resection is recommended for patients with PRETEXT stages I and II tumours with no additional annotative risk factors. Otherwise, patients should undergo neoadjuvant chemotherapy and delayed surgery. Orthotopic liver transplantation is an ideal treatment option for patients with PRETEXT stage IV hepatoblastomas and other forms of unresectable hepatoblastomas, and can provide them with more than 80% 5-year OS in the contemporary era(7-9).

Nonetheless, the outcomes of hepatoblastoma in developing countries are still far more inferior to those in developed countries(10). Treatment abandonment among children with cancer is not an unusual phenomenon in developing countries, particularly among those with advanced stage cancers(11). Furthermore, patients in developing countries have far more limited access to liver transplantation. In order to improve the management and outcomes of hepatoblastoma in developing countries, such experiences are worth reporting. Herein, we described our experiences in treating hepatoblastoma at a tertiary hospital in South China.

Methods

The diagnosis of hepatoblastoma was initially made based on an elevated AFP level and radiographic detection of a liver mass, and confirmed via pathological examination of samples obtained via either biopsy or primary liver resection. Only hepatoblastoma patients who underwent liver resection were included for statistical analysis. Patients who abandoned treatment were excluded from further analysis. Patients with hepatocellular carcinoma and other liver malignancies were excluded.

One hundred and thirty-five children were diagnosed with hepatoblastoma at our institution between January 2010 and December 2017. Forty-two cases were excluded from the analysis mainly due to treatment abandonment, including 6 cases who died due to aggressive tumour progression prior to treatment and 36 cases that received no further treatment after diagnosis. The demographic and clinical characteristics of these excluded patients was presented in the supplementary Table 1. Our study analysed 93 cases that were treated according to the institutional protocol and underwent liver resection. The chemotherapy regimens of COG (Children's Oncology Group), SIOPEL (International Childhood Liver Tumors Strategy Group), and our national regimens were used. All these chemotherapy regimens were cisplatin-based and were reported to have similar effects and achieved similar survival outcomes(12). Patients were followed up at the clinic and via regular telephone calls. The primary outcome was to evaluate the event-free survival and overall survival of hepatoblastoma resected in our institution. The secondary outcome was to analyse factors that would impact survival in this cohort of patients. The OS duration was defined as the interval between the time of diagnosis and the time of death, and event-free survival (EFS) as the interval between the time of diagnosis and the time of the first occurrence of tumour progression, relapse, or death, whichever occurred first.

We collected information regarding patients' demographic data, including age and gender; clinical data including AFP level, radiographic findings, pre-treatment extent of tumour (PRETEXT) and post-treatment extent of tumour (POST-TEXT) staging, preoperative management strategy (neoadjuvant chemotherapy and transarterial chemoembolisation [TACE]), and liver resection technique; pathological findings including pathological subtype, surgical margin status, microvascular involvement, and lymph node involvement; and clinical outcomes including disease relapse and death.

A standard data extraction form with a logical organisation similar in flow to the format of the original medical charts, was used to collect data. Two trained data abstractors, who were blinded to the study hypothesis, independently reviewed the original medical charts and collected data. Explicit criteria for extracting data regarding variables were applied. Any discrepancies between the abstractors were reviewed jointly and discussed to clarify any issues(13).

A senior radiologist, who was blinded to the study objective, retrospectively reviewed patients' computed tomography (CT) and magnetic resonance imaging (MRI) data. The radiologist defined the PRETEXT/POST-TEXT system and annotation factors according to the PRETEXT staging system(14). Not all patients had CT/MRI images stored in the electronic database; only patients who underwent CT/MRI scans at our institution had their radiographic images stored.

The study protocol was approved by the institutional review board of Guangzhou Women and Children's Medical Centre. The need for informed consent was waived on account of the retrospective nature of the demographic, clinical, and outcome data. All patients' data were de-identified prior to the analysis.

Statistical analysis

Categorical variables are presented as numbers and percentages. Continuous variables are presented as medians and ranges. The PRETEXT and POST-TEXT stages were compared using the McNemar chi-square test. The comparison of different management strategies was analysed using the Wilcoxon signed-rank test. The probabilities of OS and EFS were computed using the Kaplan-Meier method and compared using the log-rank test. Statistical significance was set at $p < 0.05$ and p -values of the paired tests in the log-rank test were adjusted using the Bonferroni method. All statistical analyses were performed using SAS 9.4 for Windows (SAS Institute Inc., Cary, NC, USA).

Results

Patients' demographic and clinical characteristics

Of the 93 patients who underwent liver resection, 66 (60.2%) were male and 37 (39.8%) were female (Table 1). The median age at diagnosis was 11 (range, 1.7–87) months. The median AFP level was 76,131 (range, 10–1,881,360) ng/ml and the median tumour diameter was 10.6 (range, 5.1–15.8) cm. Fifty-seven (61.3%) patients had unifocal tumours, 7 (7.5%) had multifocal tumours, and 29 (31.2%) had tumours with unspecified focality.

Thirty-six (38.7%) patients had PRETEXT stage II tumours, 23 (24.7%) had stage III tumours, 3 (3.2%) had stage IV tumours, and 31 (33.3%) had tumours with unspecified PRETEXT stages. Seven (7.5%) patients had ruptured tumours, 9 (9.7%) had lung metastasis [1 (1.1%) had portal vein thrombosis concurrently]. Sixteen (17.2%) patients underwent primary liver resection. Twenty-two patients (23.7%) received cisplatin-based neoadjuvant chemotherapy and delayed surgery, 25 (26.9%) received preoperative transarterial chemoembolism (TACE) and delayed surgery, and 30 (32.3%) received a combination of cisplatin-based neoadjuvant chemotherapy, TACE, and delayed surgery. The median number of treatment cycles was 2.5 (range, 1–8) for neoadjuvant chemotherapy and 2 (range, 1–7) for preoperative TACE.

Forty patients had information regarding both PRETEXT and POST-TEXT stages available for analysis. Using the McNemar test, significant downstage was noted for the 12 cases with both PRETEXT and POST-TEXT stage information ($p<0.001$). Specifically, 2 cases from stage IV to III, 8 from stage III to II, and 2 from stage II to I. Furthermore, 10 patients with unspecified PRETEXT stage were confirmed to have POST-TEXT stages II ($n=8$) and I ($n=2$) tumours.

The detailed demographic and clinical characteristics of the excluded 42 patients were listed in supplementary table 1. The excluded patients were significantly higher in age, AFP value, and PRETEXT stage than the included 93 patients. Additionally, more patients of the excluded group had lung metastases and portal vein thrombosis. The overall outcomes of these patients were largely unknown, and these patients were excluded from further analysis.

Surgery and outcomes

Thirty-seven (39.8%) patients underwent hemihepatectomy, 17 (18.3%) underwent wedge resection, 13 (14.0%) underwent trisectionectomy, 9 (9.7%) underwent bisegmentectomy (left lateral sectionectomy), and 2 (2.2%) underwent central hepatectomy. Fifteen (16.1%) patients underwent tumour resection at other institutions. The operative time, estimated volume of blood lost, and volume of red blood cells transfused were 290 (range, 100–510) minutes, 100 (range, 20–1,000) ml, and 1 (range, 0–6) units, respectively. There were 24 (25.8%) cases of epithelial variant hepatoblastoma, 11 (11.8%) cases of mixed epithelial hepatoblastoma, and 41 (44.1%) cases of mixed epithelial and mesenchymal hepatoblastoma; 17 cases were not subclassified. Seven (7.5%) cases had positive surgical margins, 69 (74.2%) had negative surgical margins, and 17 (18.3%) had unspecified surgical margin status. Twelve (12.9%) patients had microvascular involvement, 43 (46.2%) had no microvascular involvement, and 38 (40.9%) cases had unspecified microvascular status. Thirty-one patients underwent lymph node dissection, none of whom had positive lymph node involvement.

Sixty-three (67.7%) patients received cisplatin-based postoperative chemotherapy, with a median of 6 (range, 1–12) cycles. Twenty-seven (29.0%) patients received no postoperative chemotherapy. During a median follow-up duration of 30.5 (range, 0.7–105.1) months, 84 (90.3%) cases survived without relapse, 9 (9.7%) experienced disease recurrence, and 4 (5.4%) died.

Subgroup analysis of managements

In this study, the differences in management between patients without metastasis and patients with metastasis (1 of them had portal vein thrombosis at the same time) [cycle of neoadjuvant chemotherapy: 1(0-6) vs 2(0-8), $p=0.060$; cycle of preoperative TACE: 0(0-5) vs 1(0-7), $p=0.589$; cycle of postoperative chemotherapy: 6(0-12) vs 6(2-10), $p=0.817$], and patients with negative surgical margin and positive surgical margins [cycle of neoadjuvant chemotherapy: 1(0-8) vs 1(0-3), $p=0.482$; cycle of preoperative TACE: 1(0-5) vs 2(0-7), $p=0.081$; cycle of postoperative chemotherapy: 6(0-12) vs 7(2-12), $p=0.946$] were not statistically significant.

Failure among patients with tumour recurrence

Among the 9 patients with tumour recurrence, the median time from diagnosis to recurrence was 8.5 (range, 0.7–22.4) months. Among the 4 patients who died as a result of tumour recurrence, the median time from diagnosis to death was 11.3 (range, 3.6–21.4) months. Their treatment and outcome information are summarised in Table 3. Five patients underwent wedge resection, and 1 underwent left hepatectomy associated with a positive surgical margin.

Survival

The 2-year event-free survival (EFS) and overall survival (OS) rates were $89.4 \pm 3.4\%$, and $95.2 \pm 2.4\%$ (Figs. 1A and 2A), respectively. The 2-year EFS and OS rates were significantly better among patients without metastasis (no metastasis vs metastasis: EFS, $93.5 \pm 3.7\%$ vs $46.7 \pm 19.0\%$, $p=0.004$, OS, $97.6 \pm 2.4\%$ vs $61.0 \pm 18.1\%$, $p=0.003$) (Figs. 1C and 2C) than among patients with metastasis. The differences in the 2-year EFS and OS rates of patients with PRETEXT stage IV hepatoblastoma (None vs None vs None vs Unknown: EFS, $84.0 \pm 6.7\%$ vs $95.7 \pm 4.3\%$ vs $66.7 \pm 27.2\%$ vs $93.1 \pm 4.7\%$, $p=0.216$. OS, $90.1 \pm 5.5\%$ vs $95.5 \pm 4.4\%$ vs 100.0% vs 100.0% , $p=0.140$), positive surgical margins (negative vs positive vs Unknown: EFS, $92.0 \pm 3.5\%$ vs $64.3 \pm 21.0\%$ vs $87.5 \pm 8.3\%$, $p=0.259$. OS, $95.0 \pm 2.8\%$ vs $83.3 \pm 15.2\%$ vs 100.0% , $p=0.226$), and microvascular involvement (No vs Involvement vs Unknown: EFS, $95.3 \pm 3.3\%$ vs $67.3 \pm 16.0\%$ vs $88.7 \pm 5.4\%$, $p=0.07$. OS, $97.7 \pm 2.3\%$ vs $90.0 \pm 9.5\%$ vs $93.9 \pm 4.2\%$, $p=0.601$) were not statistically significant. The 2-year EFS and OS rates were also similar among patients treated with different preoperative strategies (Chemotherapy only vs TACE only vs Both: EFS, $94.7 \pm 5.1\%$ vs $91.7 \pm 5.6\%$ vs $85.6 \pm 6.7\%$, $p=0.542$. OS, $94.1 \pm 5.7\%$ vs $95.7 \pm 4.3\%$ vs $96.7 \pm 3.3\%$, $p=0.845$) (Figs. 1D and 2D).

Discussion

Here, we reported the outcomes of resected hepatoblastoma at a tertiary children's institution in a developing country. The 2-year EFS and OS rates among patients who underwent hepatic resection were satisfactory. Patients associated with distant metastasis had a worse prognosis, with 2-year EFS and OS rates of about $46.7 \pm 19.0\%$ and $61.0 \pm 18.1\%$, respectively. Neoadjuvant chemotherapy and TACE seem to have similar effects on the 2-year EFS and OS.

Both cisplatin-based neoadjuvant chemotherapy and preoperative TACE were used at our institution as preoperative strategies to shrink the tumour and downstage the tumour(15). However, our results showed no significant differences regarding the effect of neoadjuvant chemotherapy and TACE on 2-year EFS and OS. Similarly, evidence from the Japanese Study Group for Paediatric Liver Tumour (JPLT) and our institution showed that TACE was as effective as neoadjuvant chemotherapy in shrinking and downstaging tumours(15, 16). However, the JPLT study showed that the OS was inferior to that of those who underwent neoadjuvant chemotherapy(16). TACE could be an option for patients who fail to respond to neoadjuvant chemotherapy. Furthermore, TACE is particularly useful for patients who experience tumour rupture(17). Currently, neoadjuvant chemotherapy is considered the first choice for the preoperative management of hepatoblastoma. However, no prospective study has compared the effect of neoadjuvant chemotherapy and TACE on hepatoblastoma. It would be valuable to compare these two strategies in a prospective or randomized trial.

Patients with tumour metastasis had significantly lower 2-year EFS and OS. The 2-year EFS and OS for patients with metastatic disease were only about $46.7 \pm 19.0\%$ and $61.0 \pm 18.1\%$, respectively. Our result was consistent with the SIOPEL experiences, which showed that hepatoblastoma with metastasis has a 3-year EFS of 49%(18). However, we failed to demonstrate that patients with PRETEXT stage IV tumours had significantly worse EFS and OS probabilities than those with tumours of other stages. However, our cohort only had 3 cases with PRETEXT stage IV tumours. Two cases were down-staged to POST-TEXT stage III, and the other died.

The 2-year EFS and OS for patients with positive surgical margins or microvascular involvement were lower than those of their counterparts, but the differences were not statistically significant. The evidence suggested that positive surgical margin might not affect the EFS and OS in the setting of neoadjuvant chemotherapy(19). However, this might not be true in the setting of primary resection. Complete resection with a negative resection margin should always be pursued. Microvascular involvement was suggested to be a poor prognostic factor in a retrospective study(20). In our cohort, twelve (12.9%) patients had microvascular involvement, 43 patients had no microvascular involvement, and 38 patients had tumours with unspecified microvascular status. Our data suggested that although patients with microvascular involvement had lower 2-year EFS and OS than those without microvascular involvement, the differences were non-significant. Again, in the current Children's Hepatic tumours International Collaboration classification system, microvascular involvement is not considered as a risk factor(6, 21).

Hepatoblastoma seemed not to spread through the lymph nodes. None of the 31 patients who underwent lymph node biopsy had positive lymph node involvement.

Five out of 9 patients who experienced relapse or died underwent wedge resection. This suggests that wedge resection might be associated with worse outcomes. Standard hepatic resection should always be pursued in any possible scenario.

Due to the retrospective nature of this study, we were unable to retrieve some of the important information. For example, some of the patients did not undergo preoperative CT or MRI scans for PRETEXT staging. Furthermore, a large proportion of the patients abandoned or discontinued treatment after the establishment of the diagnosis. These patients will most likely fall into the high-risk group (Supplemental Table 1). In fact, the excluded patients were significantly higher in age and PRETEXT stage than included patients. Among the excluded patients, more patients had metastasis and portal vein thrombosis. Overall, the excluded patients mostly had advanced stage hepatoblastoma, and would have much worse survival. Unfortunately, we were not able to follow these excluded patients. The exclusion of these patients will incur selection bias. Treatment abandonment is not an unusual phenomenon in developing countries, which underscores the need for more attention and funding for this vulnerable population(22, 23). Furthermore, the follow-up duration was not long enough, and the EFS and OS might either be overestimated if patients abandoned treatment due to poor results, or underestimated if patients abandoned treatment because their parents prematurely assumed they were cured. An assessment of the interactions between different characteristics requires more stable follow-up with larger samples.

Conclusions

The overall outcomes for those who underwent liver resection was satisfactory. However, the abandonment of treatment by patients with hepatoblastoma was common. A large proportion of patients discontinued treatment after the diagnosis.

Abbreviations

AFP alpha-fetoprotein

CT computed tomography

CHIC Children's Hepatic tumors International Collaboration

COG Children's Oncology Group

EFS event-free survival

JPLT Japanese Study Group for Pediatric Liver Tumor

MRI magnetic resonance imaging

OS overall survival

PRETEXT pre-treatment extent of tumour

POST-TEXT post-treatment extent of tumour

SIOPEL International Childhood Liver Tumors Strategy Group

TACE transarterial chemoembolisation

Declarations

Ethics approval and consent to participate

Not applicable

Consent to publish

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due to patient privacy but are available from the corresponding author on reasonable request.

Competing interests

All authors declare no competing interests.

Funding

None

Authors' Contributions

Tianyou Yang and Yan Zou conceptualized and designed the study, Jiahao Li and Huixian Li drafted the initial manuscript, Tianyou Yang, Yan Zou reviewed and revised the manuscript.

Jiahao Li, Huixian Li, Huiying Wu, Huilin Niu, Haibo Li, Jing Pan, Jiliang Yang, Tianbao Tan, Chao Hu, Tao Xu, Xiaohong Zhang, Manna Zheng, Kuanrong Li designed the data collection instruments, collected data, carried out the initial analyses, and reviewed and revised the manuscript.

Tianyou Yang coordinated and supervised data collection, and critically reviewed the manuscript for important intellectual content.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Acknowledgements

None

References

1. Darbari A, Sabin KM, Shapiro CN, Schwarz KB. Epidemiology of primary hepatic malignancies in U.S. children. *Hepatology*. 2003;**38**:560-6 doi: 10.1053/jhep.2003.50375 [published Online.
2. Linabery AM, Ross JA. Trends in childhood cancer incidence in the U.S. (1992-2004). *Cancer*. 2008;**112**:416-32 doi: 10.1002/cncr.23169 [published Online First: 2007/12/13].
3. Hung G-Y, Lin L-Y, Yu T-Y, Lee C-Y, Yen H-J, Horng J-L. Hepatoblastoma incidence in Taiwan: A population-based study. *Journal of the Chinese Medical Association*. 2018;**81**:541-7 doi: 10.1016/j.jcma.2017.11.012 [published Online.
4. Hafberg E, Borinstein SC, Alexopoulos SP. Contemporary management of hepatoblastoma. *Curr Opin Organ Transplant*. 2019;**24**:113-7 doi: 10.1097/mot.0000000000000618 [published Online First: 2019/02/15].
5. Weldon CB, Madenci AL, Tiao GM, *et al*. Evaluation of the diagnostic biopsy approach for children with hepatoblastoma: A report from the Children's Oncology Group AHEP 0731 Liver Tumor Committee. *J Pediatr Surg*. 2019; doi: 10.1016/j.jpedsurg.2019.05.004 [published Online First: 2019/05/28].
6. Meyers RL, Maibach R, Hiyama E, *et al*. Risk-stratified staging in paediatric hepatoblastoma: a unified analysis from the Children's Hepatic tumors International Collaboration. *Lancet Oncol*. 2017;**18**:122-31 doi: 10.1016/s1470-2045(16)30598-8 [published Online First: 2016/11/26].
7. Lim IIP, Bondoc AJ, Geller JI, Tiao GM. Hepatoblastoma-The Evolution of Biology, Surgery, and Transplantation. *Children (Basel, Switzerland)*. 2018;**6** doi: 10.3390/children6010001 [published Online.
8. Busweiler LA, Wijnen MH, Wilde JC, *et al*. Surgical treatment of childhood hepatoblastoma in the Netherlands (1990-2013). *Pediatric surgery international*. 2017;**33**:23-31 doi: 10.1007/s00383-016-3989-

- 8 [published Online First: 2016/10/13].
9. Ezekian B, Mulvihill MS, Schroder PM, *et al.* Improved contemporary outcomes of liver transplantation for pediatric hepatoblastoma and hepatocellular carcinoma. *Pediatr Transplant.* 2018;**22**:e13305 doi: 10.1111/petr.13305 [published Online First: 2018/10/21].
 10. Magrath I, Steliarova-Foucher E, Epelman S, *et al.* Paediatric cancer in low-income and middle-income countries. *Lancet Oncol.* 2013;**14**:e104-16 doi: 10.1016/S1470-2045(13)70008-1 [published Online].
 11. Friedrich P, Lam CG, Itriago E, Perez R, Ribeiro RC, Arora RS. Magnitude of Treatment Abandonment in Childhood Cancer. *PLoS one.* 2015;**10**:e0135230 doi: 10.1371/journal.pone.0135230 [published Online].
 12. Yuan XJ, Wang HM, Jiang H, *et al.* Multidisciplinary effort in treating children with hepatoblastoma in China. *Cancer Lett.* 2016;**375**:39-46 doi: 10.1016/j.canlet.2016.02.051 [published Online First: 2016/03/08].
 13. Yang T, Li H, Li J, *et al.* Surgical risk factors of retroperitoneal teratoma resection in children. *Journal of Pediatric Surgery.* 2018; doi: <https://doi.org/10.1016/j.jpedsurg.2018.09.020> [published Online].
 14. Towbin AJ, Meyers RL, Woodley H, *et al.* 2017 PRETEXT: radiologic staging system for primary hepatic malignancies of childhood revised for the Paediatric Hepatic International Tumour Trial (PHITT). *Pediatric radiology.* 2018;**48**:536-54 doi: 10.1007/s00247-018-4078-z [published Online].
 15. Tan X, Zhang J, Wen Z, *et al.* Preoperative transcatheter arterial chemoembolization of hepatoblastoma in infants. *J Vasc Interv Radiol.* 2014;**25**:1029-35 doi: 10.1016/j.jvir.2014.03.032 [published Online First: 2014/05/21].
 16. Sasaki F, Matsunaga T, Iwafuchi M, *et al.* Outcome of hepatoblastoma treated with the JPLT-1 (Japanese Study Group for Pediatric Liver Tumor) Protocol-1: A report from the Japanese Study Group for Pediatric Liver Tumor. *J Pediatr Surg.* 2002;**37**:851-6 Online.
 17. Yang T, Tan T, Yang J, *et al.* Ruptured hepatoblastoma successfully treated with cisplatin monochemotherapy: A case report. *Molecular and clinical oncology.* 2018;**9**:223-5 doi: 10.3892/mco.2018.1643 [published Online First: 2018/08/14].
 18. Maibach R, Roebuck D, Brugieres L, *et al.* Prognostic stratification for children with hepatoblastoma: the SIOPEL experience. *European journal of cancer (Oxford, England : 1990).* 2012;**48**:1543-9 doi: 10.1016/j.ejca.2011.12.011 [published Online].
 19. Ren X, Li H, Diao M, Chen L, Xu H, Li L. Results of surgical resections with positive margins for children with hepatoblastoma: Case series from a single Asian center. *Pediatric blood & cancer.* 2019;**66** doi: 10.1002/pbc.27479 [published Online].
 20. Shi Y, Commander SJ, Masand PM, Heczey A, Goss JA, Vasudevan SA. Vascular invasion is a prognostic indicator in hepatoblastoma. *Journal of Pediatric Surgery.* 2017;**52**:956-61 doi: 10.1016/j.jpedsurg.2017.03.017 [published Online].
 21. Czauderna P, Haeberle B, Hiyama E, *et al.* The Children's Hepatic tumors International Collaboration (CHIC): Novel global rare tumor database yields new prognostic factors in hepatoblastoma and becomes a research model. *European journal of cancer (Oxford, England : 1990).* 2016;**52**:92-101 doi: 10.1016/j.ejca.2015.09.023 [published Online].

22. Friedrich P, Lam CG, Kaur G, Itriago E, Ribeiro RC, Arora RS. Determinants of Treatment Abandonment in Childhood Cancer: Results from a Global Survey. *PloS one*. 2016;**11**:e0163090 doi: 10.1371/journal.pone.0163090 [published Online.
23. Arora RS, Eden T, Pizer B. The problem of treatment abandonment in children from developing countries with cancer. *Pediatric blood & cancer*. 2007;**49**:941-6 doi: 10.1002/pbc.21127 [published Online.

Tables

TABLE 1 Demographic, clinical, radiological, and pathological characteristics of the study cohort

Characteristics	Number or as shown	Proportion (%)
All	93	100
Gender		
Male	56	60.2
Female	37	39.8
Age [median (range)], months	11 (1.7-87)	-
AFP level [median (range)], ng/ml	76,131 (10-1,881,360)	-
Maximum tumour diameter [median (range)], cm	10.6 (5.1-15.8)	-
Focality		
Unifocal	57	61.3
Multifocal	7	7.5
Unknown	29	31.2
PRETEXT stage		
I	0	0.0
II	36	38.7
III	23	24.7
IV	3	3.2
Unknown	31	33.3
Rupture		
Yes	7	7.5
No	56	60.2
Unknown	30	32.3
Metastasis		
Yes	9	9.7
No	55	59.1
Unknown	29	31.2
Portal vein thrombosis		
Yes	1	1.1
No	63	67.7
Unknown	29	31.2
Hepatic vein thrombosis		
Yes	0	0.0
No	64	68.8
Unknown	29	31.2
Primary resection		
Yes	16	17.2
No	77	82.8
Neoadjuvant chemotherapy		
Yes [n, median (range)]	52, 2.5 (1-8)	55.9
No	41	44.1
Preoperative TACE, cycles		
Yes [n, median (range)]	55, 2 (1-7)	59.1
No	38	40.9
POSTTEXT ^a stage (n=77)		
I	4	5.2
II	36	46.8
III	9	11.7
IV	1	1.3
Unknown	27	35.1

^aSixteen children underwent primary tumour resection (with no neoadjuvant chemotherapy and no preoperative TACE), and did not need to undergo POST-TEXT stage evaluation.

Abbreviations: AFP, alpha-fetoprotein; PRETEXT, pre-treatment extent of disease system; TACE, transarterial chemoembolisation; POST-TEXT, post-treatment extent of disease system

TABLE 2: Surgical and pathological outcomes of patients managed for hepatoblastoma

Characteristics	Number or as shown	Proportion (%)
Liver resection		
Hemihepatectomy (left hepatectomy + right hepatectomy)	37	39.8
Wedge resection	17	18.3
Trisectionectomy (left trisectionectomy + right trisectionectomy)	13	14.0
Bisegmentectomy (left lateral sectionectomy)	9	9.7
Central hepatectomy	2	2.2
Others	15	16.1
Operative time [median (range)], minutes	290 (100-510)	-
Estimated blood loss [median (range)], ml	100 (20-1000)	-
Volume of red blood cells transfused [median (range)], units	1 (0-6)	-
Pathologic subtype	93	
Epithelial variants	24	25.8
Pure foetal variant with low mitotic activity	3	-
Foetal variant, mitotically active	10	-
Unspecified	11	-
Epithelial mixed	11	11.8
Mixed epithelial and mesenchymal	41	44.1
Without teratoid features	2	-
With teratoid features	15	-
Unspecified	24	-
Unknown	17	18.3
Surgical margin		
Positive	7	7.5
Negative	69	74.2
Unknown	17	18.3
Microvascular involvement		
Yes	12	12.9
No	43	46.2
Unknown	38	40.9
Lymph node status (n=31)		
Positive	0	0.0
Negative	31	100.0
Postoperative chemotherapy		
Yes [n, median (range)]	63, 6 (1-12)	67.7
No	27	29.0
Unknown	3	3.2
Outcomes		
Survived without relapse	84	90.3
Survived with relapse	5	5.4
Died from relapse	4	4.3
Median follow-up duration [median (range)], months	30.5 (0.7-105.1)	-

TABLE 3: Detailed information of patients who experienced tumour relapse or death

^anull, unknown; -, no need to fill in; ^bW, wedge resection; LH, left hepatectomy; B,

Characteristics	Patients ^a								
	P1	P2	P3	P4	P5	P6	P7	P8	P9
Age, months	7	24	9	5	19	41	46	6	87
Gender	Female	Female	Male	Male	Male	Female	Female	Male	Male
PRETEXT stage	II	III	II	II	IV	II	Null	Null	II
Multifocal tumour	No	No	No	No	Yes	No	Null	Null	No
Metastasis	Yes	Yes	No	Yes	Yes	No	Null	Null	No
Neoadjuvant chemotherapy, cycles	0	2	0	8	4	4	2	0	0
Preoperative TACE	0	2	3	0	7	1	4	5	0
POSTTEXT stage	-	III	II	II	III	II	Null	Null	-
Surgical resection ^b	W	W	B	LH+W	LH	W	RT	Null	W
Surgical margin status ^c	N ⁻	P ⁺	N ⁻	N ⁻	P ⁺	N ⁻	Null	Null	N ⁻
Postoperative pathologic subtype ^d	Foetal	With TF	EV	MEM	EV	EM	Null	Null	EM
Postoperative chemotherapy, cycles	3	0	6	4	2	Null	Null	4	4
Relapse	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Time from diagnosis to relapse, months	0.7	3.3	13.4	8.5	11.3	6.4	14.1	7.3	22.4
Further anti-tumour treatment ^e	C	Not appropriate	C	Gave up	C	S	TACE	C	S
Death	Yes	Yes	Yes	Yes	No	No	No	No	No
Time from diagnosis to death, months	7.7	3.6	21.4	14.8	-	-	-	-	-

bisegmentectomy; RT, right trisectionectomy; ^cN⁻, negative; P⁺, positive; ^dEV, epithelial variant; With TF, with teratoid features; MEM, mixed epithelial and mesenchymal; EM, epithelial mixed; ^eC, chemotherapy; TACE, transarterial chemoembolisation; S, surgery
P8 was operated at another institution.

Figures




Figure 1

Kaplan-Meier estimates of event-free survival probabilities.




Figure 2

Kaplan-Meier estimates of overall survival probabilities.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [20191211SupplementalTable1tyedited.pdf](#)