

# Cold application for pain and anxiety reduction following chest tube removal : A systematic review and meta-analysis

**Chia-Te Chen**

National Cheng Kung University Medical Center: National Cheng Kung University Hospital

**Heng-Hsin Tung** (✉ [shannontung719@gmail.com](mailto:shannontung719@gmail.com))

National Yang Ming Chiao Tung University <https://orcid.org/0000-0001-9843-9924>

**Yen-Chin Chen**

National Cheng Kung University Medical Center: National Cheng Kung University Hospital

**Jiun-Ling Wang**

National Cheng Kung University Medical Center: National Cheng Kung University Hospital

**Sheng-Han Tsai**

National Cheng Kung University Medical Center: National Cheng Kung University Hospital

**Yen-Ta Huang**

Taipei Veterans General Hospital

**Teh-Fu Hsu**

National Cheng Kung University Medical Center: National Cheng Kung University Hospital

---

**Research**

**Keywords:** chest tube removal, cold application, pain, anxiety, systematic review, meta-analysis

**Posted Date:** June 14th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-608013/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

**Background and Objective:** Data on the effects of cold application on reducing pain and anxiety after chest tube removal (CTR) are inconsistent. This study aimed to conduct a systematic review and meta-analysis to evaluate the effects of cold application on pain and anxiety reduction after CTR.

**Methods:** We searched six databases, including Embase, Ovid Medline, Cochrane Library, Scopus, the Index to Taiwan Periodical Literature System, and Airiti Library, to identify relevant articles up to the end of February 2021. We limited the language to English and Chinese and the design to randomized controlled trials (RCTs). All studies were reviewed by two independent investigators. The Cochrane Collaboration's tool was used to assess the risk of bias, and Review Manager 5.4 was used to conduct the meta-analysis.

**Results:** Ten RCTs with 623 participants were included in the meta-analysis. The use of cold application could effectively reduce immediate pain and had persistent effects on pain after CTR. There were significant effects of cold application on reducing anxiety. The meta-regression showed that a drop in skin temperature to the 13°C target of cold application was significantly more effective for the immediate reduction in pain intensity compared with receiving up to 20 minutes target of cold application.

**Conclusion:** Cold application is a safe and easy-to-administer nonpharmacological method with immediate and persistent effects on pain and anxiety relief after CTR. In particular, skin temperature drops to the 13°C target of cold application were effective for immediate reduction of pain intensity following CTR.

## Introduction

A primary goal of chest tube removal (CTR) is the removal of tubes without the introduction of air or contaminants into the pleural space [1, 2]. The removal of a chest drain is a painful and frightening experience; however, most guidelines regarding CTR do not mention this. The act of removing the chest tube often causes pulling of the endothelial tissue adhered to the chest tube and stimulates intercostal nerves and inflamed pleura, all of which can cause pain [3-5]. A study conducted by Al-Otaibi et al.[6] found that the procedure of removing a chest tube not only causes moderate to severe pain but also increases anxiety. The source of anxiety may be related to fear and worry about the pain caused by the upcoming chest tube removal procedure.[6]

Currently, analgesics, including opioid and nonopioid drugs, are routinely prescribed to patients who are about to undergo CTR to alleviate pain during the procedure [6]. However, the use of analgesics may cause side effects such as hypoventilation, nausea, allergic reactions, and gastrointestinal bleeding [7]. Nonpharmacological interventions for pain related to chest tube removal, including music therapy, aromatherapy, and cold application, have been investigated [4, 7, 8]. Few studies have focused on the effects of music therapy and aromatherapy for reducing pain after CTR.

Cold application is a common nonpharmacological intervention used to relieve the pain caused by CTR [9]. Some studies have shown that cold application can also decrease anxiety levels [4, 8]. The mechanism of cold application mainly involves a slowing of the metabolism in skin tissue, an acceleration of nerve conduction and an increase the pain threshold, thereby reducing pain [10, 11]. Lee et al. [12] proposed that there was a residual effect of cold application on nerve velocity that could last 30 minutes after the removal of ice. However, the effectiveness of cold application for pain and anxiety level reductions are inconclusive. In addition, the chest tube management guideline does not mention pharmacological or nonpharmacological pain control for chest tube removals [13, 14]. The aim of the present systematic review and meta-analysis was to evaluate recent RCTs that analyze the effects of cold application on pain and anxiety after CTR. Furthermore, we also explored the effective period of time and target temperature of cold application related to pain relief after CTR.

## Methods

### Search strategy

This study followed the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement [15] (Supplementary Table 1 shows the PRISMA checklist) to explore the efficacy of cold application on pain intensity and anxiety levels in adults after CTR. Six databases, including Embase, Ovid Medline, Cochrane Library, Scopus, the Index to Taiwan Periodical Literature System, and Airiti Library, were systematically searched to identify relevant articles published from 1975 to February 2021. Reference lists of relevant articles were reviewed to identify additional studies. Four main concepts, chest tube removal, cold application, pain, and anxiety search strategy used combination with controlled vocabulary (11 MeSH terms and 5 Emtree terms) and free-text terms (19 synonyms plus truncation symbols) (Supplementary Table 2 shows the search strategy). The study protocol was registered in the PROSPERO international prospective register of systematic review (CRD42021245482).

### Eligibility criteria

The inclusion criteria included (1) publications in either the Chinese or English language, (2) randomized controlled trials (3) adults (aged 18 years and over) who had at least one chest tube removed from the pleural or mediastinal spaces in which (4) the experimental group received cold application and usual care and (5) the control group received usual care (or no treatment), and (6) the study outcome must have included a measurement of pain or anxiety. The exclusion criteria were as follows: (1) the experimental and control groups received different analgesic dosages before CTR; or (2) the experimental group received two or more nonpharmacological interventions.

### Data extraction and quality assessment

Two reviewers (C. T. Chen and S. H. Tsai) independently extracted and assessed the quality of eligible articles using the Cochrane Collaboration's tool. The seven domains were rated as either low risk, unclear, or high risk [16]. Any disagreements were resolved by consensus or consultation with a third author (J. L.

Wang).

## Data synthesis and analysis

We used Review Manager 5.4 to analyze the effect of cold application on pain intensity and anxiety levels in adults after CTR. The mean difference (MD) and 95% confidence intervals (CIs) were used to analyze continuous outcomes of pain intensity and anxiety level. Heterogeneity was assessed using Cochran's Q statistic and I-squared [17]. If the  $I^2$  value > 50% or P value < 0.10 was defined as statistical heterogeneity, sensitivity analysis with meta-regression was conducted by the software Open MetaAnalyst with a random-effects model [18]. A funnel plot was applied to evaluate publication bias.

## Result

### Literature search

A total of 110 articles were retrieved from six electronic databases and websites, and 35 articles were removed due to duplications. After the title, abstract and keywords were reviewed, 52 articles were discarded as not being sufficiently relevant, and 5 articles were abstracts for conferences. A total of 16 full-text articles were left for consideration. Eight articles were excluded for the following reasons: non-RCTs (n=3), irrelevant outcomes (n=2), same study reported twice (n=1), study without any report of standard deviation and 95% confidence intervals (n=3), and high risk of bias (n=2). Finally, 10 articles met the inclusion criteria and were included in the qualitative synthesis (Fig 1).

### Methodological quality assessment

Among the 10 studies, 5 studies (50%) were rated as having a low risk of bias for random sequence generation because they were randomly employed using random number generators, random number tables, block randomization, eight-member block randomization, or computer programs [7, 8, 19-21], and 5 studies were rated as having an unclear risk of random sequence generation [3, 4, 22-24]. One study was rated as having an unclear risk of concealment because the author did not provide a detailed explanation [8], and 9 studies (90%) were rated as having high risks of allocation concealment [3, 4, 7, 19-24]. Two studies were rated as having unclear risks of blinding of participants or researchers [8, 23], and 8 studies (80%) were rated as having high risks of blinding of participants or researchers [3, 4, 7, 19-22, 24]. Because cold application is a nonpharmacologic therapy, it is difficult to blind both participants and researchers. Two studies were rated as low-risk for the blinding of the outcome assessments [8, 19], and 8 studies (80%) were rated as high-risk for the blinding of the outcome assessments [3, 4, 7, 20-24]. All studies (100%) were rated as low-risk for incomplete outcome data, selective reporting, and other biases. The results are depicted in Supplemental Figure 1.

### Study characteristics

The characteristics of the ten included RCTs are summarized in Table 1. Five intervention studies used cold application [8, 20-22, 24], and 5 intervention studies used cold application plus a standard drug treatment [3, 4, 7, 19, 23]. The intervention durations varied from 10 minutes to 20 minutes [3, 4, 8, 11, 20-23], and 2 studies showed that cold application applied to the skin dropped the skin temperature to 13.6°C before CTR [7, 24]. The cold application was applied to the area surrounding the chest tube with 9-inch [8], 7-cm [24], 5-cm [23], and 6-inch [22] diameters. The major outcomes included the pain intensity and anxiety level. Pain intensities were measured by using the vertical visual analog scale (VAS), Numeric Rating Scale (NRS), and McGill Melzack Pain Questionnaire (MPQ). Anxiety levels were measured by the State Trait Anxiety Invent (STAI) and Hamilton Anxiety Scale (HAS). No adverse events were reported from any study.

### Cold application and immediate pain intensity

Ten studies using the VAS or NRS measurement were included in the meta-analysis to assess pain intensity. Due to high heterogeneity ( $P < 0.00001$ ,  $I^2 = 83\%$ ), the random effects model was selected. The results showed that cold application significantly reduced immediate pain intensity after CTR (MD: 1.32, 95% CI = -1.77 to -0.87,  $P < 0.00001$ ) (Fig 2a).

### Meta-regression of time and temperature target of cold application

Due to the high heterogeneity and variety of cold applications, covariates for meta-regression were divided into time up to 20 minutes and skin temperature drops to the 13°C target of cold application. The results showed that skin temperature drops to the 13°C target of cold application were more effective for the immediate reduction of pain intensity, compared to receiving cold application over a duration up to 20 minutes after CTR (coefficient: -0.979, Std. error 0.376,  $P = 0.009$ ) (Fig 2b).

### Cold application and persistent effects (15 minutes) on pain intensity

Six studies were included in the meta-analysis, and the results revealed that cold applications had significantly persistent effects (15 minutes) on pain intensity (MD: -0.50, 95% CI = -0.95 to -0.05,  $P = 0.03$ ) after CTR (Fig 3).

### Cold application and anxiety level

Three studies were included in the meta-analysis to assess anxiety levels. The random effects model was selected due to high heterogeneity ( $P = 0.02$ ,  $I^2 = 76\%$ ). The results revealed that cold application significantly reduced anxiety levels after CTR (SMD: -0.74, 95% CI = -1.40 to -0.07,  $P = 0.03$ ) (Fig 4). Two studies showed that cold application did not significantly reduce immediate anxiety levels after CTR (SMD: -0.50, 95% CI = -1.41 to -0.41,  $P = 0.28$ ). Only one

study showed a significant difference between the anxiety scores of the cold application and control groups 20 minutes after CTR (SMD: -1.19, 95% CI = -1.74 to -0.64,  $P < 0.0001$ ) (Fig 4).

### Publication bias

Funnel plots were investigated through visual inspection to assess the publication bias of the included studies. The shape of the funnel plot showed evidence of obvious symmetry (Supplemental Figure 2).

## Discussion

Several randomized studies have discussed the effect of cold application on pain caused by CTR. However, owing to the small sample sizes of the various studies, different strategies of cold application, and different time pain evaluation points considered after CTR, no certain effect of cold application alleviation after CTR can be confirmed. This study is the first to use a meta-analysis to investigate the effect of cold application on pain caused by CTR. The results of the meta-analysis show that cold application can effectively reduce immediate pain intensity after CTR. This result is similar to those from the two studies conducted by Sajad et al. [20] and Ertug and Ulker [24], respectively. In addition to pain relief after CTR, cold application was also reported to reduce pain before arterial puncture and intramuscular injection, but few studies have been reported [25, 26]. Cold application can lower the skin tissue temperature, impede the sensory nerve response and nerve conduction speed, and produce anesthetic effects, resulting in analgesia [27].

Currently, there is no consensus on the strategy of cold application regarding pain relief after CTR. Our meta-regression showed that a skin temperature drop to a target of 13°C effectively and immediately reduced the pain intensity following CTR. A study by Lowitzsch et al.[28] found that applying cold application to the skin and causing its temperature to drop to 27°C led to a change in the nerve conduction velocity. When the skin temperature drops to 10°C–15°C, cell metabolism slows, and the anti-inflammatory effect is augmented [27, 29-31]. If the skin temperature drops to 13.6°C or the cold application lasts 20 minutes, an analgesic effect can then be achieved [12, 27, 32]. Based on the abovementioned research results and arguments, it is recommended to check whether the regional skin temperature has reached 13.6°C when cold application is applied as a method to reduce pain in patients undergoing CTR. If the equipment is not available for checking skin temperature, 20 minute of cold application is recommended before performing CTR [3, 20].

In terms of the effect of persistent cold application (15 minutes) on pain after CTR, the meta-analysis shows that there is a significant difference in the pain intensity between the experimental group and control group. This result is consistent with the findings of Mokadem et al. [4] Lee et al.[12] pointed out that even after the cold application is removed, a residual effect reducing the nerve conduction velocity remains, such that analgesia is maintained.

Regarding the effect of cold application on anxiety reduction among patients receiving CTR, the result shows a significant difference but high heterogeneity. We supposed that this may be related to different tools, varying time points of anxiety evaluation, and the limited number of 160 participants. The results should be explained with caution. Further studies are required to confirm the effects of cold application on reducing anxiety levels after CTR.

Among the papers included in this systematic literature study, nine out of the ten included studies did not mention whether there were any complications of wound infection or frostbite in the experimental groups in which patients received cold application. Only one study indicated that no side effects were noted [24]. According to a systematic literature review by Greenstein [27], if a cold application causes the skin temperature to drop to -5°C or cold is applied for > 30 minutes, temporary or permanent frostbite or nerve damage may occur. Therefore, before and during cold application, it is necessary to closely monitor patients' skin temperature and the application duration. In addition, for patients with a medical history of Raynaud's disease, rheumatoid arthritis, cryoglobulinemia, cold urticaria, arteriosclerosis, or vascular injury, cold application is not recommended owing to contraindication concerns [27, 33].

### Limitations

This study has some limitations. First, the search strategies were limited to two languages, Chinese and English, and we only included ten studies with small sample sizes, which may affect the precision of our outcomes. Second, various studies have used different cold application strategies combined with different types of medications; therefore, we could not evaluate the effectiveness of different types of cold application on pain intensity in detail, and we do not know if combinations with other relaxation methods help. Third, the initial subjective pain intensity of the participants in all studies was not equal, which may contribute to heterogeneity. Fourth, most of the studies lacked a good concealment and blinding method; therefore, the results should be interpreted with caution. Finally, most eligibility criteria included participants over the age of 18 who underwent cardiothoracic surgery and had at least one chest tube; therefore, the generalization of the results may not be suitable to pediatric or medical patients with CTR.

## Conclusion

The results from this meta-analysis revealed that cold application can efficiently reduce pain intensity after CTR. When the skin temperature dropped to the target of 13°C with cold application, the effect was a more immediate reduction in pain intensity following CTR. Cold application is an effective and safe, nonpharmacological strategy that is easy to administer by a healthcare worker. Cold application could be included in the standard nonpharmacological management for reducing pain during CTR.

## Declarations

### Ethical Approval and Consent to participate

Not applicable.

## Consent for publication

All authors have read and agreed to the submission of the manuscript.

## Availability of supporting data

All data used and analyzed during this study are included in this published article and additional files.

## Acknowledgements

This research was supported by National Cheng Kung University Hospital Center for Clinical Medicine Research (NCKUH- T11008016). We thank all the colleagues and experts who joined in the study that greatly assisted the research.

## Author Contributions:

CT Chen and SH Tsai were responsible for this study's design. CT Chen completed the analysis of the data and manuscript preparation. YC Chen and HH Tung were responsible for revising the manuscript. CT Chen, HH Tung, YC Chen, JL Wang, SH Tsai, YT Huang, and TF Hsu approved the final manuscript.

## Funding

This research was supported by National Cheng Kung University Hospital Center for Clinical Medicine Research (NCKUH- T11008016)

## Conflict of interest:

The authors declare no potential conflicts of interest in this study.

## References

1. Bruce EA, Howard RF, Franck LS. Chest drain removal pain and its management: a literature review. *J Clin Nurs*. 2006;15:145-54. <https://doi.org/10.1111/j.1365-2702.2006.01273.x>
2. Kirkwood P. Chest tube removal (perform). *AACN Procedure Manual for High Acuity, Progressive, and Critical Care-E-Book*. 2016:184
3. Payami MB, Daryei N, Mousavinasab N, et al. Effect of cold application in combination with Indomethacin suppository on chest tube removal pain in patients undergoing open heart surgery. *Iran J Nurs Midwifery Res*. 2014;19:77
4. Mokadem NME, Shimaa E, Ibraheem S. Cold application and breathing exercises to reduce pain and anxiety during chest tube removal. *Am J Nurs*. 2017;6:285-92. <https://doi.org/10.11648/j.ajns.20170604.12>
5. Göttgens KW, Siebenga J, Belgers EH, et al. Early removal of the chest tube after complete video-assisted thoracoscopic lobectomies. *Eur J Cardiothorac Surg*. 2011;39:575-8. <https://doi.org/10.1016/j.ejcts.2010.08.002>
6. Al-Otaibi R, Mokabel F, Al-Ghuneimy Y. The effect of cold application on pain and anxiety during chest tube removal. *J Am Sci*. 2013;9:13-23
7. Hasanzadeh F, Kashouk NM, Amini S, et al. The effect of cold application and lavender oil inhalation in cardiac surgery patients undergoing chest tube removal. *EXCLI J*. 2016;15:64. <https://doi.org/10.17179/excli2015-748>
8. Aktaş YY, Karabulut N. The use of cold therapy, music therapy and lidocaine spray for reducing pain and anxiety following chest tube removal. *Complement Ther Clin Pract*. 2019;34:179-84. <https://doi.org/10.1016/j.ctcp.2018.12.001>
9. Mohammadi N, Pooria A, Yarahmadi S, et al. Effects of cold application on chest tube removal pain in heart surgery patients. *Tanaffos*. 2018;17:29
10. Ernst E, Fialka V. Ice freezes pain? A review of the clinical effectiveness of analgesic cold therapy. *J Pain Symptom Manage*. 1994;9:56-9. [https://doi.org/10.1016/0885-3924\(94\)90150-3](https://doi.org/10.1016/0885-3924(94)90150-3)
11. Hsieh L-Y, Chen Y-R, Lu M-C. Efficacy of cold application on pain during chest tube removal: a randomized controlled trial: A CONSORT-compliant article. *Medicine*. 2017;96. <https://doi.org/10.1097/MD.00000000000008642>
12. Lee JM, Warren MP, Mason SM. Effects of ice on nerve conduction velocity. *Physiotherapy*. 1978;64:2-6. <https://doi.org/10.2522/ptj.20090131>.
13. Huggins JT, Carr SR, Woodward GA, et al. Thoracostomy tubes and catheters: Management and removal.
14. Laws D, Neville E, Duffy J. BTS guidelines for the insertion of a chest drain. *Thorax*. 2003;58 Suppl 2:ii53-9. [https://doi.org/10.1136/thorax.58.suppl\\_2.ii53](https://doi.org/10.1136/thorax.58.suppl_2.ii53)
15. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *PLoS Med*. 2021;18:e1003583. <https://doi.org/10.1371/journal.pmed.1003583>
16. Higgins J, Thomas J. *Cochrane handbook for systematic reviews of interventions (Version 6)*. 2019

17. Melsen WG, Bootsma MC, Rovers MM, et al. The effects of clinical and statistical heterogeneity on the predictive values of results from meta-analyses. *Clin Microbiol Infect.* 2014;20:123-9. <https://doi.org/10.1111/1469-0691.12494>
18. Wallace BC, Dahabreh IJ, Trikalinos TA, et al. Closing the gap between methodologists and end-users: R as a computational back-end. *J Stat Softw.* 2012;49:1-15. <https://doi.org/10.18637/jss.v049.i05>
19. Gorji MH, Nesami MB, Ayyasi M, et al. Comparison of ice packs application and relaxation therapy in pain reduction during chest tube removal following cardiac surgery. *Am J Med Sci.* 2014;6:19. <https://doi.org/10.4103/1947-2714.125857>
20. Yarahmadi S, Mohammadi N, Ardalan A, et al. The combined effects of cold therapy and music therapy on pain following chest tube removal among patients with cardiac bypass surgery. *Complement Ther Clin Pract.* 2018;31:71-5. <https://doi.org/10.1016/j.ctcp.2018.01.006>
21. Sajedi-Monfared Z, Rooddehghan Z, Haghani H, et al. Cold therapy and respiratory relaxation exercise on pain and anxiety related to chest tube removal: A clinical trial. *Iran J Nurs Midwifery Res.* 2021;26:54. [https://doi.org/10.4103/ijnmr.IJNMR\\_228\\_19](https://doi.org/10.4103/ijnmr.IJNMR_228_19)
22. Sauls J. The use of ice for pain associated with chest tube removal. *Pain Manag Nurs.* 2002;3:44-52. <https://doi.org/10.1053/jpmn.2002.123017>
23. Demir Y, Khorshid L. The effect of cold application in combination with standard analgesic administration on pain and anxiety during chest tube removal: a single-blinded, randomized, double-controlled study. *Pain Manag Nurs.* 2010;11:186-96. <https://doi.org/10.1016/j.pmn.2009.09.002>
24. Ertuğ N, Ülker S. The effect of cold application on pain due to chest tube removal. *J Clin Nurs.* 2012;21:784-90. <https://doi.org/10.1111/j.1365-2702.2011.03955.x>
25. Shirvani MA, Ganji Z. The influence of cold pack on labour pain relief and birth outcomes: a randomised controlled trial. *J Clin Nurs.* 2014;23:2473-9. <https://doi.org/10.1111/jocn.12413>
26. Haynes JM. Randomized controlled trial of cryoanalgesia (ice bag) to reduce pain associated with arterial puncture. *Respir Care.* 2015;60:1-5. <https://doi.org/10.4187/respcare.03312>
27. Greenstein G. Therapeutic efficacy of cold therapy after intraoral surgical procedures: a literature review. *J Periodontol.* 2007;78:790-800. <https://doi.org/10.1902/jop.2007.060319>
28. Lowitzsch K, Hopf H, Galland J. Changes of sensory conduction velocity and refractory periods with decreasing tissue temperature in man. *J. Neurol.* 1977;216:181-8. <https://doi.org/10.1007/BF00313619>
29. Hubbard TJ, Aronson SL, Denegar CR. Does cryotherapy hasten return to participation? A systematic review. *J Athl Train.* 2004;39:88
30. Mac Auley DC. Ice therapy: how good is the evidence? *Int J Sports Med.* 2001;22:379-84. <https://doi.org/10.1055/s-2001-15656>
31. Sapega A, Heppenstall R, Sokolow D, et al. The bioenergetics of preservation of limbs before replantation. The rationale for intermediate hypothermia. *JBJS.* 1988;70:1500-13
32. Bugaj R. The cooling, analgesic, and rewarming effects of ice massage on localized skin. *Phys Ther.* 1975;55:11-9. <https://doi.org/10.1093/ptj/55.1.11>
33. McMaster WC. A literary review on ice thera in injuries. *Am J Sports Med.* 1977;5:124-6. <https://doi.org/10.1177/036354657700500305>

## Table

Table 1. characteristics of included studies

| Author, year              | Sample (I/C)  | Age (mean±SD)   | Intervention Description (before CTR)   | Control Descriptio (before CTR)  |
|---------------------------|---|---|---|--|
| Sauls, 2002               | 50 (25/25)  | T:59.7±13.06,   | Ice pack 10 mins  | Water pac from 87° I to 89° F  |
| Demir and Khorshid, 2010  | 90 (30/30/30)   | T:53.4±14.04  | Cold application 4C° 20 mins plus SDT   | C <sup>1</sup> : warm applicatio: 18-24C° plus SDT, C <sup>2</sup> : SDT |
| Ertug and Ulker, 2012     | 140 (70/70)   | I:48.74±16.59, C:49.17±16.20  | Ice bag until skin temperature reached 13°C   | No interventio   |
| Payami et al., 2014       | 66 (32/34)  | I:57.31±7.8, C:60.18±6.2  | Cold pack 4C°20 mins plus SDT   | Room temperatur get pack plus SDT  |
| Gorji et al., 2014        | 90 (30/30/30)   | I:57.75±9.38, C:58.48±9.78  | I <sup>1</sup> :Cool gel packs until skin temperature reached 13° plus SDT<br>I <sup>2</sup> : Relaxation skill 5mins plus SDT  | SDT  |
| Hasanzadeh et al., 2016   | 80 (I <sup>1</sup> :20,I <sup>2</sup> :20,I <sup>3</sup> :20/C:20)  | T:54.3±10.9   | I <sup>1</sup> : Cooling gel pack until skin temperature reached 13°C plus SDT<br>I <sup>2</sup> :inhaled lavender oil 20 mins plus SDT<br>I <sup>3</sup> :Cold with cooling gel pack plus inhaled lavender oil 20mins plus SDT | SDT  |
| Mokadem et al., 2017      | 120 (I <sup>1</sup> :30,I <sup>2</sup> :30,I <sup>3</sup> :30/C:30) | I <sup>1</sup> :40.25±10.98I <sup>2</sup> :37.55±14.12,I <sup>3</sup> :39.15±10.99 C:41.25±10.7   | I <sup>1</sup> :Cool gel pack 15 mins plus SDT<br>I <sup>2</sup> :Breathing exercises plus SDT<br>I <sup>3</sup> :Cool get pack and breathing exercises plus SDT  | SDT  |
| Yarahmadi et al., 2018    | 174 (I <sup>1</sup> :43,I <sup>2</sup> :44,I <sup>3</sup> :43/C:44) | I <sup>1</sup> :58.9±8.3,I <sup>2</sup> :60±5.7,I <sup>3</sup> :59.4±6.3,C:56.8±9.2               | I <sup>1</sup> :Ice pack -5°C 20 mins<br>I <sup>2</sup> :Listening to music 30 mins<br>I <sup>3</sup> :Ice pack plus music  | No interventio   |
| Aktas and Karabulut, 2019 | 120 (I <sup>1</sup> :30,I <sup>2</sup> :30,I <sup>3</sup> :30/C:30) | I <sup>1</sup> :62.6±12.11,I <sup>2</sup> :64.13±9.59,I <sup>3</sup> :64.8±7.07,C:65.8±7.23       | I <sup>1</sup> :Cold gel packet 20 mins<br>I <sup>2</sup> :Music therapy 20 mins<br>I <sup>3</sup> :lidocaine spray   | No interventio   |
| Zahra et al., 2021        | 120 (I <sup>1</sup> :30,I <sup>2</sup> :30,I <sup>3</sup> :30/C:30) | I <sup>1</sup> :57.70±10.16,I <sup>2</sup> :59.43±11.78,I <sup>3</sup> :51.26±15.77,C:53.83±12.25 | I <sup>1</sup> :Ice bag 20 mins<br>I <sup>2</sup> :Respiratory exercise 20 mins<br>I <sup>3</sup> :Ice bag and respiratory exercise   | No interventio   |

I/C: intervention/control, T:toal, SD: standard deviation, VAS: Visual Analogue Scale, NRS: Numeric Rating Scale, CTR: chest tube removed, SF-MPQ: Short-form McGill Pain Questionnaire, STAI: State Trait Anxiety Invent, HAM-A: Hamilton Anxiety Scale, SDT: standard drug treatment,

## Figures






Figure 1

PRISMA diagram for the literature search and selection process.



(a)



(b)

Figure 2

(a) Mean difference in the effects of cold application on reducing pain intensity (Visual Analog Scale, VAS; Numerical Rating Scale, NRS) immediately after CTR compared with the control. (b) Bubble plots of covariate meta-regression between time up to 20 minutes and skin temperature drops to the 13°C target of cold application.




Figure 3

Mean difference in the effects of cold application persistent effect (15 minutes) on pain intensity after CTR compared with the control.




Figure 4

Standard mean difference in the effects of cold application on anxiety level after CTR compared with the control.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Coldapplicationsupplemental.docx](#)