

Maternal factors of newborn low birthweight in malaria endemic settings of Nanoro, rural Burkina Faso

Moussa Lingani (✉ lingani10@yahoo.fr)

Institut de Recherche en Sciences de la Sante <https://orcid.org/0000-0002-5651-2035>

Serge H. Zango

Institut de Recherche en Sciences de la Sante

Innocent Valéa

Institut de Recherche en Sciences de la Sante

Daniel Valia

Institut de Recherche en Sciences de la Sante

Maïmouna Sanou

Institut de Recherche en Sciences de la Sante

Sékou O. Samandoulougou

Quebec Heart and Lung Institute

Annie Robert

Universite Catholique de Louvain (UCL)

Halidou Tinto

Institut de Recherche en Sciences de la Sante

Michèle Dramaix

Universite Libre de Bruxelles Ecole de Sante Publique

Philippe Donnen

Universite Libre de Bruxelles Ecole de Sante Publique

Research article

Keywords: Low birthweight, malaria, intermittent preventive treatment, sulfadoxine-pyrimethamine, Burkina Faso

Posted Date: November 2nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-61707/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background. In sub-Saharan Africa, the intermittent preventive treatment of malaria in pregnancy using sulphadoxine pyrimethamine (IPTp-SP) strategy is recommended to limit malaria adverse effects on birth outcomes. Ten year after IPTp-SP was adopted in Burkina Faso, we assessed the magnitude and maternal factors of low birthweight (LBW) in Nanoro.

Methods. A secondary analysis of data from a cross-sectional study was carried out in women who gave birth at Nanoro peripheral health centers using a binary multivariate logistic regression. Maternal socio-demographic factors, gynecological history and relevant medical characteristics were evaluated to identify associated factors. A p-value less than 0.05 was considered statistically significant.

Results. Of 291 delivery records examined, 14 % of women received three or more doses of SP while 80% used bed nets the night before their admission for delivery. Malaria was detected in 36.1% and anemia in 52.9% of women. The average neonate birthweight was 2933 g and 12 % were born with a low birthweight. By multivariate analysis, first delivery (OR = 8.84, [95% CI: 3.72-21.01]), and being multiparous with history of stillbirth (OR = 5.03, [95% CI: 1.54-16.40]) were significantly associated with an increased risk of LBW.

Conclusion. LBW was still prevalent in rural Nanoro and the uptake of three or more doses of SP for the IPTp was low. In addition, to improving the coverage of the 3-dose IPTp-SP, it may be necessary to target interventions aiming to reduce maternal anemia such as adequate nutrients uptake to mitigate the issue of adverse birth outcomes.

1.background

Despite halving the global neonatal mortality rate from 36.6 to 18.0 per 1000 livebirths between 1990 and 2017 [1], an estimated 2.5 million neonates died in the world in 2018 and this situation threatens the achievement of the third sustainable development goal (SDG), to end preventable child deaths by 2030 [2]. Low birthweight (LBW) - a birthweight below 2500 grams - is an important risk factor of neonatal deaths [3,4], and is prevalent in sub-Saharan Africa although half of infant birthweight is not recorded [5]. Noteworthy, neonate who survive from low birthweight immediate consequences are still at higher risk of short and long term health consequences such as stunted growth [6], lower intelligence quotient (IQ, an IQ test score of 70 or below) [7,8] or adult-onset chronic conditions [9]. Regional data suggest that the global burden of LBW is heavily skewed towards sub-Saharan Africa and south Asia due to high risk of infections [3,10]. Maternal infections including malaria and sexually transmitted infections (STI) are important risk factors of low birthweight through intrauterine growth retardation (birthweight below the 10th percentile for the gestational age) or prematurity (birth before 37 weeks of gestational age) [11–18]. Maternal malaria causes 20% of LBW in malaria endemic area and 6% neonatal deaths [18]. Potentially, 100,000 infant lives could be saved each year if adequate malaria prevention measures were

implemented during pregnancy. In addition, 25% of all stillbirth occurring in sub Saharan Africa [19] and 40% of all premature births follow a maternal infection with malaria or STI [20,21].

Sulfadoxine-pyrimethamine (SP) is recommended to prevent malaria adverse effects in pregnancy in sub Saharan Africa through the intermittent preventive treatment of pregnant women (IPTp) [22]. The strategy entails the administration of a full treatment course of SP to pregnant women starting early from the second trimester and administered monthly until delivery [23]. Despite the overwhelming evidence supporting the use of SP, the spread of *P. falciparum* resistance combined to low IPTp-SP uptake however threatens the antimalarial effectiveness of the strategy [24].

In Burkina Faso, IPTp-SP was adopted in 2005 starting at the second trimester of pregnancy with the requirement of 3 doses of SP [25]. Seven years later, the magnitude of LBW did not decrease substantially (from 15.8% in 2005 to 13.4% in 2012) and since then no other evaluation was carried out although it is still the recommended strategy [26,27]. A reevaluation of the impact of the current IPTp strategy is needed to understand its utility and prevent the effects of non-efficacy. The main goal of this analysis was to measure the prevalence of term LBW and to identify associated risk factors in Nanoro ten years after IPTp-SP policy was adopted.

2. Methods

2.1. Study site: The study was conducted in Nanoro demographic and health surveillance system (HDSS) area, a sentinel site for the "Roll Back Malaria" initiative. It is situated in the centre-west region of Burkina Faso at about 90 km from Ouagadougou the capital city with a population of 63,000 inhabitants [28]. Malaria transmission is holo-endemic, with transmission peaking during the rainy season (June-October). The commonest malaria vectors are *Anopheles gambiae*, *A. funestus* and *A. arabiensis* and *P. falciparum* is the predominant malaria parasite. Malaria is the main reason for visiting health centres with a case-fatality rate between 5% and 30% [29].

2.2. Study design: A secondary data analysis was conducted from data originally collected to evaluate malaria infection among women who recently gave birth in Nanoro health centers through a cross sectional survey. Nursing mothers and their babies were enrolled into the study. Data on demographic, gyneco-obstetric characteristic and relevant medical history were collected from the mothers onto a case report form. The participant antenatal care (ANC) cards were reviewed and data extracted from September 2013 to March 2014.

2.3. Inclusion/exclusion criteria: Women were eligible if they were aged between 16 and 45 years and gave birth to a neonate within the last 24 hours from a singleton pregnancy with a minimum gestational age of 37 weeks (using last menstrual period or the Ballard score)[30], lived in Nanoro HDSS, and have signed an inform consent form. Mothers who delivered non-singleton babies or those not willing to participate were not eligible. Of the 323 women, 9.9% (32/323) were not included, due to reasons including 4 (1.2%) non-singleton pregnancies, 6 (1.8%) stillbirths, 3 (0.9%) neonate deaths in the first two hours of life, 2 (0.6%) very ill newborns that were referred before clinical examination, 5 (1.5%) missing

birthweight records, 3 (0.9%) concurrent participation to other study and 9 (2.3%) delivery before 37 weeks of gestational age. Finally, 291 (90.1%) women were included. In this analysis, term low birthweight, birthweight between 500 and 2500 grams from a gestational age over 37 weeks was considered the main outcome.

2.4 Sample size: At the time of policy change, term LBW proportion was 15.8% [27]. We hypothesized that the policy change would reduce this proportion to 10% within the ten-year period of implementation. The sample size was estimated to test the hypothesis, with a precision $i = 3.5\%$, a confident interval of 95 % ($\alpha = 0.05$, with a critical value $Z^2 = 1.96$), the minimum sample size required was $n = 282$ participants using the following formula $n = Z^2 * p * (1-p) / i^2$. This was a secondary analysis of database collected on 291 participants recruited to estimate malaria prevalence. Therefore, the available sample size was adequate to test the hypothesis.

2.5 Data collection procedures and variables collected. Data were collected from the health center birth registries, the antenatal care cards and by neonate examination. Information not available in these documents were collected by one-on-one person interviews using a standardized questionnaire. Neonate examination was conducted within 24 hours of delivery either in the health facility or by home visit by trained midwives. Variables collected for mothers are presented in table 1 and included age, occupation, education level, parity, history of miscarriage, history of stillbirth, the number of living children at the time of the survey, the use of insecticide treated net (ITN) the night before visiting the health facility for delivery, the number of ANC visits performed, knowledge of the recommended protocol of 3 SP received during the pregnancy. Neonate birthweights were measured using calibrated Seca® 384 electronic scale with 10 g resolution and a precision less than 10 grams (seca gmbh & co. kg, Germany). In addition, blood samples were obtained from finger pricks to measure maternal hemoglobin level using a portable spectrophotometer (HemoCue, Ängelholm, Sweden) and a malaria rapid diagnostic test was performed using SD-Bioline Malaria Antigen *Pf*[®] test strips (sensitivity 99.7% and specificity of 99.5 %) for the identification of *P. falciparum*.

Table 1. List of exploratory variables

No	Variable	Definition	Categories
<i>Socio-demographic's characteristics</i>			
1	Age	Mother age (years)	15 - 19 ; 20 - 34 ; and 35 - 42
2	Education level	Educational level	None, primary or more
3	Occupation	Mother's occupation	Unemployed, employed/self employed
4	Parity	Number of birth that a woman had after 28 weeks of gestation	1; 2 - 4; 5 or more
5	Stillbirth	Maternal history of stillbirth	Yes / No
6	Miscarriage	Maternal history of miscarriage	Yes / No
7	Kids	Number of living kids at the time of the survey	One or two; three or four; Five or more
<i>Attitude and knowledge regarding malaria prevention</i>			
8	Required_IPTp	Knowledge of the minimum required 3 IPTp-SP doses	Yes / No
<i>Use of malaria prevention measures and relevant medical conditions</i>			
9	ITN use	Use of ITN the previous night	Yes / No
10	IPTp-SP	Number of IPTp - SP doses received	1; 2; 3 or more
11	ANC	Number of ANC visits performed	Less than 4; 4 or more
12	Anemia	Anemia (Hemoglobin level < 11g/dL)	No / Yes
13	Malaria	Positive malaria rapid diagnostic test results	No / Yes
14	Birthweight	Weight of neonate at birth in grams	Low (<2500); Normal (>=2500)

ANC: antenatal care, ITN: Insecticide treated bed net, IPTp-SP: Intermittent preventive treatment with sulfadoxine-pyrimethamine, g/dL: Gram per deciliter

2.6. Data processing and analysis: data were entered into a REDCap database, cleaned and imported to Stata version 15 (StataCorp. 2017, TX, USA) for analysis. The uptake of IPTp-SP was grouped in one, two, and three or more doses. Anemia was defined as hemoglobin level < 11g/dL, malaria as a positive rapid diagnosis test result. Because ultrasound was not available, pregnancy duration was assessed using the last menstrual period (LMP) or the Ballard score obtained by neonate examination whenever there was uncertainty. A term low birthweight was defined as any birthweight from a singleton pregnancy of at least 37 weeks of gestational age that was below 2500 g [16,30]. Odds ratios (OR) of LBW and 95% confident intervals (95% CI) were calculated according to each maternal factor using univariate logistic regression. Adjusted odd ratios (aOR) and 95% CI were derived by a backward elimination regression of variables with a p-values < 0.20 and retention of variables with statistically significant p-values. For the multivariable analysis, variable age was not included due to its strong correlation with the parity. Because of a comparable proportion of LBW if more than one delivery, mothers were subsequently grouped in single and multiple deliveries. In addition, variable history of stillbirth was combined to parity (first

delivery, and multiple delivery with history of stillbirth). A p-value < 0.05 was considered statistically significant.

2.7. Ethical considerations: The study protocol was discussed with local health authorities and community leaders to obtain their assent. Permission was obtained from the national ethics committee of Burkina Faso by the principal investigator (CERS 018.6.078). Informed consent was obtained from all women prior to participation. All participants with anemia or positive malaria test were treated according to the national guidelines.

3. Results

3.1. Study participants' characteristics: Table 2 summarizes study participants' characteristics. In brief, of the 291 mothers included, the average age (\pm standard deviation) was 26 (\pm 6) years and most of them were unemployed (85.2%). The median parity stood at three births (interquartile range, 2-5 births) and the median number of living children was three. One third of participant were tested positive for peripheral malaria (with a lower percentage among bed net users 33% versus 41% among non-users although this difference was not statistically significant) and half study participants were anemic. More than 96% of women reported three as the minimum SP doses required during the course of a pregnancy contrasting with the low rate (14%) of those that effectively received at least three doses

Table 2: Socio-demographic, gynecologic-obstetric and relevant medical history at inclusion (n=291).

Characteristics	items	Percentage
<i>Age group (years)</i>	< 20	20.3
	20-34	67.7
	>=35	12.0
<i>History of stillbirth</i>	No	89.3
	Yes	10.7
<i>History of miscarriage</i>	No	84.5
	Yes	15.5
<i>Parity</i>	1	22.7
	2 - 4	49.8
	5 or more	27.5
<i>Anemia</i>	No	47.1
	Yes	52.9
<i>Malaria</i>	Negative	63.9
	Positive	36.1
<i>ITN use</i>	Yes	81.6
	No	18.4
<i>Number of ANC visits performed</i>	≤ 1	5.2
	2	17.5
	≥ 3	77.3
<i>Knowledge of required SP doses</i>	≥ 3	96.6
	< 3	3.4
<i>Number of IPTp-SP doses received</i>	≤ 1	31.3
	2	54.6
	≥ 3	14.1
<i>Education level</i>	None,	83.8
	Primary or more	16.2
<i>Mother occupation</i>	Unemployed	85.2
	Employed/self employed	14.8

Abbreviation. ANC: antenatal care, SP: sulfadoxine-pyrimethamine,

3.2. Low birthweight prevalence and maternal factors: Of the 291 neonates, the mean birth weight (\pm standard deviation) was 2933 ± 390 g and 12.0 % (35/291) were born with a birthweight below 2500 g. This proportion was higher among teenage mothers (28.8%). **Table 3** presents the prevalence of LBW in relation to maternal sociodemographic characteristics, obstetrical history as well as the univariate odds ratios. Younger maternal age, and being primigravid or a multigravida with history of stillbirth were positively associated with the risk of low birthweight, while sleeping under bed net was significantly associated with a lower prevalence. Uptake of more than 3 doses of SP was not significantly association with LBW despite the downward trend of LBW prevalence observed as the uptake of SP increased. The prevalence of low birthweight was higher among anemic mothers although this association was not statistically significant ($p=0.052$). However, the risk of anemia was significantly increased mother with malaria compare to those without malaria (66.7% versus 44.2%, $p = 0.01$)

Table 3: Low birthweight prevalence and maternal factors, Burkina Faso 2013-14 (n=286)

Characteristics	Total	LBW (%)	OR	95%CI	p. value
<i>Age group (years)</i>					
< 20	59	28.8	4.0	1.9-8.5	<0.001*
20 - 34	197	9.2	1.0	ref	
≥ 35	35	0.0	-	-	
<i>Educational level</i>					
None	244	11.5	0.7	0.3-1.8	0.509
Primary or more	47	14.9	1.0	Ref	
<i>Occupation</i>					
Housewives	248	12.5	1.0	Ref	0.539
Others	43	9.3	0.7	0.3-2.1	
<i>Parity</i>					
Primipara	61	29.5	8.8	3.7 - 21.0	<0.001*
Multipara with history of stillbirth	26	19.2	5.0	1.5 - 16.4	
Multipara	199	4.5	1.0	Ref	
<i>History of miscarriage</i>					
No	246	11.4	1.0	Ref	0.443
Yes	45	15.6	1.4	0.6-3.5	
<i>Anemia</i>					
No	137	8.0	1.0	Ref	0.052
Yes	154	15.6	2.1	0.9-4.5	
<i>Malaria</i>					
No	186	10.8	1.0	Ref	0.378
Yes	105	14.3	1.4	0.7-2.8	
<i>ITN use</i>					
yes	227	9.7	1.0	Ref	0.018*
No	51	21.6	2.6	1.2-5.7	
<i>Number of ANC visits</i>					
≤1	15	13.3	1.2	0.3-5.5	0.902
2	51	13.7	1.2	0.5-3.0	
≥3	225	11.5	1.0	Ref	
<i>Number of SP doses received</i>					
≤ 1	91	14.3	1.5	0.5-5.1	0.703
2	159	11.3	1.2	0.4-3.7	
≥3	41	9.8	1.0	Ref	

Legend: *statistically significant

Abbreviations. SP: sulfadoxine-pyrimethamine; CI: confidence interval; ANC: antenatal care; LBW: low birthweight; Ref: reference group, ITN: insecticide treated bed nets,

The multivariate analysis showed that first delivery (OR = 8.83, [95% CI: 3.71-21.01]), and subsequent deliveries when there is an obstetrical history of stillbirth (OR = 5.03, [95% CI: 1.54-16.40]) were significantly associated with an increased risk of LBW.

4. Discussion

Low birthweight prevalence was still high in Nanoro particularly given that low birthweight from preterm births were not counted. However, the observed prevalence was 3.8 percentage points lower than the 15,8 % reported among term neonates at the start of the new strategy in the same region [27]. Lower prevalence than that reported in Nanoro was observed in other African countries with a similar malaria patterns and comparable intervention packages for antenatal care services. Indeed, in Ethiopia (10%) in 2014 [31], and Nigeria (7%) in 2013 [32] the reported prevalence was lower than that of the current

evaluation. However these estimates could be understated as in the Nigerian's study, data were collected retrospectively from the routine surveillance system and it is reported that half of delivery data in Africa is not recorded due to weaknesses in the reporting system [5]. Similarly, the study in Ethiopia was carried out in an urban area where the economic level of women tend to be higher than those from the rural area and the impact of better economic status on the prevalence of low birthweight is already described [33,34]. In contrast, rate in excess were reported in South Africa (38.5%) in 2014 [35], and in a study conducted in Zimbabwe (16.7%) [36]. The reason behind this higher prevalence in the South African and Zimbabwean studies could be related to the higher prevalence of human immuno-deficiency virus (HIV) infection [37]. HIV is reported in many African studies as a possible risk factor of LBW although the mechanism has not yet been fully elucidated [36,38]. This illustrates the importance of testing HIV when evaluating the impact of interventions allocated to pregnant women.

It is reported that the maternal obstetrical history and events occurring during the course of the pregnancy impact the outcome of pregnancies [39]. However, the specific factors reducing the birthweight are not always identified. Our results showed that combined parity and obstetrical history of stillbirth were significantly associated with the occurrence of low birthweight in rural Burkina Faso. Similarly to Hoquet *et al.* report, our study showed that the risk of LBW significantly increased at the first delivery [38]. The impact of primiparity on low birthweight was reported by several authors [40–42]. The biological ground for first delivery children to be at higher risk of low birthweight may be related to the placenta of first pregnancy women being at higher risk of parasite-infected red blood cells sequestration, disrupting thus nutrients supply to the fetus [43] or because some physiologic changes occurring during pregnancy to facilitate fetal growth with the process incompletely reversing in postpartum may create a more efficient environment for subsequent fetuses [44]. Indeed, uteroplacental blood flow, which is responsible for delivering oxygen and nutrients to the fetus, is greater during subsequent pregnancies compared to the first pregnancy [45,46]. In addition, there may be structural factors that limit the uterine capacity in the first pregnancy as described in animal models [47]. Furthermore, infants born from first pregnancy may have developed in a different maternal immune environment, contributing to relative growth restriction, compared to subsequent pregnancies [48]. Along with primiparity, mother with multiple pregnancy and an obstetrical history of stillbirths also presented a higher risk of delivering low birthweight neonates compared to multiple pregnancy without history of stillbirths. Similar observation was reported by Chen *et al.* in 2018 and Rozi *et al.* in 2016 [49,50]. Ahrens *et al.* have noticed that the risk of low birth weight was highest after history of three stillbirths [51]. Particular attention to pregnant women with such background characteristics during their antenatal care visits could be of greater impact in mitigating the risk of low birthweight among them.

Poor maternal nutritional status is also described as a key factor of poor intrauterine fetal growth [52]. Even though that characteristic was not assessed in the current study, the high prevalence of anemia could indicate a poor nutritional status [53]. Indeed, a study conducted in the same site showed that low maternal weights, low maternal body mass indexes and low brachial perimeters were significantly associated with anemia [48]. In the current study, anemia was positively associated with an increased risk of low neonate weights at univariate analysis ($p = 0.052$) however that trend weakened with adjustment

to parity by multivariate analysis and also because of reduced sample size secondary to missing values. Further assessment building on the scope of this finding may be necessary to show the adequate impact of anemia and poor nutritional status on the risk of the occurrence of LBW.

Studies reported that using bed net throughout pregnancy reduces the risk of adverse delivery outcomes including low birthweight [54]. A similar trend was observed by univariate analysis in the current study ($p=0.018$) which showed bed net use to efficaciously prevent adverse birth outcomes by reducing up to 11% of malaria infections. However, this could not be confirmed by multivariate analysis because of the reduced sample size due to missing values. Studies building on a larger sample size may be required to assess the actual impact of ITN use on the reduction of LBW

A number of limitations in this study are worth noting. The study was conducted on women who have delivered at term excluding low birthweight from preterm deliveries and that could understate the actual extent of the problem. It remains to bear in mind that gestational age record was collected from parental assessment of last menstrual period (LMP), which could be prone to some errors, as majority of the study population did not recall their LMP. In addition, SP is not indicated during the first trimester of pregnancy and malaria infection may have already occurred at the start of its administration from the second trimester. Although the analysis had limitations, it is worthy identifying the population that need special attention during antenatal care. Larger studies building on the scope of these factors examined in this report will be valuable in guiding efforts to curtail the impact of low birthweight in sub-Saharan Africa.

5. Conclusions

This report showed that low birthweight prevalence remained high among neonates in rural Nanoro although its magnitude has considerable decreased. It also showed that the intake of IPTp-SP was still low among the women. Improving the coverage of IPTp combined to interventions aiming to reduce poor maternal nutritional status may be key to successfully reduce the magnitude of LBW in such settings.

List Of Abbreviations

SP: sulfadoxine pyrimethamine, **IPTp**: intermittent preventive treatment in pregnancy, **LBW**: low birthweight, **ANC**: antenatal care, **OR**: odds ratio, **CI**: confident interval, **SDG**: sustainable development goals, **NMR**: neonatal mortality rate, **SGA**: small-for-gestational-age, **HDSS**: health and demographic surveillance system, **HIV**: human immunodeficiency virus, **WHO**: world health organization, **LMP**: late menstrual period,

Declarations

Acknowledgement

We thank the study participants and the health staff of Nanoro Health District.

Funding

The field work in this project was supported by the Clinical Research Unit of Nanoro, Burkina Faso. ML and HSZ are supported by a PhD scholarship from the Belgian Universities Cooperation for the Development (<https://www.ares-ac.be/fr/cooperation-au-developpement>). The funding bodies had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

ML, DV, IV and HT have conducted the study, conceptualized and written large parts of the manuscript. ML, MWD, and AR performed the statistical work. MS, SOS AR, MWD and DP have made important comments to the manuscript. All authors read and approved the final manuscript.

Availability of data and materials

The dataset used and analyzed during the current study is available from the corresponding author.

Ethics approval, consent to participate

The study was approved by the national ethics committee of health science research of Burkina Faso (CERS-201-6-078)

Consent to publication

Not applicable

References

1. Hug L, Alexander M, You D, Alkema L. National, regional, and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: a systematic analysis. *Lancet Glob Heal*. 2019;7:e710–20.
2. WHO-newborns: reducing mortality. 2019. <https://www.who.int/news-room/fact-sheets/detail/newborns-reducing-mortality>. Accessed 30 Oct 2019
3. Lawn JE, Blencowe H, Shefali O, You D, Lee AC, Waiswa P, et al. Every newborn: progress, priorities, and potential beyond survival. *Lancet*. 2014;384:189–205.
4. McCormick M c. The contribution of low birth weight to infant mortality and childhood morbidity. *N Engl J Med*. 1985;312:82–90.

5. UNICEF-Low birthweight. 2019. <https://data.unicef.org/topic/nutrition/low-birthweight/>. Accessed 23 May 2019
6. Christian P, Lee SE, Angel MD, Adair LS, Arifeen SE, Ashorn P, et al. Risk of childhood undernutrition related to small-for-gestational age and preterm birth in low- and middle-income countries. *Int J Epidemiol*. 2013;42:1340–55.
7. Gu H, Wang L, Liu L, Luo X, Wang J, Hou F, et al. A gradient relationship between low birth weight and IQ: a meta-analysis. *Sci Rep*. 2017;7:1–13.
8. Trahan L, Stuebing KK, Hiscock MK, Fletcher JM. The Flynn Effect: a meta-analysis. *Psychol Bull*. 2014;140:1332–60.
9. Jornayvaz FR, Vollenweider P, Bochud M, Mooser V, Waeber G, Marques-Vidal P. Low birth weight leads to obesity, diabetes and increased leptin levels in adults: the CoLaus study. *Cardiovasc Diabetol*. 2016;15:1–10.
10. Lee AC, Katz J, Blencowe H, Cousens S, Kozuki N, Vogel JP, et al. National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. *Lancet Glob Heal*. 2013;1:e26–36.
11. Wijesooriya NS, Rochat RW, Kamb ML, Turlapati P, Temmerman M, Broutet N, et al. Global burden of maternal and congenital syphilis in 2008 and 2012: a health systems modelling study. *Physiol Behav*. 2016;4:e525-33.
12. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population-representative data. *Lancet Glob Heal*. 2013;1:16–25.
13. Desai M, ter Kuile FO, Nosten F, McGready R, Asamoia K, Brabin B, et al. Epidemiology and burden of malaria in pregnancy. *Lancet Infect Dis*. 2007;7:93–104.
14. Zaba B, Calvert C, Marston M, Isingo R, Nakiyingi-Miiro J, Lutalo T, et al. Effect of HIV infection on pregnancy-related mortality in sub-Saharan Africa: secondary analyses of pooled community based data from the network for analysing longitudinal population-based HIV/AIDS data on africa (ALPHA). *Lancet*. 2013;381:1763–71.
15. Poespoprodjo JR, Fobia W, Kenangalem E, Lampah DA, Warikar N, Seal A, et al. Adverse pregnancy outcomes in an area where multidrug-resistant *Plasmodium vivax* and *Plasmodium falciparum* infections are endemic. *Clin Infect Dis*. 2008;46:1374–81.
16. Cutland CL, Lackritz EM, Mallett-Moore T, Bardaji A, Chandrasekaran R, Lahariya C, et al. Low birth weight: case definition & guidelines for data collection, analysis, and presentation of maternal immunization safety data. *Vaccine*. 2017;35:6492–500.
17. Jefferson KK. The bacterial etiology of preterm birth. *Adv Appl Microbiol*. Academic Press; 2012. p. 1–22.
18. Guyatt HL, Snow RW. Impact of malaria during pregnancy on low Birth weight in Sub-Saharan Africa. *Clin Microbiol Rev*. 2004;17:760–9.

19. Goldenberg RL, Thompson C. The infectious origins of stillbirth. *Am J Obstet Gynecol*. 2003;189:861–73.
20. Kemp MW. Preterm birth, intrauterine infection, and fetal inflammation. *Front Immunol*. 2014;5:574.
21. Agrawala V, Hirscha E. Intrauterine infection and preterm labor. *Semin Fetal Neonatal Med*. 2012;17:12–9.
22. WHO | updated WHO policy recommendation: intermittent preventive treatment of malaria in pregnancy using sulfadoxine-pyrimethamine (IPTp-SP). 2012. https://www.who.int/malaria/publications/atoz/who_iptp_sp_policy_recommendation/en/. Accessed 30 Oct 2019
23. WHO | Intermittent preventive treatment in pregnancy (IPTp). 2019. https://www.who.int/malaria/areas/preventive_therapies/pregnancy/en/. Accessed 4 May 2020.
24. van Eijk AM, Larsen DA, Kayentao K, Koshy G, Slaughter DEC, Roper C, et al. Effect of *Plasmodium falciparum* sulfadoxine-pyrimethamine resistance on the effectiveness of intermittent preventive therapy for malaria in pregnancy in Africa: a systematic review and meta-analysis. *Lancet Infect Dis*. 2019;19:546–56.
25. Gansané A, Nébié I, Soulama I, Tiono A, Diarra A, Konaté AT, et al. Change of antimalarial first-line treatment in Burkina Faso in 2005. *Bull la Soc Pathol Exot*. 2009;102:31–5.
26. He Z, Bishwajit G, Yaya S, Cheng Z, Zou D, Zhou Y. Prevalence of low birth weight and its association with maternal body weight status in selected countries in Africa: A cross-sectional study. *BMJ Open*. 2018;8:1–8.
27. Kabore P, Donner P, Dramaix-wilmet M. Obstetrical risk factors for lowbirth-weight in a rural Sahelian area. *Sante Publique (Paris)*. 2007;19:489–97.
28. Derra K, Rouamba E, Kazienga A, Ouedraogo S, Tahita MC, Sorgho H, et al. Profile: Nanoro Health and Demographic Surveillance System. *Int J Epidemiol*. 2012;41:1293–301.
29. WHO - Burkina Faso African region. 2019. https://www.who.int/malaria/publications/country-profiles/profile_bfa_en.pdf?ua=1. Accessed 23 May 2019.
30. Rosenberg RE, Ahmed ASMNU, Ahmed S, Saha SK, Chowdhury MAK, Black RE, et al. Determining gestational age in a low-resource setting: validity of last menstrual period. *J Heal Popul Nutr*. 2009;27:332–8.
31. Gebregzabihher Y, Haftu A, Weldemariam S, Gebrehiwet H. The prevalence and risk factors for low birth weight among term newborns in Adwa general hospital, northern Ethiopia. *Obstet Gynecol Int*. 2017;2017:7.
32. Dahlui M, Azahar N, Oche OM, Aziz NA. Risk factors for low birth weight in Nigeria: evidence from the 2013 Nigeria demographic and health survey. *Glob Health Action*. 2016;9:28822.
33. Kaur S, Ng CM, Badon SE, Jalil RA, Maykanathan D, Yim HS, et al. Risk factors for low birth weight among rural and urban Malaysian women. *BMC Public Health*. 2019;19:1–10.

34. Rezende Chrisman J, Mattos IE, Koifman RJ, Koifman S, Moraes Mello Boccolini P, Meyer A. Prevalence of very low birthweight, malformation, and low Apgar score among newborns in Brazil according to maternal urban or rural residence at birth. *J Obstet Gynaecol Res.* 2016;42:496–504.
35. Tshotetsi L, Dzikiti L, Hajison P, Feresu S. Maternal factors contributing to low birth weight deliveries in Tshwane district, South Africa. *PLoS One.* 2019;14:1–13.
36. Feresu SA, Harlow SD, Woelk GB. Risk factors for low birthweight in Zimbabwean women: a secondary data analysis. *PLoS One.* 2015;10:1–17.
37. Eaton JW, Rehle TM, Jooste S, Nkambule R, Kim AA, Mahy M, et al. Recent HIV prevalence trends among pregnant women and all women in sub-Saharan Africa. *Aids.* 2014;28:S507–14.
38. Hoque ME, Towbola OA, Mashamba TJ, Monokoane T. Comparison of adverse pregnancy outcome between teenage and adult women at a tertiary hospital in South Africa. *Biomed Res.* 2014;25:167–72.
39. Scrimshaw SC, Backes EP. Committee on assessing Health outcomes by birth settings A Consensus Study report. 2020. <https://doi.org/10.17226/25636>.
40. Bihoun B, Zango SH, Coulibaly MT, Rouamba T, Zemba D, Tahita MC, et al. Low birth weight and prematurity in teenage mothers in rural areas of Burkina Faso. *J Pregnancy Child Heal.* 2017;04:344.
41. Fall CHD, Osmond C, Sachdev HS, Sinha S, Restrepo-Mendez MC, Victora C, et al. Association between maternal age at childbirth and child and adult outcomes in the offspring: a prospective study in five low-income and middle-income countries (COHORTS collaboration). *Lancet Glob Heal.* 2015;3:e366–77.
42. Restrepo-Méndez MC, Lawlor DA, Horta BL, Matijasevich A, Santos IS, Menezes AMB, et al. The association of maternal age with birthweight and gestational age: a cross-cohort comparison. *Paediatr Perinat Epidemiol.* 2015;29:31–40.
43. Duffy PE. *Plasmodium* in the placenta: Parasites, parity, protection, prevention and possibly preeclampsia. *Parasitology.* 2007;134:1877–81.
44. Soma-Pillay P, Nelson-Piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr.* 2016;27:89–94.
45. Clapp JF, Capeless E. Cardiovascular function before, during, and after the first and subsequent pregnancies. *Am J Cardiol.* 1997;80:1469–73.
46. Prefumo F, Bhide A, Sairam S, Penna L, Hollis B, Thilaganathan B. Effect of parity on second-trimester uterine artery Doppler flow velocity and waveforms. *Ultrasound Obstet Gynecol.* 2004;23:46–9.
47. Igwebuike UM. A review of uterine structural modifications that influence conceptus implantation and development in sheep and goats. *Anim Reprod Sci.* 2009;112:1–7.
48. Moffett A, Hiby SE, Sharkey AM. The role of the maternal immune system in the regulation of human birthweight. *Philos Trans R Soc B Biol Sci.* 2015;370.

49. Rozi S, Butt ZA, Zahid N, Wasim S, Shafique K. Association of tobacco use and other determinants with pregnancy outcomes: a multicentre hospital-based case-control study in Karachi, Pakistan. *BMJ Open*. 2016;6:e012045.
50. Chen S, Yang Y, Qu Y, Zou Y, Zhu H, Yang H, et al. Both maternal and paternal risk factors for term singleton low birthweight infants in rural Chinese population: a population-based, retrospective cohort study. *Sci Rep*. 2018;8:1–7.
51. Ahrens KA, Rossen LM, Branum AM. Pregnancy loss history at first parity and selected adverse pregnancy outcomes. *Ann Epidemiol*. 2016;26:474-481.e9.
52. Han Z, Mulla S, Beyene J, Liao G, McDonald SD. Maternal underweight and the risk of preterm birth and low birth weight: A systematic review and meta-analyses. *Int J Epidemiol*. 2011;40:65–101.
53. Meda N, Coulibaly M, Nebie Y, Diallo I, Traore Y, Ouedraogo LT. Magnitude of maternal anaemia in rural Burkina Faso: contribution of nutritional factors and infectious diseases. *Adv Public Heal*. 2016;2016:1–7.
54. Gamble C, Ekwaru JP, ter Kuile FO. Insecticide-treated nets for preventing malaria in pregnancy. *Cochrane database Syst Rev*. 2009;38:34–5.