

Petri net Modeling and Property Analysis of C3 plant Photosynthesis and Photorespiration

Gajendra Pratap Singh (✉ gajendra@mail.jnu.ac.in)

Jawaharlal Nehru University <https://orcid.org/0000-0002-7642-2715>

Madhuri Jha

Jawaharlal Nehru University

Research article

Keywords: Petri net Modeling, C3 plant, Photosynthesis, Photorespiration

Posted Date: October 2nd, 2019

DOI: <https://doi.org/10.21203/rs.2.15608/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Metabolic pathway - Modeling and Analysis, is the emerging research area in Biochemistry over the past few years. C_3 plants are the best carbohydrate producers and also are the most common plants found in the nature. These plants are very much efficient in photosynthesis in cool and wet climate while, in very hot climatic condition it undergoes photorespiration process. In this paper we model the process of making food by C_3 plants to study the behavior in different climate conditions.

Background

In biochemistry a metabolic pathway is a combination of chemical reactions inside a cell. There are various chemical reactions which occurs in our body as well in the plants to release usable energy. These chemical reactions drive the processes that keep the living organism 'alive'. There are two types of metabolic pathways found, anabolic and catabolic pathways depending on their ability to either combines the molecules with the use of energy or splitting/breakdown of complex molecules by producing some energy in the process. The reactant, product and the intermediates may or may not be same. Sometimes these compounds act as enzymes to perform the certain reactions. Somewhere the product of one reaction acts as enzyme for other one. Some of the major metabolic networks in our body are glycolysis, citric acid cycle, urea cycle etc. Some metabolic pathways are present in the nature too such as photosynthesis and photorespiration. The food making process by plants is the most basic metabolic pathways found in environment but still have a greater importance. Increase in demand of food especially Carbohydrate rich food in India always encourage researchers to develop strategies to increase in food production. So, we are focusing on the carbohydrate giving plants and that is C_3 plants. There are three different type of plants occurs in nature namely C_3 , C_4 , and Crassulacean Acid Metabolism (CAM) plants, which named on the number of carbon present in the first stable compound formed during the photosynthesis process. Three photosynthesis pathways exist among terrestrial plants. Different plants have different networks depending on certain conditions [1,2]. Graph theory results are also available in several biological networks and in photosynthetic effect [3, 4]. Several other mathematical models are proposed in order to understand the network of plants photosynthesis and explain the stomatal behavior of leaf-level C_3 photosynthesis but they all are analytical model [5,6]. In this paper we propose a Descriptive mathematical model which approaches through graphical representation and draws some better qualitative and quantitative properties of any biological model or any complex system network. In order to understand the dynamic property and behavior of C_3 - plants Carboxylation process (Photosynthesis) and the Oxygenase process (Photorespiration), we propose a mathematical and graphical approach of modeling and that is Petri Net (PN). Due to its adaptability Petri net is very much efficient to model any biological system and can handle any concurrent and distributed complex system perfectly This model is Petri net (PN) which is a discrete event system first proposed by Carl Adam Petri in early sixties to model the concurrent, asynchronous, distributed system [7,8]. Petri net is also used in several other field of sciences determining binary vectors or can be used as a recommender system in

several networks [9,14]. Petri net has also application in biological networks like in Cardiovascular disease or in tuberculosis [15,18]

In this paper we focus on C_3 plants metabolic network and will compare the networks with C_4 plants using Petri net model. There are several software available to trace Petri net model and validate the system modeled and provide several results to explain the behavior of network considered. In this paper we are using PIPE v4.3.0 to draw the network and uses the results to explain the C_3 plant network [19]

Photosynthesis is the most common and most important biochemical reaction occurring in nature. Almost all types of plants undergo photosynthesis under different conditions. C_3 plants are the most common plants found in nature. Most of the carbohydrate producers are the C_3 plants for example Beans, Rice, Wheat, Potato etc, which are also called energy giving plants. The process in which these plants produce energy is Photosynthesis. This process has a network explained in Figure 1, as they are efficient at photosynthesis in cool and wet climates. C_3 plants do photosynthesis using NORMAL CALVIN CYCLE. Photosynthesis is the process where Reduced Nicotinamide Adenine Diphosphate (NADPH) and Adenosine Diphosphate (ATP) are created in CBB (Calvin Benson Bassham). In the suitable weather CO_2 enters through the tiny pores on leaves called **stomata**. This carbon dioxide then combines with an enzyme called RuBisCO to produce Sugar. This process produces a molecule made up of three carbon atoms, and that is why C_3 plants named. For C_3 plants, this process takes place in a chloroplast layer, the green cells present in plants which helps in photosynthesis which makes any plant green. The cycle continues and with the help of energy in the form of light from the sun, sugar is made as well as RuBP is also produced for the future use [20]. Sometimes the C_3 plant consumes oxygen too, which can slow down the process of photosynthesis. This process is called Photorespiration also called oxidative photosynthetic carbon cycle, or C_2 cycle. C_2 cycle is named so as the first stable compound formed is a two carbon molecule. In Figure 2, one can observe the network involved in producing energy in C_3 plants during Photorespiration. It is the process of taking O_2 instead of CO_2 in the process of Photosynthesis. Instead of fixing carbon when Rubisco fixes oxygen under certain conditions results in Photorespiration (PR). Sometimes it is also called anti-photosynthesis process. Under hot conditions RuBisCo has more affinity towards oxygen, so under hot weather when stomata closed so CO_2 can't diffuse in and O_2 can't diffuse out. Earlier it was said that this process is harmful for plants but recent researches shows the importance of this process. The process PR converts 2-phosphoglycolate (2PGP) into 3-phosphoglycerate (3PGA) and is escorted by O_2 intake and CO_2 release. PR plays the role of salvage or metabolic repair process which convert the toxic compound produced i.e., PGP into a useful compound PGA which produced during CBB cycle. PR also leads to the loss of CO_2 and NH_3 . Eventually, PR decreases the rate of photosynthesis by 30% in current atmospheric concentration of CO_2 and O_2 [21].

Results And Discussion

While modeling the process we are considering the stable intermediate products as places and the enzymes responsible for the reactions are taken as transitions. Tokens on the places are the number of

molecules available of the corresponding compounds. The weight of the arc is according to the number of molecules required in the reactions. Flow of tokens implies the formation of one product to another product according to the firing rule of Petri net. In Figure 3, we have shown the Petri net modeling of metabolic network in producing energy by C_3 plants in both conditions either involving light reaction or dark reaction. In Table 1, the description of places and transitions are shown according to the model.

Here we have modeled both the process of i.e., photosynthesis and photorespiration in a single model. When CO_2 is present in the stomata of the plant cell then the normal photosynthesis process will take place as t_1 will fire and further CO_2 combines with RuBp which is already present in the plant cells to form PGA then G3P followed by the formation of Glucose to produce energy for the plant.

Hence the path of firing of these transitions leads to normal photosynthesis $t_0, t_{11}, t_{12}, t_{13}, t_{14}, t_{15}, t_{16}$.

This cycle of model is validated from PIPE v4.3.0 software and some results follows which can be explained by the properties of Petri net.

- The model is *bounded*, signifies the compounds formed during photosynthesis is in limited number of molecules. There is no accumulation of any product formed during the process either the product has been used for the further process or being used by the plants growth.
- The number of molecules formed is different for different compound, so this model is not *Safe*.
- In this model we are not concerning after the formation of glucose so this model must have a *deadlock*. But the product RuBp is produced again during the process so the process is also *Live*.

In other case when there is no CO_2 enters through stomata then tokens on p_0 will be zero. Then t_1 attached by the inhibitor arc will fire to extract O_2 which is already present there in the cells of the plants and the tokens shifts to place (O_2) according to the weight associated with the preceding arcs to process the further reactions for photorespiration. The compound RuBp is already present in the plant, So to start the process we are taking the six molecules of RuBP i.e., $\mu^0(RuBp) = 6$, and for all other places in P , $\mu^0(p_i) = 0$.

Here, only transition t_1 will fire due to inhibitor arc property and consume O_2 to start the process. Now, O_2 combines with RuBp in the presence of RuBisCo enzyme to form two compounds PGA and PGP. PGA will again start normal Calvin cycle and PGP goes through a certain number of steps to form PGA back to follow Calvin cycle to produce Glucose.

Availability of sufficient amount of CO_2 results in Photosynthesis and insufficient amount of CO_2 results in photorespiration with availability of O_2 . In either case the Petri net is **non-terminating**. Also, the process is **Live** always. We can infer that the formation of glucose which is very essential for plants will never stop and the production of food given by C_3 plants can be increased by maintaining a suitable atmospheric composition. During modeling of metabolic pathways in photosynthesis and photorespiration, as the starting molecule is produced back to start the reactions again so the process is always **reversible** [25].

Conclusions And Scope

We have discussed the modeling of metabolic pathways as discrete-event systems. The benefit of modeling the bio-chemical process in Petri net graph is to analyze in a better way about the different stable compounds formed and the inter-conversion of one compound to another compound. The reachability graph makes a clear concept of formation of glucose from a certain available gases from air. C_3 plants like beans, rice, wheat, potatoes etc are the major plants for the survival of human being. One can model the Rice plant system or any other carbohydrate giving plants network. This will help to check how nitrogen can improve the productivity just by placing one more place in the model without affecting the basic properties of the model [26,27].

For any specific C_3 plant like Soya bean (a high source of Protein) which is a very important C_3 plant in nature, one can model the photosynthesis process for the plant in such a way to know the exact way in order to increase the production. Certain nanoparticles are also available by the addition which the productivity of soyabean plant or certain other C_3 plants can be increased. For examples Cerium Oxide (CeO_2NP_5) has the average size 19nm is a useful nanoparticles for C_3 plants. One can also model the other type of photosynthesis like C_3 and CAM plants photosynthesis network and can compare that by C_3 plants network. These plants are capable of resisting the photorespiration process by storing CO_2 in different layer. One can model for these plants that in the absence of CO_2 how these plants process the further reactions to produce Glucose. These days the modeling of C_4 pathways in the existing C_3 crops is very important by observing in these plants how RuBisCo not combines with O_2 can be modeled using Petri Net graph.

With modeling of all these networks, validation and simulation is also very important with Petri net software like PIPE or in MATLAB with Petri Net tool box.

Methods

- **Petri Net (PN)**

Petri net is a special kind of directed graph where two kinds of nodes are present, one is place and another is transition. While drawing the graphical representation of any model, places are drawn as a circle, and transitions as rectangle. It describes the complex system in graphical manner and gives an easy visualization of the system. Peterson describes further advantages of Petri net as a model of concurrent and conflict systems. The power of Petri net means the properties of the net can help to describe the modeled system in a better way. An advance Petri net analysis of techniques provides systematic and qualitative results of the modeled system. Petri net model analysis is drawing attention because of its simple graphical approach and certain high level Petri nets like timed Petri net, stochastic Petri net and colored Petri nets are proved very useful to study about any complex system. Petri net is 5-tuple set, consisting of a finite set of places, a finite set of transitions, a positive incidence function that denotes the number of arc from any transition to place, a negative incidence function that denotes the number of arc from any place to transition and an initial marking which represents the number of tokens on each place initially.

Mathematically represented as $PN = (P, T, I, I^+, \mu_0)$. Any transition fires at the given marking if and only if the outgoing arcs from all the places is less than and equal to the present marking of that place. After firing the tokens are moved from the input places and deposited on the output places accordingly.

The Reachability Graph of a Petri net is denoted by $R(PN, \mu)$ where, all the markings μ are taken as the vertices and the transition responsible for the corresponding marking is taken as edges. It helps in drawing the results of reachability of one state to another state. One state is reachable from another state if there is a directed connected path found between them [22,23].

- **Properties of Petri Net**

- **Reachability Graph/ Coverability Graph:** The graphical representation of the firing sequences and all possible marking states is called Reachability Graph of the Petri net. Here, nodes are the marking states and the arcs show the corresponding transition which fires to attain that state. In biological system the reachability graphs help to find the possible ways in the formation of the required product.
- **Boundedness:** A Petri net is Bounded if the number of tokens on any of the places involved never exceeds a finite or a fixed amount. In biological system it helps to imply that the number of molecules formed in the product compound is limited. Accumulation of any compound is not taking place. Hence it shows the property of conservation of mass.

- **Liveness:** A Petri net is live for μ (initial marking) , if there exists a firing sequence to reach to one marking from any other marking by a suitable firing sequence . in biological network it implies that the involved reactions will occurs repeatedly which contributes to the development depending on time.
- **Reversible:** A Petri Net is reversible if the initial marking is reachable from all the other reachable markings. In biological system it implies that the compound which starts the reaction is formed or produced again within the process
- **Extended Petri-net:** The extended Petri net is somehow an extension of standard Petri net where it has a form of arc called an inhibitor arc. An inhibitor arc is from a place p to a transition t which enables only when there are no tokens on p . This makes the concurrent phenomena much easier to model [24].

Abbreviations

RuBp: Ribulose 1,5-bisphosphate, RB: Ribulose 5-phosphate, RuBisCO: Ribulose-1,5-bisphosphate carboxylase Oxygenase, CO₂: Carbon dioxide, O₂: Oxygen, PGA: Phosphoglyceric acid, PGP:Phosphoglycolate phosphatase, G3P: Glyceraldehyde 3- Phosphate, H₂O₂:Hydogen peroxide, H₂O: Water, NH₃: Amonia Gas, NH₂: Amino redical, HPR:Hydroxypyruvate reductase, ATP: Adenosine triphosphate, ADP: Adenosine diphosphate, NADPH: Reduced Nicotinamide Adenine Diphosphate, PR: pyruvate.

Declarations

Acknowledgement

The authors thank Dr. Mamtesh Singh (Assistant professor , Gargi college) for validating syntax for reduced notations.

Authors' Contributions

GPS has modeled the process with helping in manuscript writing while MJ validated the model with its properties and wrote the paper. Final manuscript approval has been done by both the authors after reading the paper.

Funding

The corresponding author thanks to DST purse and UPOE-2,(Id No: 257) and a project under DST-SERB (Id No: ECR/2017/003480/PMS) for providing the research facility.

Availability of data and material

The software PIPE v4.3.0 used in the paper is easily available on the link <http://pipe.allapp.biz/downloaded.html>.

Ethics approval and consent to participate

Not applicable

Consent for Publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Author details

^{1,2}Mathematical Sciences and Interdisciplinary Research Lab, School of Computational and Integrative Sciences, Jawaharlal Nehru University, New Delhi-110067, India, E-mail: gajendra@mail.jnu.ac.in, jhamadhuri81@gmail.com

References

1. Yu H., Li X., Duchoud F., Chuang D.S. and Liao J.C.2018. Augmenting the Calvin –BensonBassham cycle by a synthetic malyl-CoA-glycerate carbon fixation pathway. Nature Communications volume 9, Article number: 2008 (2018).
2. Lawson, T., Kramer, D. M. and Raines, C. A. Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. Current Opinion in Biotechnology 23, 215U220” (2012).
3. Wang, Chuanli, et al. "Systematic comparison of C3 and C4 plants based on metabolic network analysis." BMC systems biology. Vol. 6. No. 2. BioMed Central, 2012.
4. Gao, Wei, et al. "Study of biological networks using graph theory." Saudi journal of biological sciences 25.6 (2018): 1212-1219.
5. Morgan, John A., and David Rhodes. "Mathematical modeling of plant metabolic pathways." Metabolic engineering 4.1 (2002): 80-89.
6. Bellasio, Chandra. "A generalised dynamic model of leaf-level C 3 photosynthesis combining light and dark reactions with stomatal behaviour." Photosynthesis research (2018): 1-20.
7. Brauer, Wilfried, and Wolfgang Reisig. "Carl adam Petri and “Petri nets”." Fundamental Concepts in Computer Science 3.5 (2009): 129-139.

8. Murata, Tadao. "Petri nets: Properties, analysis and applications." Proceedings of the IEEE 77.4 (1989): 541-580.
9. Kansal, S., Singh, G.P., Acharya, M. 2011. 1-Safe Petri nets generating every binary n-vector exactly once, *Scientiae Mathematicae Japonicae*, 74,(1), 29-36, e-2011, 127-134.
10. Kansal, S., Acharya, M. and Singh, G.P., Boolean Petri nets. In: *Petri nets - Manufacturing and Computer Science* (Ed.: Pawel Pawlewski), 381-406; Chapter 17. InTech Global Publisher, 2012, ISBN 978-953-51-0700-2.
11. Singh, G.P. and Kansal, S., "Basic results on Crisp Boolean Petri Nets", In *Modern Mathematical Method and High Performance in Science and Technology*, [Springer Proceedings in Mathematics & Statistics](#) Springer, Singapore, ISBN: 978-981-10-1453-6, 171 (2016), 83-88.
12. Sakshi Gupta, GajendraPratap Singh, SunitaKumawat. "Petri Net Recommender System to Model Metabolic Pathway of Polyhydroxyalkanoates" *International Journal of Knowledge and Systems Science*, 10(2), (2019): 42-59.
13. Gajendra Pratap Singh, Sujit Kumar Singh. "Petri Net Recommender System for Generating of Perfect Binary Tree" *International Journal of Knowledge and Systems Science*, 10(2), (2019), 1-12.
14. Singh, G.P., " Application of Petri Nets in Electrical, Electronics and Optimizations", IEEE Xplore, [International Conference on Electrical, Electronics, and Optimization Techniques \(ICEEOT\)](#), ISBN: 978-1-4673-9940-1 (2016) .
15. GajendraPratap Singh, Agraj Gupta, "A Petri Net Analysis to Study the Effects of Diabetes on Cardiovascular Diseases", IEEE Xplore, ISBN: 978-93-80544-36-6, (2019).

16. Gajendra Pratap Singh, Madhuri Jha, "Petri Net Modeling of Drug Resistance mechanism by First Line Drug in Tuberculosis", (In process)
17. Peterson, James L. "Petri nets." ACM Computing Surveys (CSUR) 9.3 (1977): 223-252.
18. Jensen, Kurt. "High-level Petri nets." Applications and theory of Petri nets. Springer, Berlin, Heidelberg, 1983. 166-180.
19. Bonet, Pere, et al. "PIPE v2. 5: A Petri net tool for performance modelling." Proc. 23rd Latin American Conference on Informatics (CLEI 2007). 2007.
20. Bauwe H. 1984. Photosynthetic enzyme activities and immunofluorescence studies on the localization of ribulose-1,5-bisphosphate carboxylase/oxygenase in leaves of C3, C4, and C3-C4 intermediate species of Flaveria (Asteraceae). Biochemie und Physiologie der Pflanzen 179, 253-268.
21. Lorimer, G.H. (1981) The carboxylation and oxygenation of ribulose 1,5 bisphosphate: the primary events in photosynthesis and photorespiration. Annu. Rev. Plant Physiol.
22. Murata T. Petri nets: Properties, analysis and applications. Proceedings of IEEE. 1989; 90:541-580. doi: 10.1109/5.24143
23. Pastor, Enric, et al. "Petri net analysis using boolean manipulation." International Conference on Application and Theory of Petri Nets. Springer, Berlin, Heidelberg, 1994.
24. Dugan, Joanne Bechta, et al. Extended Stochastic Petri Nets: Applications and Analysis. WISCONSIN UNIV-MADISON MOTOR BEHAVIOR LAB, 1984.
25. Reddy, Venkatramana N., Michael L. Mavrovouniotis, and Michael N. Liebman. "Petri net representations in metabolic pathways." ISMB. Vol. 93. 1993.

26. Lawson, T., Kramer, D. M. and Raines, C. A. Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. Current Opinion in Biotechnology 23, 215U220(2012).
27. Misra, B. B. , Chen, S. Advances in understanding CO₂ responsive plant metabolomes in the era of climate change. Metabolomics. 11, 1478U1491 (2015).

Tables

Table 1: Name and Description of Places and Transitions taken in the model in Figure 3

Places	Description	Transitions	Description
p ₀	CO ₂ entering from stomata	t ₀	Extract CO ₂
p ₁	CO ₂	t ₁	Extract O ₂
p ₂	O ₂	t ₂	RuBisCo performing Carboxylation
p ₃	RuBP	t ₃	RuBisCo as Oxygenase
p ₄	PGA	t ₄	Phosphoglycollate phosphatase enzyme
p ₅	PGP	t ₅	Carboxylation with NADPH to release NADP ⁺
p ₆	Glycolate	t ₆	Catalysation
p ₇	H ₂ O ₂	t ₇	Deanimation
p ₈	Glyoxylate	t ₉	Decarboxylase and methyltransferase
p ₉	H ₂ O	t ₁₀	Catalyzation in presence of ATP
p ₁₀	NH ₂	t ₁₁	Reduction in presence of phosphatase
p ₁₁	Glycine	t ₁₂	Reduction in the presence of triosephosphate dehydrogenase
p ₁₂	NH ₃	t ₁₃	Reduction and rearrangement in the presence of some metabolic enzymes
p ₁₃	Serine	t ₁₄	Reduction and isomerization in the presence of phosphatase
p ₁₄	HPR	t ₁₅	ATP + Phosphate pentose Kinase
p ₁₅	Glycerate	t ₁₆	Reduction and isomerization
p ₁₆	G3P		
p ₁₇	Fructose		
p ₁₈	RB		
p ₁₉	Glucose		

Figures




Figure 1

C3 photosynthesis




Figure 2

C3 photorespiration




Figure 3

PN modeling of C3Photosynthesis and Photorespiration