

Ideal Cardiovascular Health at Age 5-6 years and Cardiometabolic Outcomes in Preadolescence

Hester Jaspers Faijer - Westerink

Department of Public and Occupational Health, Amsterdam Public Health Research Institute, Amsterdam UMC, University of Amsterdam

Mette Stavnsbo

Western Norway University of applied Sciences, Department of Sport, Food and Natural Sciences

Barbara A. Hutten

Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Amsterdam Cardiovascular Sciences Research Institute, Amsterdam Public Health Research Institute, Amsterdam UMC, University of Amsterdam

Mai Chinapaw

8122-Netherlands School of Public and Occupational Health

Tanja GM Vrijkotte (✉ t.vrijkotte@amsterdamumc.nl)


Amsterdam Public Health Research Institute, Amsterdam UMC, University of Amsterdam, The Netherlands

<https://orcid.org/0000-0003-3641-4048>

Research

Keywords: children, ideal cardiovascular health, cardiovascular risk, epidemiology

DOI: <https://doi.org/10.21203/rs.3.rs-67772/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background- The American Heart Association (AHA) developed a definition of ideal cardiovascular health (ICH) based on the presence of both ideal health behaviours (diet, physical activity, weight status and smoking) and ideal health factors (glucose, total cholesterol and blood pressure levels). However, research of ICH in the paediatric population is scarce. We aimed to study ICH at age 5-6 years by extending the original ICH score with the health behaviours: sleep duration, screen time and prenatal smoke exposure, and to evaluate its association with cardiometabolic outcomes at age 11-12.

Methods- A total of 1,666 children aged 5-6 years were selected from the prospective ABCD-cohort study. Data on health behaviours and factors were used to calculate the ICH scores (original and extended) by adding the frequency of scoring 'healthy' on each indicator, based on international cut-offs. The children were followed up for 6 years and cardiometabolic outcomes (carotid intima-media thickness (CIMT), blood pressure, glucose and lipids) were measured. Associations between ICH (both original and extended) and cardiometabolic outcomes were examined using multivariable regression models.

Results- At age 5-6 years, 11% scored poor (score 1-5), 56% intermediate (score 6-7) and 33% good (score 8-9) on extended ICH. Healthy diet and normal total cholesterol concentrations were the least prevalent. Neither the original nor the extended ICH scores were associated with CIMT at age 11-12. A higher score on the extended ICH was associated with lower total cholesterol (p for trend <0.001), lower systolic (p for trend $=0.012$) and diastolic blood pressure (p for trend $=0.011$), and lower body mass index (BMI) ($p<0.001$) at age 11-12. The original ICH score was associated with lower total cholesterol ($p<0.001$) and BMI ($p<0.001$) only.

Conclusion- Our findings suggest that extending the ICH score in young children with additional health behaviours improves prediction of some cardiometabolic outcomes, but not CIMT in preadolescence, compared to the original ICH score. Further research with longer follow-up is needed to find out whether associations persist into adulthood.

Introduction

Cardiovascular diseases (CVDs) are the leading cause of death and remain one of the most important health challenges worldwide.¹ Although most children are born with optimal cardiovascular health, clustering of CVDs (or cardiometabolic) risk factors already occurs in childhood.² As both health behaviours as well as cardiometabolic risk factors track from childhood into adulthood, maintaining cardiovascular health in children is an important step to take.³ However, to develop successful strategies with special focus on cardiovascular health in children, the most important CVD risk factors in children need to be identified.

To define and measure cardiovascular health of individuals, the American Heart Association (AHA) created the construct of ideal cardiovascular health (ICH). The ICH construct consists of seven metrics: four health behaviours (not smoking, having a healthy weight status, being sufficiently physically active and eating a cardiovascular-healthy diet), and three health factors (healthy levels of total cholesterol, blood pressure and glucose).⁴

The construct of ICH is already used in population-based studies, especially focusing on adults. Reviews of these studies show that there is an inverse association between ideal ICH metrics and the incidence of CVDs.⁵⁻⁸ There is also evidence that even one point higher on the ICH metrics, gives a reduction of 19% in cardiovascular related mortality.⁷ Prevention strategies focusing on the factors included in the ICH metrics is thus highly recommended. However, studies on the prevalence of ICH in children are scarce⁹⁻¹²; only three have been conducted in children > 12 years of age.^{9,11,12} Further, few studies have investigated the relationship between ICH and cardiometabolic outcomes in populations under the age of 25, but these studies mainly focus on individual components.¹³⁻¹⁸

The definition of ICH in children differs slightly from the adult definition.⁴ Smoking is excluded in the ICH definition in children, as children younger than 12 years normally do not smoke. Also, recommended physical activity levels are higher for children: ≥ 60 minutes of moderate to vigorous physical activity per day.

Growing evidence shows that the health behaviours included in the ICH construct are not the only risk factors for CVD. In adults, excessive sedentary time and short sleep duration have i.e. shown to be risk factors too. The longitudinal evidence for health effects of sedentary behaviour is inconsistent, but the evidence for screen time alone is more convincing.^{19,20} As health behaviours lead to changes in health factors, adding more components to the ICH construct could make the construct more sensitive for the prediction of cardiometabolic outcomes. Thus we propose an extended version of the ICH definition for the purpose of this study by adding screen time, sleep duration, and prenatal smoke exposure. All three components have been associated with cardiovascular health in children.^{21,22}

The aim of our study was to apply this extended ICH score to a cohort of 5–6 year old children and evaluate its association with cardiometabolic outcomes at age 11–12 years. As children aged 11–12 years old do not yet experience cardiovascular diseases, we focused on the carotid intima-media thickness (CIMT), a surrogate marker for CVD, as our primary outcome. We compared the results with the original ICH definition.

Methods

Study population and design

This study was part of the Amsterdam Born Children and their Development (ABCD)-study, a prospective cohort study aiming at examining the associations between maternal lifestyle, health and psychosocial circumstances during pregnancy and the child's health at birth and later in life.²³ Between January 2003 and March 2004, 12,373 pregnant women were approached (> 99% of the target population) during their first prenatal visit to an obstetric care provider, of which 8,266 agreed to participate and filled out a pregnancy questionnaire.

Two weeks after their child's fifth birthday, mothers who gave permission for follow-up and whose addresses could be retrieved ($n = 6,161$) received a questionnaire on the health, development and behaviour of their child and an invitation for the child's health check. The questionnaire was returned for 4,488 children, of which 2,041 participated in the health check consisting of among other, a finger prick for blood collection after an overnight fast, measurements of height, weight, waist circumference and measurements of blood pressure. For those children who participated in the health check, mothers were also asked to fill in a validated food frequency questionnaire (FFQ).²⁴ Information on all ICH components was available from 1,666 children at age 5–6 years. A randomly selected subgroup of 1,081 children were approached for participation in follow-up health measurements at age 11–12 years (March 2015 to June 2016). The health check was similar to the health check at the age of 5, but ultrasound measurements of CIMT were added. From this group, 607 children participated at both time points and 559 children had information on the cardiovascular profile. Information from these 559 children were used to investigate the association between ICH at age 5–6 and cardiovascular health at age 11–12. The information of 459 children could be used for the association between ICH at age 5–6 and CIMT. Flowchart of the study population is presented in Fig. 1.

For inclusion in stage 1 of the current study the children had to meet the following criteria: 1) availability of information on maternal smoking status during pregnancy, 2) participating in the five year health check with cardiovascular measurements, 3) filled out FFQ, 4) filled out questionnaire on physical activity, sedentary behaviour and sleep duration. For inclusion in stage 2 of the study, children had to participate in the eleven year health check with cardiovascular measurements. Exclusion criteria's from participating in the study were: non-fasting blood samples, congenital heart malformations or use of medication altering blood pressure or lipid concentrations.

Demographic information

Ethnicity was based on the mother and her mother's country of birth, derived from the pregnancy questionnaire and categorized into Dutch, Turkey, Moroccan and Surinamese (the three most common non-Western ethnicities in the Netherlands). The remaining ethnicities were categorized into western and non-western ethnicities. Information on mother's and father's weight and height to calculate parent's body mass index (BMI) was derived from the questionnaire mother's received after their child's fifth birthday. Information on socioeconomic status (maternal education level) and cardiovascular diseases in the immediate family was also derived from this questionnaire. Maternal educational level was divided into low (primary school, technical secondary education, lower vocational secondary education), moderate (degree higher vocational secondary education, academic secondary education, intermediate vocational education) or high (degree higher vocational education, university). Cardiovascular diseases in the immediate family were answered by marking 'no', 'yes' or 'don't know' for the following diseases by the mother, the child's biological father and their first degree relatives: overweight, hypertension < 55 years, high cholesterol, stroke < 55 years, heart attack < 55 years and dead from heart attack < 55 years.²⁵ Information on the child's sexual maturation at the age of 11–12 was assessed by the Tanner scale, scored by the mother.²⁶

Health behaviours

Considered health behaviours were weight status, physical activity, diet, screen time, sleep duration and prenatal smoke exposure, apart from weight status all reported by the mother. Weight and height were measured to calculate BMI (weight in kg/height in cm²). Weight was measured to the nearest 100 g by using a Marsden weighing scale (model MS-4102, Oxfordshire, UK) and height to the nearest mm with a Leicester portable height measure (Seca, Hamburg, Germany). For defining metabolic syndrome, waist circumference was measured placing a Seca measuring tape around the abdomen between the costal border and iliac crest. Children aged 5–6 wore underwear only, children at age 11–12 wore sport outfits. Information on physical activity included both recreational and organized physical activity outside school hours as well as transport to and from school cycling. Hours of recreational physical activity was assessed by asking the hours of playing outside at schooldays and in weekends for both summer and winter. The total hours of organized sport activities per week and the total hours of cycling to and from school were added to a total physical activity score.²⁷

The FFQ that was used to evaluate the children's food consumption consisted of questions regarding the 71 most consumed food items by Dutch children.²³ For the present study, we included information regarding fruits and vegetable intake, fish intake, fibre intake and intake of sugar-sweetened beverages in line with the healthy diet definition of the AHA.⁴ The AHA guideline looks at the intake of fibre-rich whole grains, while the FFQ used in this present study measures total intake. Fibre intake was considered sufficient if it accounted 3 g/MJ, recommended by the Health council of the Netherlands.²⁸ Sodium intake could not be assessed with our FFQ. For other components besides fibre and sodium, the AHA cut-offs for a healthy diet were used: ≥ 450 g of fruits and vegetables/day, ≥ 2 servings of fish/week with a total weight of ≥ 200 g and ≤ 450 kcal of sugar-sweetened beverages/week based on a 2000 kcal diet. A comparison of our definition with the AHA definition can be found in supplementary table 2.

For the assessment of screen time, the number of hours per day children were watching TV, DVD or VIDEO and how many hours they played games on computer, XBOX or other types of videogames were asked. Hours of screen time for both schooldays and weekends were ranked in six categories from '(almost) never' to 'five or more hours per day'.²⁹ Information on children's sleep duration included the total hours of sleep per day, separately for schooldays and in the weekends. For all the above stated behaviours, a weighted weekly average was calculated. Mother's smoking habits during pregnancy were assessed from the pregnancy-questionnaire in which mothers scored the number of cigarettes smoked daily.

Health Factors

At age 5–6, plasma glucose, serum lipids and triglyceride concentrations were determined in capillary blood, collected after an overnight fast by a finger puncture by using a validated kit, developed for ambulatory purposes (Demecal: LabAnywhere,

Haarlem, The Netherlands).³⁰ At age 11–12, capillary blood was collected by finger puncture after 3 hours fasting and analysed by the point-of-care analyser Alere Cholestech LDX machine using Lipid Profile and GLU cassettes (Cholestech Alere Health Hayward, CA, USA).³¹ At both time points, blood pressure was measured by the Omron 705 IT (Omron Health Inc, Bannockburn, IL, USA)³² in sitting position with the arm supported at the heart level. After a test reading and fifteen minutes of rest, blood pressure was measured twice on the upper right arm with an appropriate cuff. If these measurements differed > 10 mmHg, a third measurement was performed.³³ The systolic and diastolic blood pressure were determined by averaging the two closest measurement.

Defining ICH at age 5–6 years

On each ICH component, children could score 'ideal' (score = 1) or 'non-ideal' (score = 0). To define ICH components as ideal, the AHA recommendation was used for the following components: BMI < 85th percentile according to the WHO growth reference³⁴, ideal diet when three out of four components were met,⁴ fasting plasma glucose < 5.6 mmol/L, total cholesterol < 4.40 mmol/L and blood pressure < 90th percentile based on reference values from the guideline of the American Academy of Pediatrics³⁵. Physical activity was considered ideal according to the AHA when children were physically active at moderate or vigorous intensity for at least 60 minutes per day. In our study the children of mother's who reported a total physical activity score of ≥ 7 hours per week were considered ideal.⁹

For the extended ICH score, a score on three additional health behaviours were added. There is no evidence-based cut-off for screen time. Based on a commonly used cut-off, we considered ≤ 2 hours of screen time per day as ideal.²⁹ Sleep duration was considered ideal if children slept ≥ 10 hours per night as recommended by the National Sleep Foundation.³⁶ Children from mothers who reported not having smoked during pregnancy received an ideal smoking score. The definition of each component according to AHA and the additions are listed in supplementary table 2.

The score on extended ICH ranged from zero to nine; zero to five points was considered poor, six and seven points intermediate and eight or nine points good ICH. This classification is made to compare characteristics of the most cardiovascular healthy population (high ICH-score), with the once with the fewest items on an ideal level.

Carotid intima-media thickness

Ultrasound measurements of the common carotid artery (CCA) were performed with the automated Panasonic Cardio Health Station V1.8 (Diagnostic Ultrasound System GM-72P00A). During this measurement, children were in lying position with their head at a 45 degree angle. CIMT was measured bilaterally at angles 150, 120 and 90 on the right CCA and at 210, 240 and 270 on the left CCA. The software automatically identified the region of interest and froze the image. Mean CIMT was calculated by averaging the measured angles, with a minimum of at least three angles. Out of 559 individuals, 459 individuals had a CIMT measurement with at least three angles, whereof 442 individuals had all six angles measured. Increased risk CIMT was defined as a CIMT of ≥ 90 th percentile of the cohort.

Cardiometabolic outcomes

Secondary outcomes were glucose, total cholesterol, high-density lipoprotein cholesterol (HDL-C), triglycerides (TG), systolic and diastolic blood pressure and BMI at the age of 11–12 (all continuous). The following outcome variables were dichotomized: low HDL-C (< 10th percentile), high TG (≥ 75 th percentile), (pre) hypertension (systolic and/or diastolic blood pressure ≥ 90 th percentile) and overweight (BMI ≥ 85 th percentile). Participants were considered having metabolic syndrome when they met three or more of the following criteria: waist circumference ≥ 75 th percentile, (pre) hypertension, low HDL-C, high TG and glucose ≥ 75 th percentile.³⁷ All percentiles in this paragraph are based on data of the own cohort.

Statistical analyses

We investigated the representativeness of the included subgroup by comparing the baseline characteristics between the non-included group (i.e. those who were approached at age 5–6, but did not participate in the questionnaire and/or health

check at age 5–6) and the included group (i.e. those who participated both in the questionnaire and health check at age 5–6) by an independent samples t-test for continuous variables and the χ^2 test for categorical variables.

Stage 1 - Differences in demographic characteristics between the poor, intermediate and good ICH groups (based on extended ICH score) were tested by one-way analysis of variance (ANOVA) for continuous variables and the χ^2 test for the categorical variables. The association of individual health behaviours, health factors and lipids with the extended ICH score at age 5–6 was evaluated by means of sex- and age adjusted multivariable linear regression models. *P*-for-trend was calculated considering ICH as a continuous variable.

Stage 2 - We studied the association between the extended ICH score at age 5–6 with CIMT and cardiometabolic outcomes at age 11–12 by means of multivariable linear regression models for continuous outcomes and multivariable logistic regression models for dichotomous outcomes. We considered the ICH variable as a continuous variable to calculate p-for-trend. For these analyses, all models included age, sex and maturation. To examine whether the association with respect to the dichotomous outcomes was present in children with a healthy weight at baseline, we repeated the logistic regression analysis excluding children with overweight at age 5–6.

All analyses were repeated with the original ICH score.

IBM SPSS Statistic software version 20.0 (IBM Corp, Armonk, NY, USA) was used for the analysis. A p-value of < 0.05 was considered statistically significant.

Results

Comparison of included and non-included children of the ABCD-cohort

Compared to those who were not included (N = 2,822), the included children (N = 1,666) were younger, had more often mothers with Dutch ethnicity and a higher educational level and reported more often family history of CVD. No differences were observed in parental weight status and boy-girl distribution. (Supplementary Table 1).

STAGE 1

Characteristics of the study subjects

Mean age (standard deviation (SD)) of the children at baseline was 5.7 (0.4) years. The prevalence of individual ideal ICH components were: normal glucose concentration (97.1%), no prenatal smoke exposure (92.3%), adequate physical activity (88.2%), healthy weight status (87.6%), normal blood pressure (86.9%), sufficient sleeping hours (81.4%), screen time < 2 hours (80.4%), normal total cholesterol concentration (69.7%) and healthy diet (2.0%). According to the extended ICH the majority of children had either intermediate (56%) or good (33%) ICH, which means respectively 6–7 or 8–9 ideal components. Compared to children with poor ICH (11%, 1–5 ideal components), children with a good ICH were more likely to have a Western ethnicity, higher educated mothers, normal weight parents and less frequent have a family history of CVD, see Table 1.

Table 2 shows the results of the associations between extended ICH scores and cardiovascular health outcomes at age 5–6 years. Children with higher scores had lower levels of health factors and higher percentages of ideal health behaviours.

Table 1
 Characteristics per ICH subgroup according to the extended ICH definition

N	All				P-value*
	1,666	Poor (1–5)	Intermediate (6–7)	Good (8–9)	
		188	934	544	
Age at first measurement (years), mean (SD)	5.7 (0.43)	5.7 (0.44)	5.7 (0.44)	5.6 (0.42)	0.055
Male gender, n (%)	846 (50.8)	101 (53.7)	471 (50.4)	274 (50.4)	0.692
Ethnicity	1210 (72.7)	88 (47.1)	677 (72.5)	445 (81.8)	< 0.001
Dutch, n (%)	26 (1.6)	13 (7.0)	11 (1.2)	2 (0.4)	< 0.001
Turkey, n (%)	59 (3.5)	16 (8.6)	39 (4.2)	4 (0.7)	< 0.001
Moroccan, n (%)	64 (3.8)	20 (10.7)	40 (4.3)	4 (0.7)	< 0.001
Surinamese, n (%)	205 (12.3)	10 (5.3)	118 (12.6)	77 (14.2)	
Other western, n (%)	101 (6.1)	40 (21.4)	49 (5.2)	12 (2.2)	
Other non-Western, n (%)	145 (8.7)	57 (31.0)	78 (8.4)	10 (1.8)	
Maternal educational level	318 (19.2)	56 (30.4)	194 (20.8)	68 (12.5)	
Low, n (%)	1195 (72.1)	71 (38.6)	659 (70.8)	465 (85.6)	
Middle, n (%)	792 (48.5)	60 (33.0)	434 (47.6)	298 (55.3)	
High, n (%)	622 (38.1)	75 (41.2)	361 (39.6)	186 (34.5)	
Parent weight status	218 (13.4)	47 (25.8)	116 (12.7)	55 (10.2)	
Normal weight, n (%)	829 (49.8)	68 (36.2)	470 (50.3)	291 (53.5)	
One overweight parent, n (%)	590 (35.4)	79 (42.0)	322 (34.5)	189 (34.7)	
Two overweight parents, n (%)	247 (14.8)	41 (21.8)	142 (15.2)	64 (11.8)	
Family history of CVD					
None, n (%)					
One parent, n (%)					
Two parents, n (%)					

Abbreviations: ICH, ideal cardiovascular health; n, number; SD, standard deviation; CVD, cardiovascular diseases.
 *Adjusted for age and sex.

Table 2

Association between extended ICH score and cardiovascular health outcomes at age 5-6 years

N, %	All									P for trend
	1,666	2	3	4	5	6	7	8	9	
		2 (0.1)	15 (0.9)	35 (2.1)	136 (8.2)	355 (21.3)	579 (34.8)	527 (31.6)	17 (1.0)	

Age at first measurement (years), mean (SD)	5.7 (0.4)	5.5 (0.2)	5.8 (0.5)	5.8 (0.4)	5.7 (0.4)	5.6 (0.4)	5.6 (0.4)	5.6 (0.4)	5.7 (0.5)	0.360
		0 (0.0)	3 (20.0)	21 (60.0)	77 (56.6)	188 (53.0)	283 (48.9)	263 (49.9)	11 (64.7)	0.055
Male gender, n (%)	846 (50.8)	4.6 (0.3)	4.8 (0.1)	4.9 (0.1)	4.7 (0.0)	4.6 (0.0)	4.6 (0.0)	4.5 (0.0)	4.4 (0.1)	< 0.001
Ideal health factors at age 5/6 years	4.6 (0.0)	5.4 (0.4)	4.4 (0.2)	4.3 (0.1)	4.2 (0.1)	4.3 (0.0)	4.1 (0.0)	3.7 (0.0)	3.6 (0.2)	< 0.001
Glucose (mmol/L), mean (SE)	4.0 (0.0)	98.4 (5.5)	105.7 (2.0)	103.7 (1.3)	101.1 (0.7)	98.4 (0.4)	96.8 (0.3)	95.3 (0.3)	92.7 (1.9)	< 0.001
Total cholesterol (mmol/L), mean (SE)	97.2 (0.2)	64.3 (5.0)	66.2 (1.8)	64.2 (1.2)	60.0 (0.6)	58.5 (0.4)	57.1 (0.3)	55.5 (0.3)	56.6 (1.7)	< 0.001
Systolic BP (mmHg), mean (SE)	57.4 (0.2)	1.4 (0.2)	1.4 (0.1)	1.4 (0.0)	1.3 (0.0)	1.3 (0.0)	1.3 (0.0)	1.2 (0.0)	1.3 (0.1)	< 0.001
Diastolic BP (mmHg), mean (SE)	1.3 (0.0)	0.8 (0.2)	0.7 (0.1)	0.8 (0.0)	0.7 (0.0)	0.6 (0.0)	0.6 (0.0)	0.6 (0.0)	0.6 (0.0)	< 0.001
HDL-C (mmol/L), mean (SE)	0.6 (0.0)	18.9 (0.9)	17.9 (0.3)	17.3 (0.2)	16.1 (0.1)	15.6 (0.1)	15.3 (0.1)	15.1 (0.1)	15.3 (0.3)	< 0.001
Triglycerides (mmol/L), mean (SE)	15.4 (0.0)	0 (0.0)	8 (53.3)	23 (65.7)	107 (78.7)	304 (85.6)	554 (95.7)	524 (99.4)	17 (100.0)	< 0.001
Ideal health behaviors 5/6y	1537 (92.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (0.5)	13 (2.5)	17 (100.0)	< 0.001
BMI, mean (SE)	33 (2.0)	1 (50.0)	7 (46.7)	21 (60.0)	90 (66.2)	286 (80.6)	521 (90.0)	526 (99.8)	17 (100.0)	< 0.001
Ideal smoking status, n (%)	1469 (88.2)	0 (0.0)	4 (26.7)	8 (22.9)	49 (36.0)	228 (64.2)	506 (87.4)	527 (100.0)	17 (100.0)	< 0.001
Ideal healthy diet score, n (%)	1339 (80.4)	0 (0.0)	2 (13.3)	14 (40.0)	60 (44.1)	244 (68.7)	493 (85.1)	526 (99.8)	17 (100.0)	< 0.001
Ideal physical activity, n (%)	1356 (81.4)									
Ideal screen time, n (%)										
Ideal sleep behavior, n (%)										

STAGE 2

Extended ICH score and carotid intima-media thickness

The mean CIMT at age 11–12 years was 0.462 mm (SE = 0.001). The association between extended ICH score at age 5–6 years and CIMT was not statistically significant (p for trend = 0.12; Table 3). Further, no statistical significant association was observed between extended ICH and the dichotomous variable ‘increased risk CIMT’ (odds ratio (OR) [95% confidence interval (95%CI)]: 1.04 [0.77–1.40], p = 0.80 (Table 4).

Extended ICH score and cardiometabolic outcomes

The extended ICH score at age 5–6 was significantly associated with total cholesterol (p for trend < 0.001), systolic blood pressure (p for trend = 0.01), diastolic blood pressure (p for trend = 0.01) and BMI (p for trend < 0.001) at age 11–12. No significant association was found for glucose (p for trend = 0.92), HDL-C (p for trend = 0.31) and triglycerides (p for trend = 0.47) (Table 3).

Table 4 shows OR (95%CI) for the extended ICH score and the prediction of cardiometabolic outcomes at age 11–12 years, adjusted for age-, sex- and maturation. Extended ICH was inversely associated with overweight (OR [95% CI]: 0.54 [0.43–0.68], p < 0.001). When children with overweight at age 5–6 were excluded, the association of ICH at age 5–6 and overweight at age 11–12 remained significant (OR: 0.66 [0.48–0.91], p = 0.01). No significant association was found between extended ICH score and metabolic syndrome, (pre-)hypertension, low HDL-C and high triglycerides.

Comparison between the extended and original ICH score

Overall the results were similar regardless of using the original or extended ICH score (Supplementary Table 3). However, the prospective results revealed no statistically significant association between the original ICH score and systolic- (p for trend = 0.056) and diastolic blood pressure (p for trend = 0.212) at age 11–12 years (Supplementary Table 4). The original ICH score was

inversely associated with overweight (OR [95%CI]: 0.52 [0.37–0.74], p < 0.001) but no longer significant when excluding children with overweight (OR [95%CI]: 0.84 [0.52–1.36], p = 0.47).

Table 3

Association between extended ICH score at age 5–6 years and cardiovascular health outcomes at age 11–12 years

N, %	All 559									P for trend
		2	3	4	5	6	7	8	9	
		2 (0.4)	4 (0.7)	6 (1.1)	38 (6.8)	119 (21.3)	198 (35.4)	187 (33.5)	5 (0.9)	
Age at first measurement (years), mean (SD)	5.7 (0.4)	5.5 (0.2)	5.7 (0.4)	5.7 (0.5)	5.8 (0.5)	5.7 (0.4)	5.7 (0.5)	5.7 (0.4)	5.6 (0.5)	0.943
Age at follow up (years), mean (SD)	11.8 (0.4)	12.1 (0.3)	11.4 (0.2)	12.0 (0.4)	11.9 (0.3)	11.7 (0.4)	11.8 (0.4)	11.8 (0.3)	11.5 (0.3)	0.011
Maturation, mean (SD)	4.9 (0.0)	4.5 (0.4)	4.6 (0.3)	4.8 (0.2)	5.2 (0.1)	4.9 (0.1)	4.9 (0.0)	4.9 (0.0)	4.6 (0.3)	0.107
Cardiovascular health 11/12y	4.1 (0.0)	5.0 (0.4)	4.5 (0.3)	4.1 (0.3)	4.1 (0.1)	4.3 (0.1)	4.1 (0.0)	3.9 (0.0)	3.5 (0.3)	0.920
Glucose (mmol/L), mean (SE)	105.2 (0.4)	110.8 (5.8)	110.6 (4.1)	113.0 (3.7)	106.8 (1.4)	105.2 (0.8)	105.1 (0.6)	104.2 (0.6)	109.3 (3.7)	<
Total cholesterol (mmol/L), mean (SE)	60.1 (0.3)	64.4 (4.6)	63.2 (3.2)	68.9 (2.9)	60.6 (1.1)	60.5 (0.6)	60.1 (0.5)	59.1 (0.5)	66.5 (2.9)	0.001
Systolic BP (mmHg), mean (SE)	1.0 (0.0)	1.0 (0.4)	0.8 (0.3)	1.2 (0.2)	1.1 (0.1)	1.0 (0.1)	0.9 (0.0)	1.0 (0.0)	0.5 (0.4)	0.123
Diastolic BP (mmHg), mean (SE)	17.6 (0.1)	20.3 (1.6)	20.2 (1.1)	22.1 (1.0)	18.2 (0.4)	18.4 (0.2)	17.2 (0.2)	17.2 (0.2)	17.1 (1.0)	
HDL-C (mmol/L), mean (SE)	0.462 (0.001)	0.445 (0.020)	0.485 (0.015)	0.429 (0.020)	0.455 (0.006)	0.458 (0.003)	0.462 (0.002)	0.464 (0.002)	0.457 (0.013)	
Triglyceriden (mmol/L), mean (SE)										
BMI, mean (SE)										
CIMT (mm), mean (SE)										

Abbreviations: N, number; SD, standard deviation; SE, standard error; BP, blood pressure; HDL-C, high-density lipoprotein cholesterol; BMI, body mass index; CIMT, carotid intima-media thickness.

Results are adjusted for age, sex and maturation. †There were no children with 0 or 1 ICH point.

Table 4
Odds ratios and 95% confidence intervals for the association between ICH score (extended and original) at age 5–6 years and cardiometabolic outcomes at age 11–12 years

	OR*	95% CI	P	OR*	95% CI	P
(Pre)Hypertension	0.85	0.66–1.09	0.203	0.89	0.62–1.28	0.526
Metabolic Syndrome	0.81	0.66–1.01	0.056	0.89	0.64–1.22	0.456
Low HDL-C	1.05	0.78–1.41	0.757	1.30	0.84–2.02	0.232
High triglycerides	1.00	0.82–1.23	0.940	1.02	0.77–1.36	0.879
Overweight	0.54	0.43–0.68	< 0.001	0.52	0.37–0.74	< 0.001
Increased risk CIMT	1.04	0.77–1.40	0.802	1.01	0.67–1.53	0.956

Abbreviations: HDL-C, high-density lipoprotein cholesterol; CIMT, carotid intima-media thickness;

OR, odds ratio; CI, confidence interval. * Adjusted for age, sex and maturation

Discussion

In this study, we applied an extended ICH score to a large cohort of children aged 5-6 years and investigated to what extent this score was related to cardiometabolic health at age 11-12 years. The results were compared to the original ICH score. To our knowledge this was the first time that additional components were added to the original ICH score to examine if this would improve the prediction of cardiometabolic health in children.

We found that ICH score at age 5-6 was not significantly associated with CIMT at age 11-12 according to both the extended and original ICH definition. Both definitions showed associations of ICH score at age 5-6 with total cholesterol and BMI at age 11-12, while associations with systolic- and diastolic blood pressure were only found when using the extended definition. Both ICH-scores predicted subsequent overweight. When restricted to normal weight children at age 5-6 only, the extended ICH score still predicted overweight at age 11-12 whereas this association was no longer significant according to the original definition.

Interpretation of findings

STAGE 1

Ideal health factors were more common than ideal health behaviours, but most children aged 5-6 years still scored ideal on healthy weight status (87.6%), sleeping hours (81.4%) and screen time (80.4%). Extra attention is needed for a healthy diet, since only 2% of the children met the healthy diet criteria. In line with our expectations, health factors were more often ideal than health behaviours as ideal behaviours precede changes in health factors. Other studies on ICH in child populations have shown similar findings.⁹⁻¹²

The low prevalence of a healthy diet is in line with other studies on ICH.¹⁰⁻¹² Another Dutch longitudinal study on the health of 8 year old Rotterdam born children found no children meeting Dutch dietary guidelines regarding intake of vegetables and sugar sweetened beverages.³⁸ As the Dutch dietary cut-offs were fairly similar to the AHA diet guideline³⁹, a similar percentage of ideal diet according to the AHA diet guideline can be expected. Dietary intake of both the Amsterdam and Rotterdam born children can thus be accounted as suboptimal.

In our study, the second most important ICH component was total cholesterol. We found 30.3% of 5-6 year old children with high total cholesterol concentrations. As lipoprotein concentrations change with growth and maturation (during puberty total cholesterol declines by 10-15%), it is questionable if the setpoint of cholesterol measurement at the age of 5-6 is representative for cardiovascular health.^{3,40} Though in studies of children >12 years of age, the prevalence of high total cholesterol was 21.5% to 66.8%^{9,11,12}, which indicate that our findings are similar to findings of studies that examine older children. Next, of the 559 children included in stage 2 of this study, 31.7% had high total cholesterol concentrations at age 5-6, which declined to 26.8% at age 11-12.

STAGE 2

Ideal cardiovascular health in childhood has shown to be associated with CIMT measurements in adulthood.⁹ Nevertheless, clustering of CVD risk factors starts in childhood and vascular changes may therefore also occur at younger ages. In our sample ICH score at age 5-6 was not related to CIMT measurements at follow-up six years later. The variation of CIMT in this group was small, which can be explained by the fact that we examined a relative young and healthy population of the same age. According to the systematic review of van den Munckhof et al.⁴¹ on CIMT in an apparently healthy adult population, 84% of CIMT can be explained by age. In contrast, several other studies in children found associations of ICH components and CIMT in childhood.^{14,15,37,42} The children included in these studies were mainly older at baseline (range 8 to 11) than our sample and CIMT was also measured at older ages (range 15 to 21)^{14,15,37} However, the study by Geerts et al.⁴² found an association between prenatal exposure to smoking and CIMT measurements at the age of 5 years old. The inconsistency of findings may be explained by the way in which CIMT was measured; in our study a hand guided ultrasound was used, which has a lower sensitivity than automatic devices. In addition, we performed measurements at three different angles in one segment (common carotid artery), while other studies performed measurements at one angle in three different segments (common carotid artery, internal carotid artery, carotid bulb). Lastly, different sonographers can have different findings and all studies used different readers. This inevitably makes results difficult to compare between studies.

The extended ICH definition at age 5-6 has stronger associations with systolic- and diastolic blood pressure and BMI at age 11-12 than the original definition and remained when testing only in non-overweight individuals aged 5-6 years. This difference cannot be explained by only one of the three additional components used in the extended definition. The difference is thus due to the clustering of additional components. This might be because of the bi-directional relationship between sleep duration and physical activity and sedentary behavior.⁴³ The implication for our study is that children with ideal levels of one of the three behaviours (sleep duration, physical activity or screen time) were more likely to have ideal levels of the other cardiovascular components, which resulted in a considerably higher ICH-level. As sleep duration, physical activity and screen time are linked, the extended definition gives a larger variance between ICH levels of those with and without ideal physical activity compared to the original ICH definition. Since there is a larger variation in ICH-level, associations of ICH-level with cardiometabolic outcomes are more likely. Furthermore, as unhealthy behaviour leads to changes in blood cholesterol concentrations which in turn leads to increased CIMT, it could be that adding more behaviours to the ICH construct will make the ICH construct more sensitive for predicting cardiovascular health.

The association between cardiometabolic outcomes and the additional components, namely: screen time, sleep behaviour and prenatal smoke exposure have been examined in previous studies.^{21,22,44} In children, associations between sedentary

behaviour and blood pressure are inconsistent²¹, although there is some evidence for a relationship between short sleep duration in children and increased blood pressure.²² Prenatal smoke exposure is not associated with an increased blood pressure in childhood according to a Portuguese study in 3-10 year old children (n ≈ 2500). However prenatal smoke exposure is associated with an increase in BMI (n = 17286).⁴⁴ Sleep duration in childhood is also found to be inversely associated with BMI²², while the evidence for sedentary behaviour is inconsistent.²¹ It is possible that the addition of the extra components single-handedly led to the extra association. However, it might also be that the combination of more non-ideal health behaviours is the basis of reduced cardiometabolic health which is associated with higher levels of blood pressure and BMI.

Strengths and limitations

Our study has some important strengths, including the longitudinal study design and its large sample size. Measurements took place at the same timepoint, by the same researchers with the same measuring instruments. Further, extensive measurements were done with the validated FFQ at stage 1 in the study.²⁴ The present study is the first to examine ICH in childhood with CIMT measurement in children as young as 11-12 years old.

The present study also has several limitations. First, our sample differed from the ABCD-cohort in terms of ethnicity, maternal educational level and family history of CVD. As supplementary Table 1 shows, precisely those participants are more likely to have a poor cardiovascular health. Therefore, our results probably overestimate the prevalence of ideal cardiovascular health in the ABCD-cohort. Not all children participating at age 5-6 also participated 5-6 years later, though the study cohort still consisted of 559 individuals. Information on health behaviours were reported by parents and thus all subjective. It is conceivable that parents perceive or present their child's behaviour in a more positive manner than the actual situation is. This could lead to an overestimation of healthy behaviour. Health factors were measured during the health check and automatic oscillometric devices were used for blood pressure measurements, while the blood pressure table of the American Academy of Paediatrics was made by auscultatory measurements.³⁵ Compared to blood pressure values measured by the auscultatory method, the oscillometric blood pressure device used in the present study found on average (SD) 4.6 (4.9) mmHg higher systolic blood pressure values and 3.3 (5.4) mmHg lower diastolic blood pressure values.³² This could have led to an overestimation of (pre-)hypertension because of higher levels of systolic blood pressure. However, a systematically overestimation of blood pressure values would not affect the present study. Many paediatric studies now use automatic devices in combination with the standard blood pressure tables⁴⁵, thus findings of this study are comparable with other available research. For outcome measures, our study was limited to CIMT measurement and CVD risk factors since cardiovascular events could not be studied as clinical outcome at this age.

Implications

Our study implicates that to raise a healthy generation, there is need for dietary interventions in Dutch children. Adherence to the healthy diet guidelines among the children was extremely low, which is unfortunately since dietary habits have shown to track into adulthood.⁴⁶

As the extended definition did not improve the prediction of CIMT and most cardiometabolic factors, we do not recommend the AHA to add the extra components to their definition yet. First there is need for further research.

In compliance with other studies,^{47,48} our study shows that children with a non-western ethnicity and lower educated mothers were more likely to have a poorer cardiovascular health, public health interventions regarding cardiovascular health should focus particularly on these at risk individuals.

Recommendations for future research

Further research is needed to gain insight into the associations between lifestyle of young children and cardiovascular health later in life. We recommend other researchers to look at ICH in children under 12 years in relation to cardiometabolic outcomes at a later age and explore the addition of sleep duration, screen time and prenatal smoke exposure to ICH on these outcomes. We also encourage other researchers to examine subpopulations (i.e. overweight children).

The ABCD-cohort will also be followed up longer and report on associations of ICH at age 5-6 and cardiometabolic health at an older age.

Conclusion

This study showed that 33% of children meet the criteria for an ideal cardiovascular health. A healthy diet was least prevalent among 5-year olds (2%). Extending the ICH score in young children with additional health behaviours improved the prediction of some cardiometabolic outcomes, but not CIMT in preadolescence, compared to the original ICH score.

Declarations

Acknowledgements

The authors thank all participating hospitals, obstetric clinics, general practitioners, Youth Health Care Centres and primary schools for their contribution to data collection for the ABCD study; we thank all the participating women and children for their valuable cooperation.

Authors Contribution

All the authors have participated in the study to a significant extent and have met the criteria for authorship as established by the International Committee of Medical Journal Editors. All authors have declared that no competing interests exist.

Data availability

The individual data are not available for a public repository for ethical reasons but can be made available to other researchers for purposes of reproducing results or for collaboration. Researchers wishing to apply for the data can contact the project leader of the ABCD-study (abcd@amsterdamumc.nl).

Ethical standards

Approval for the ABCD study was obtained from the Central Committee on Research involving Human Subjects in the Netherlands, the Medical Ethical Committees of the participating hospitals, and from the Registration Committee of the Municipality of Amsterdam. Written informed consent was obtained from all participating mothers.

Sources of funding

This part of the ABCD-study was financially supported by the Netherlands Organisation for Health Research and Development (Grants 21000076, 92003489, 40-00812-98-11010), Dutch Heart Foundation (grant 2007B103) and Sarphati Amsterdam.

Disclosures

Authors have nothing to disclose.

References

1. McAloon CJ, Boylan LM, Hamborg T, et al. The changing face of cardiovascular disease 2000–2012: An analysis of the world health organisation global health estimates data. *Int J Cardiol.* 2016;224:256–64. doi:10.1016/j.ijcard.2016.09.026.
2. Erald G, Erenson SB, Athanur S, et al. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. *N Engl J Med.* 1998;338(23):1650–1656. <http://www.nejm.org.ezproxy.library.wur.nl/doi/pdf/10.1056/NEJM199806043382302>. Accessed February 26, 2018.
3. Steinberger J, Daniels SR, Hagberg N, et al. Cardiovascular Health Promotion in Children: Challenges and Opportunities for 2020 and Beyond: A Scientific Statement From the American Heart Association. *Circulation.* 2016;September(12):236–55. doi:10.1161/CIR.0000000000000441. 134) .
4. Lloyd-Jones DM, Hong Y, Labarthe D, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: The american heart association's strategic impact goal through 2020 and beyond. *Circulation.* 2010;121(4):586–613. doi:10.1161/CIRCULATIONAHA.109.192703.
5. Aneni EC, Crippa A, Osondu CU, et al. Estimates of mortality benefit from ideal cardiovascular health metrics: A dose response meta-analysis. *J Am Heart Assoc.* 2017;6(12). doi:10.1161/JAHA.117.006904.
6. Maclagan LC, Tu JV. Using the concept of ideal cardiovascular health to measure population health: A review. *Curr Opin Cardiol.* 2015;30(5):518–24. doi:10.1097/HCO.0000000000000210.
7. Guo L, Zhang S. Association between ideal cardiovascular health metrics and risk of cardiovascular events or mortality: A meta-analysis of prospective studies. *Clin Cardiol.* 2017;(October):1339–46. doi:10.1002/clc.22836.
8. 10.1016/j.mayocp.2016.01.019
Younus A, Aneni EC, Spatz ES, et al. A Systematic Review of the Prevalence and Outcomes of Ideal Cardiovascular Health in US and Non-US Populations. *Mayo Clin Proc.* 2016;91(5):649–670. doi:10.1016/j.mayocp.2016.01.019.
9. Laitinen T, Laitinen TT, Pahkala K, et al. Ideal cardiovascular health in childhood and cardiometabolic outcomes in adulthood: The cardiovascular risk in young finns study. *Circulation.* 2012;125(16):1971–8. doi:10.1161/CIRCULATIONAHA.111.073585.
10. Dong H, Yan Y, Liu J, et al. Alarming trends in ideal cardiovascular health among children and adolescents in Beijing, China, 2004 to 2014. *Int J Cardiol.* 2017;231(March 2017):264–70. doi:10.1016/j.ijcard.2016.12.027.
11. Yang Q, Yuan K, Gregg EW, et al. Trends and Clustering of Cardiovascular Health Metrics Among U.S. Adolescents 1988–2010. 2014. doi:10.1016/j.jadohealth.2014.03.013.
12. González-Gil EM, Santabárbara J, Ruiz JR, et al. Ideal cardiovascular health and inflammation in European adolescents: The HELENA study. *Nutr Metab Cardiovasc Dis.* 2017;27(5):447–55. doi:10.1016/j.numecd.2016.12.003.
13. Hanvey AN, Mensah FK, Clifford SA, Wake M. Adolescent Cardiovascular Functional and Structural Outcomes of Growth Trajectories from Infancy: Prospective Community-Based Study. *Child Obes.* 2017;13(2):154–63. doi:10.1089/chi.2016.0263.
14. Menezes AMB, da Silva CTB, Wehrmeister FC, et al. Adiposity during adolescence and carotid intima-media thickness in adulthood: Results from the 1993 Pelotas Birth Cohort. *Atherosclerosis.* 2016;255:25–30. doi:10.1016/j.atherosclerosis.2016.10.026.
15. Yajnik CS, Katre PA, Joshi SM, et al. Higher glucose, insulin and insulin resistance (HOMA-IR) in childhood predict adverse cardiovascular risk in early adulthood: the Pune Children's Study. *Diabetologia.* 2015;58(7):1626–36. doi:10.1007/s00125-015-3602-z.

16. Geerts CC, Bots ML, Grobbee DE, Uiterwaal CSPM. Parental smoking and vascular damage in young adult offspring: is early life exposure critical? The atherosclerosis risk in young adults study. *Arterioscler Thromb Vasc Biol.* 2008;28(12):2296–302. doi:10.1161/ATVBAHA.108.173229.
17. Ayer JG, Belousova E, Harmer JA, David C, Marks GB, Celermajer DS. Maternal cigarette smoking is associated with reduced high-density lipoprotein cholesterol in healthy 8-year-old children. *Eur Heart J.* 2011;32(19):2446–53. doi:10.1093/eurheartj/ehr174.
18. Ried-Larsen M, Grontveld A, Moller NC, Larsen KT, Froberg K, Andersen L. Association between objectively measured physical activity intensity in childhood and measures of subclinical cardiovascular disease in adolescence: prospective observations from the European Heart Study. *Br J Sports Med.* 2014;48:1502–7. <http://ovidsp.ovid.com/athens/ovidweb.cgi?T=JS.&NEWS=N&PAGE=fulltext&AN=3676641&D=med2..>
19. Proper KI, Singh AS, Van Mechelen W, Chinapaw MJM. Sedentary Behaviors and Health Outcomes Among Adults A Systematic Review of Prospective Studies. *Am J Prev Med.* 2011;40(2):174–82. doi:10.1016/j.amepre.2010.10.015.
20. Covassin N, Singh P. Sleep Duration and Cardiovascular Disease Risk: Epidemiologic and Experimental Evidence. doi:10.1016/j.jsmc.2015.10.007.
21. Chinapaw MJM, Proper KI, Brug J, Van Mechelen W, Singh AS. Relationship between young peoples' sedentary behaviour and biomedical health indicators: a systematic review of prospective studies. *Obes Rev.* 2011;(12):621–32. doi:10.1111/j.1467-789X.2011.00865.x.
22. Matthews KA, Pantesco EJM. Sleep Characteristics and Cardiovascular Risk in Children and Adolescents: An Enumerative Review. doi:10.1016/j.sleep.2015.06.004.
23. Van Eijsden M, Vrijkotte TGM, Gemke RBJ, Van der Wal MF. Cohort profile: The Amsterdam born children and their development (ABCD) study. *Int J Epidemiol.* 2011;40(5):1176–86. doi:10.1093/ije/dyq128.
24. Dutman AE, Stafleu A, Kruizinga A, et al. Validation of an FFQ and options for data processing using the doubly labelled water method in children. *Public Health Nutr.* 2010;14(3):410–7. doi:10.1017/S1368980010002119.
25. Silberberg JS, Wlodarczyk J, Fryer J, Robertson R, Hensley MJ. Risk Associated with Various Definitions of Family History of Coronary Heart Disease The Newcastle Family History Study II. 1998;147(12). <https://academic.oup.com/aje/article-abstract/147/12/1133/93184>.
26. Petersen AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal status: Reliability, validity, and initial norms. *J Youth Adolesc.* 1988. doi:10.1007/BF01537962.
27. Burghard M, Knitel K, Van Oost I, Tremblay MS, Takken T. Is our youth cycling to health? results from the Netherlands' 2016 report card on physical activity for children and youth. *J Phys Act Heal.* 2016;13(11):218–24. doi:10.1123/jpah.2016-0299.
28. Health Council of the Netherlands. *Guideline for Dietary Fibre Intake*; 2006. https://www.gezondheidsraad.nl/sites/default/files/Dietary_fibre_intake_0.pdf. Accessed June 15, 2018.
29. Chinapaw MJM, Altenburg TM, Van Eijsden M, Gemke RBJ, Vrijkotte TGM. Screen time and cardiometabolic function in Dutch 5–6 year olds: Cross-sectional analysis of the ABCD-study. *BMC Public Health.* 2014;14(1):1–6. doi:10.1186/1471-2458-14-933.
30. Gootjes J, Tel RM, Bergkamp FJM, Gorgels JPMC. Laboratory evaluation of a novel capillary blood sampling device for measuring eight clinical chemistry parameters and HbA1c. *Clin Chim Acta.* 2009;401(1–2):152–7. doi:10.1016/J.CCA.2008.12.016.
31. Alere. *Alere Cholestech LDX Lipid Profile GLU Cassettes CLSI*; 2004. <https://www.alere.com/en/home/product-details/cholestech-ldx-system.html>.
32. Stergiou GS, Yiannes NG, Rarra VC. Validation of the Omron 705 IT oscillometric device for home blood pressure measurement in children and adolescents: The Arsakion School Study. *Blood Press Monit.* 2006;11(4):229–34. doi:10.1097/01.mbp.0000209074.38331.16.

33. van den Berg G, van Eijsden M, Galindo-Garre F, Vrijkotte TGGR, Reinoud JBJ, Gemke. Explaining socioeconomic inequalities in childhood blood pressure and prehypertension. The ABCD study. *Hypertension*. 2013;61(1):35–41.
34. WHO Multicentre Growth Reference Study Group. (2006). WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr Suppl.* (450):76–85.
35. Flynn J, Kaelber D, Baker-Smith C, et al. Clinical Practice Guideline for Screening and Management of High Blood Pressure in Children and Adolescents. *Pediatr Transplant*. 2017;140(3):e20171904.
36. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's updated sleep duration recommendations: final report. *SLEH*. 2015;1:233–43. doi:10.1016/j.sleh.2015.10.004.
37. Juonala M, Singh GR, Davison B, et al. Childhood metabolic syndrome, inflammation and carotid intima-media thickness. The Aboriginal Birth Cohort Study □. *Int J Cardiol*. 2016;203:32–6. doi:10.1016/j.ijcard.2015.10.073.
38. van der Velde LA, Nguyen AN, Schoufour JD, et al. Diet quality in childhood: the Generation R Study. *European Journal of Nutrition*. 2018.
39. *The Wheel of Five Guidelines [Dutch: Richtlijnen Schijf van Vijf]*. The Hague, the Netherlands; 2016. <https://www.voedingscentrum.nl/Assets/Uploads/voedingscentrum/Documents/Professionals/Schijf van Vijf/Voedingscentrum Richtlijnen Schijf van Vijf 2016 4.pdf>.
40. Balder JW, Lansberg PJ, Hof MH, Wiegman A, Hutten BA, Kuivenhoven JA. Pediatric lipid reference values in the general population: The Dutch lifelines cohort study. *J Clin Lipidol*. 2018;12:1208–16. doi:10.1016/j.jacl.2018.05.011.
41. Van Den Munckhof ICL, Jones H, Hopman MTE, et al. Relation between age and carotid artery intima-medial thickness: a systematic review: Relation between age and cIMT. 2018. doi:10.1002/clc.22934.
42. Geerts CC, Bots ML, Ent van der, CK, Grobbee DE, Uiterwaal C. Parental Smoking and Vascular Damage in Their 5-year-old Children. *Pediatrics*. 2012;129(1):45–54. <http://pediatrics.aappublications.org/content/pediatrics/129/1/45.full.pdf>. Accessed March 22, 2018.
43. Lin Y, Tremblay MS, Katzmarzyk PT, et al. Temporal and bi-directional associations between sleep duration and physical activity/sedentary time in children: An international comparison. *Prev Med (Baltim)*. 2018. doi:10.1016/j.ypmed.2017.12.006.
44. Li L, Peters H, Gama A, et al. Maternal smoking in pregnancy association with childhood adiposity and blood pressure. *Pediatr Obes*. 2016. doi:10.1111/ijpo.12046.
45. Parker ED, Kharbanda EO, Sinaiko AR. Are we measuring blood pressure correctly in children, particularly in obesity? *Arch Dis Child*. 2016;101(11):990–1. doi:10.1136/archdischild-2016-310682.
46. Craigie AM, Lake AA, Kelly SA, Adamson AJ, Mathers JC. Tracking of obesity-related behaviours from childhood to adulthood: A systematic review. *Maturitas*. 2011. doi:10.1016/j.maturitas.2011.08.005.
47. Rashid V, Engberink MF, Van Eijsden M, et al. Ethnicity and socioeconomic status are related to dietary patterns at age 5 in the Amsterdam born children and their development (ABCD) cohort. *BMC Public Health*. 2018;18:115. doi:10.1186/s12889-017-5014-0.
48. Dommelen van P, Schönbeck Y, Hirasing RA, Buuren van S. Call for early prevention: prevalence rates of overweight among Turkish and Moroccan children in The Netherlands. *Eur J Public Health*. 2015;25(5):828–33. doi:10.1093/eurpub/ckv051.

Figures




Figure 1

Flowchart of study population

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalmaterialICHinChildhood.docx](#)