

Cardiovascular dysautonomia and cognitive impairment in Lewy body disease

Hisayoshi Oka (✉ h.oka@jikei.ac.jp)

,Daisan Hospital, The Jikei University School of Medicine

Tadashi Umehara

Tokyo Jikeikai Ika Daigaku

Atsuo Nakahara

Tokyo Jikeikai Ika Daigaku

Hiromasa Matsuno

Tokyo Jikeikai Ika Daigaku

Research article

Keywords: Orthostatic hypotension, Daily blood pressure fluctuation, Cognitive decline, Parkinson's disease, Dementia with Lewy bodies

Posted Date: October 23rd, 2019

DOI: <https://doi.org/10.21203/rs.2.16373/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 18th, 2020. See the published version at <https://doi.org/10.1186/s12883-020-01928-5>.

Abstract

Background Cognitive impairment may be correlated with cardiovascular dysautonomia, including blood pressure (BP) dysregulation, in Parkinson's disease (PD), but the association between these factors in dementia with Lewy bodies (DLB) is uncertain. This study aimed to clarify whether cardiovascular dysautonomia had an influence on cognitive function in Lewy body disease or not. **Methods** 99 patients with de novo PD (n=75) and DLB (n=24) were evaluated using the Mini-Mental State Examination (MMSE) and Frontal Assessment Battery (FAB). Cardiac 123I-metaiodobenzylguanidine (MIBG) scintigraphy, orthostatic hypotension (OH), supine hypertension (SH), postprandial hypotension (PPH), nocturnal BP fall in 24-hour ambulatory blood pressure monitoring (ABPM) and constipation were estimated. Associations of these factors with cognitive and executive dysfunction were examined. **Results** In DLB, MIBG uptake was reduced and OH, PPH and SH were severely disturbed, compared to PD. The nocturnal BP fall in ABPM was lower in DLB, and the failure of nocturnal BP fall in PD was associated with MMSE, after adjustment for other clinical features. FAB was significantly associated nocturnal BP fall, age and SH in PD, but no significant correlations among factors were found for DLB. **Conclusion** The significant association between nocturnal BP dysregulation and cognitive or executive decline in PD might be due to impaired microvascular circulation or invasion of α -synuclein in the CNS. The lack of a correlation of BP insufficiency with cognitive impairment in DLB suggests initial involvement of Lewy body pathology in the neocortex, regardless of Lewy body invasion of the autonomic nervous system.

Background

Parkinson's disease (PD) is commonly accepted to be associated with motor symptoms and various non-motor symptoms, including behavioral changes such as depression, sleep disturbance, fatigue and autonomic dysfunction. Autonomic impairment associated with PD is characterized by clinical features of constipation, sweating, orthostatic hypotension (OH) and postprandial hypotension (PPH), even in the early phase [1]. OH occurs through sympathetic noradrenergic dysfunction and is clinically important in 20-50% of patients with PD [1]. Survival depends on the OH status, with a greater risk of death in PD with OH than in PD without OH [2]. OH may also affect cognition [3], daily activities, and quality of life [4]. Patients with PD with OH have significantly worse sustained attention and visual episodic memory [5] and significantly lower scores on the Mini-Mental State Examination (MMSE) [6] [7].

Abnormal blood pressure (BP) fluctuations are also common in PD [8, 9]. The normal nocturnal BP fall in a healthy person disappears in PD, especially in a case with autonomic dysfunction [10, 11]. Cognitive decline in older people is associated with abnormal BP fluctuations, such as the absence of a normal nocturnal BP fall [11, 12], and cognitive impairment in PD has also been associated with abnormal BP fluctuation [13] (Tanaka *et al.*, 2018). Autonomic dysfunction in dementia with Lewy bodies (DLB) is generally more severe than that in PD [14]. The severity of autonomic failure in DLB is intermediate between those in PD and multiple system atrophy [15]. OH is likely to be severer in DLB than in PD because it is thought to be due to Lewy body involvement in the rostral ventrolateral medulla and

medullary raphe, which may control sympathetic outflow [15]. However, it is not clear whether cognitive dysfunction in DLB is correlated with blood circulatory insufficiency such as OH and abnormal BP fluctuation, as occurs in PD.

The aims of this study were to examine the association of cardiovascular dysautonomia and cognitive impairments in de novo PD and DLB using ^{123}I -metaiodobenzylguanidine (MIBG) uptake by the heart, responses of BP and plasma norepinephrine in a head-up tilt-table test (HUT), a 75-g oral glucose tolerance test (75-g OGTT) for PPH, 24-hour ambulatory blood pressure monitoring (ABPM) and constipation.

Methods

Study design and participants

The subjects were 75 de novo PD patients who were diagnosed using the criteria for PD proposed by the UK Parkinson's Disease Society Brain Bank [16], and 24 de novo DLB patients with a central feature and two or more core features in diagnostic criteria, which is sufficient for diagnosis of probable DLB [17]. All patients were examined at Disan Hospital, The Jikei University School of Medicine, between January 2012 and March 2018, and were diagnosed by at least two neurologists. We also used a 1-year rule to distinguish DLB from PD with dementia [17].

Patients with overt diabetes or clinically relevant cardiac disease, and those who had been treated with tricyclic antidepressants, tetracyclic antidepressants, serotonin reuptake inhibitors, and serotonin and norepinephrine reuptake inhibitors were excluded from the study. None of the patients had received levodopa, other anti-Parkinson drugs, or treatment for OH. Global cognition and executive function were evaluated using the MMSE and the Frontal Assessment Battery (FAB). No patient had atrophy on brain MRI of the putamen, brainstem or cerebellum. If patients were already receiving antihypertensive drugs, such drugs were withdrawn at least 48 hours before evaluation of OH. All patients received levodopa or a dopamine agonist for their parkinsonism after this study, and all had a good response.

The motor severity of PD was assessed using the Unified Parkinson's Disease Rating Scale (UPDRS) motor score. The patients were divided into tremor-dominant, akinetic-rigid, and mixed type subgroups based on the tremor and non-tremor scores, which were obtained using part III of the UPDRS [18]. This study was approved by the Ethics Committee of The Jikei University School of Medicine, and all subjects gave written informed consent before enrollment.

Cardiac ^{123}I -metaiodobenzylguanidine (MIBG) scintigraphy

Cardiac sympathetic denervation was evaluated using MIBG scintigraphy. The ratio of the average pixel count in the heart (H) to that in the mediastinum (M) (H/M ratio) at 15 min (early) and 3 hours (delayed)

after injection of 111 MBq 123I-MIBG (Fujifilm RI Pharma Co., Ltd. Tokyo, Japan) was calculated [19].

Olfactory assessment

Olfactory function was assessed by the odor stick identification test Japan (OSIT-J) (Daiichi Yakuhin Sangyo Co. Ltd., Tokyo, Japan), as described in our previous study [20]. The number of correct responses for the 12 odorants was defined as the OSIT-J score, which has been shown to be significantly correlated with those on the University of Pennsylvania Smell Identification Test (UPSIT) and the cross-cultural, smell identification test (CC-SIT) [21, 22].

Head-up tilt-table test (HUT)

All subjects underwent HUT in a silent room maintained at an ambient temperature of 23 to 26°C. After an overnight fast, the test was started at 9:00 am. After resting for 20 min in the supine position, the subject was tilted to a 60° upright position for 3 min. Brachial systolic blood pressure (SBP) and diastolic blood pressure (DBP) were measured by an automated sphygmomanometer after 20 min of rest in the supine position and every 1 min after the subject was tilted for up to 3 min. The maximum decreases in SBP and DBP during tilt were evaluated. Plasma norepinephrine concentrations in serum (NE, µg/ml) were measured with the subjects in a supine position after 20 minutes of rest and after 10 min in a tilted position. OH was defined as a fall in SBP by ≥ 20 mmHg [23]. Supine hypertension (SH) was defined as SBP >140 mmHg or DBP >90 mmHg, as measured after 20 min of rest in the supine position [24]. Neurogenic SH was defined as a case of OH with SBP >140 mmHg or DBP >90 mmHg in the supine position [25].

24-hour ambulatory blood pressure monitoring (ABPM)

24-hour ABPM was performed using a noninvasive automated portable recorder in hospitalized patients and outpatients. BP was measured every 30 min during the day (7:00-21:00) and every hour at night (22:00-6:00). SBP was used as an indicator of BP. The nocturnal fall in BP was calculated as: $\frac{SBP_{\text{day}} - SBP_{\text{night}}}{SBP_{\text{day}}} \times 100$ (%), where SBP_{day} is the mean SBP during the day, and SBP_{night} is the mean SBP at night. Cases with nocturnal falls in BP of $\geq 10\%$, $<10\%$ and no fall were defined as dipper, non-dipper, and riser types [26, 27].

75-g oral glucose tolerance test for evaluation of postprandial hypotension

After overnight fasting (except for non-caloric liquids), the 75-g OGTT was started between 9:00 and 10:00 am in a quiet room at an ambient temperature of 23 to 26°. Whenever possible, we performed HUT followed by the 75-g OGTT on the same day. If this was not possible, the 75-g OGTT was performed on the day after HUT. After 20 min resting in the supine position, the subjects drank 75 g of glucose water (calorie content, 300 kcal) and remained resting and awake in the supine position for 120 min. After 20 min and then every 10 min for the next 120 min, brachial SBP and DBP were measured by an automated sphygmomanometer in the supine position. The time to the maximum drop in SBP on the 75-g OGTT was measured. Postprandial hypotension (PPH) was defined as a maximum decrease in SBP of 20 mmHg within 2 hours after glucose intake [28, 29].

Constipation

Constipation was defined as the presence of at least one of two criteria: absence of daily defecation, and use of drugs to treat constipation [30].

Statistical analysis

Statistical analyses were performed with statistical software (Esumi Co., Ltd., Tokyo, Japan). Differences between groups were compared by Wilcoxon rank sum test for continuous variables. Pairwise comparisons were made using χ^2 tests for binary variables. Lepage analysis was used to evaluate differences in nocturnal fall in BP. Associations of MMSE and FAB scores with clinical factors such as age, gender, symptom duration, UPDRS motor score, motor subtype, olfaction, cardiac MIBG uptake, BP fall in HUT, NE at rest in the supine position in HUT, nocturnal fall in BP in ABPM, PPH, SH and constipation were evaluated by multiple regression analysis. $P < 0.05$ was considered to indicate significance.

Results

Cognitive impairment and cardiovascular dysautonomia in PD and DLB

A comparison of the characteristics of the PD and DLB patients is given in Table 1. DLB patients were older than PD patients, but there was no significant difference in the duration of PD and DLB. PD patients were significantly more frequently female, while DLB patients were more commonly male. There were no significant differences in UPDRS motor scores and motor subtype between PD and DLB cases. DLB patients had more severely impaired olfaction, lower MMSE and FAB scores, a lower H/M ratio in cardiac MIBG uptake, a lower BP fall in HUT, lower NE in a resting supine position in HUT, and a higher prevalence of OH, compared to PD cases. There was a reduced nocturnal fall in BP in ABPM and a higher percentage of non-dipper/riser types among DLB cases. BP fall in PPH and the prevalence of PPH were larger in DLB

cases. The prevalence of SH did not differ significantly, but that of neurogenic SH was higher in DLB patients. The prevalence of constipation in DLB cases was also higher than that in PD cases.

Correlations between cognitive impairment and cardiovascular dysautonomia

In multiple regression analyses, MMSE in PD was significantly associated with nocturnal fall in BP in ABPM ($p=0.0275$), after adjustment for age, disease duration, UPDRS motor score, motor subtype, olfaction, cardiac MIBG scintigraphy, BP fall in HUT, NE at rest in a supine position in HUT, PPH, SH and constipation (Table 2); and FAB in PD was significantly related with nocturnal fall in BP ($p=0.0395$), aging ($p=0.0076$) and SH ($p=0.0037$) (Table 3). In contrast, neither MMSE nor FAB in DLB was associated with nocturnal fall in BP or any other clinical variable (Tables 4, 5).

Discussion

In this study, DLB patients clearly had more severe cognitive decline compared with PD patients. Olfaction was more impaired in DLB, and olfactory dysfunction has been associated with cognitive dysfunction [20, 31]. De novo PD with mild cognitive decline shows olfactory impairment compared with patients without cognitive dysfunction [32]. This is consistent with DLB patients with impaired cognition having more severe olfactory dysfunction than that in PD patients.

MIBG uptake in scintigraphy indicating cardiac sympathetic denervation was lower in patients with DLB than in those with PD. BP fall on standing was significantly larger, NE at rest was lower, and the BP fall in PPH and SH were larger in DLB. Neurogenic SH differed significantly between PD and DLB cases, but not the prevalence of SH. The prevalence of constipation in DLB was higher than in PD, which suggests that intestinal autonomic dysfunction might be severer in DLB. Overall, our findings suggest that DLB involves wider spread severer sympathetic and parasympathetic autonomic dysfunction compared with PD, which might be due to peripheral and CNS impairments [14].

The nocturnal fall in BP in ABPM found in healthy persons was reduced more in DLB than PD. Abnormal daily BP fluctuation in PD [33, 34] has been associated with cardiovascular dysautonomia [34], but has rarely been reported in DLB. PD patients with this condition, including reduced or reverse nocturnal BP fall on ABPM, have also been found to have a higher prevalence of OH [33].

Thus, severe cardiovascular autonomic dysfunction in DLB might be linked to more profound impairment of the nocturnal fall in BP. Cognitive function has previously been linked to abnormal BP fluctuation in PD [11, 13], including an abnormality in nocturnal BP fall in ABPM. The novel finding in our study is that nocturnal BP abnormality was associated with cognitive and executive dysfunction in early stage and de novo PD after adjustment for other cardiovascular dysautonomic factors, including OH, PPH, SH and cardiac sympathetic impairment indicated by MIBG uptake insufficiency.

Several pathogeneses have been suggested for the association of cognitive dysfunction and BP abnormality, but the underlying mechanism is still unclear. The Braak hypothesis [35] suggests that Lewy body (LB) pathology initially occurs in the olfactory nucleus and dorsal motor nucleus and progressively ascends through the brain stem to the cortex, causing noradrenergic and dopaminergic neuronal degeneration that results in progression of motor, cognitive, and autonomic impairment. Cognitive decline in PD has been associated with specific patterns of LB density in the entorhinal cortex and anterior cingulate cortex [36], which plays a role in autonomic nervous system (ANS) control, including the higher centers of autonomic regulations [37]. Involvement of the anterior cingulate cortex might simultaneously cause cognitive impairment and cardiovascular sympathetic failure.

Noradrenergic projection from the locus coeruleus (LC) spreads extensively in the whole brain cortex, including the hippocampus, entorhinal and mediotemporal cortex, cingulate gyrus and neocortex. Tyrosine hydroxylase immunoreactivity is lost in neurons projecting from the LC due to the LB pathology in PD [38]. Involvement of the noradrenergic neurons of the LC is increasingly recognized as a potential major contributor to cognitive manifestations in early PD, particularly for impaired attention [39]. The LC projects to the parasympathetic neurons of the dorsal motor nucleus of the vagus (the nucleus ambiguus), while the descending pathway projects to sympathetic preganglionic neurons in the spinal cord [39]. Therefore, the LC should influence cardiovascular modulation via insufficiency of cardiac parasympathetic and cardiovascular sympathetic function. The LC also regulates part of the wake-promoting circuit with the suprachiasmatic nucleus and dorsomedial hypothalamus [40]. Therefore, spoiling of the LC may cause abnormal daily BP fluctuation, in addition to cardiovascular sympathetic failure.

BP insufficiency such as OH, including circadian rhythm failure, is associated with increased white matter hyperintensities (WMHs) on MRI, even in older people [41, 42], and cognitive impairment and WMHs are associated with OH and SH in PD. Cognitive scores and WMH volumes on MRI are severely reduced in PD patients with both OH and SH, and lability of BP circulation may be related to cognitive function and WMH volume on MRI [11]. Our study revealed that abnormal BP fluctuation, and especially a reduced nocturnal fall in BP, was associated with cognitive and executive function in PD, after adjustment for other autonomic characteristics, including cardiovascular sympathetic function reflected by cardiac MIBG uptake, OH, PPH, circulatory NE concentration, and constipation. This suggests that increased lability of daily BP and nocturnal BP is a risk factor for cognitive impairment, even in early de novo PD. Furthermore, FAB scores, but not MMSE scores, were correlated with SH and aging in our PD patients. This may indicate executive dysfunction occurred due to damage of prefrontal areas, which is readily attributable to cerebrovascular circulatory insufficiency of the cortex, white matter or age-related changes in the brain in PD [23].

In contrast to PD, we found that cognitive and executive impairments in DLB patients were not correlated with lability of BP. Dysautonomia in DLB seems to be severer than that in PD from our results and previous reports [14, 15]. It is uncertain whether PD and DLB including PDD are separate disease entities or parts of the same disease spectrum. LB pathology in PD is restricted to the brainstem and limbic

regions, while the pathology more quickly extends to the neocortex in DLB. LB pathology in PD is also not so widely distributed in autonomic nervous organs, compared with that in DLB. The discrepancy between cardiovascular and cognitive dysfunction in DLB might suggest that regional invasion of LB pathology differs between the neocortex and sympathetic autonomic center. Braak's hypothesis [35] suggests that α synucleinopathy initially involves the intestinal organ and ascends to the brainstem, including the dorsal motor nucleus of the vagus, LC, medullary reticular formation, raphe nuclei, and peripheral sympathetic nervous system. These organs are associated with modulation of cardiovascular autonomic regulation in early PD.

Cognitive dysfunction in PD might occur due to white matter damage from BP dysregulation and noradrenergic decline of the LC [38]. Cognition dysfunction should be associated with cardiovascular autonomic failure if LB pathology involves the ACC or insular cortex. Cognitive decline has already progressed due to involvement of LB pathology in brain cortex, so that cognitive impairment in DLB should not be strongly influenced by BP dysregulation. Alzheimer disease (AD) pathology with hyperphosphorylated tau and amyloid- β ($A\beta$) may also contribute to cognitive decline in DLB and PD. $A\beta$ plaques are significantly more common in cortical and subcortical regions in DLB compared to Parkinson's disease with dementia (PDD) [43, 44], and DLB displays concurrent AD-related pathology compared to PDD [45]. Cardiovascular dysautonomia including reduced cardiac MIBG uptake and OH is not as impaired in AD compared to DLB [46]. Thus, AD pathology may be not correlated with ANS effects, and the increased AD pathology may induce dissociation between cognitive and BP dysregulation in DLB compared to PD.

Conclusions

BP dysregulation, especially reduced nocturnal BP fall, was associated with cognitive and executive decline in PD, and this may be driven by impaired microvascular circulation or infiltration of α synuclein from the peripheral ANS to the CNS, such as the LC or ACC. The absence of a correlation between cognitive and BP dysregulation in DLB is due to earlier spread of LB pathology to the neocortex, while ascending LB invasion of the LC or ANS occurs in PD. The more severe AD pathology in the cortex in DLB compared to PD might also contribute to dissociation of cognitive dysfunction and BP abnormality. Therefore, treatment for BP dysregulation may prevent progression of cognitive decline in PD, but not in DLB.

Abbreviations

Alzheimer disease (AD), ambulatory blood pressure monitoring (ABPM), amyloid- β ($A\beta$), autonomic nervous system (ANS), blood pressure (BP), cross-cultural smell identification test (CC-SIT), dementia with Lewy bodies (DLB), diastolic blood pressure (DBP), Frontal Assessment Battery (FAB), heart (H), head-up tilt-table test (HUT), 123I-metaiodobenzylguanidine (MIBG), Lewy body (LB), locus coeruleus (LC), mediastinum (M), Mini-Mental State Examination (MMSE), 75-g oral glucose tolerance test (75-g OGTT), orthostatic hypotension (OH), Parkinson's disease (PD), Parkinson's disease with dementia (PDD),

postprandial hypotension (PPH), supine hypertension (SH), systolic blood pressure (SBP), Unified Parkinson's Disease Rating Scale (UPDRS), white matter hyperintensities (WMHs),

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of The Jikei University School of Medicine, and all subjects gave written informed consent before enrollment for participation and to publication.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors report no competing interests.

Funding

Funding was from institutional sources only.

Author contributions

Hisayoshi Oka: drafting/revising the manuscript for content, including medical writing; study concept and design; acquisition of data; analysis and interpretation of data. Tadashi Umehara: study concept and design; acquisition of data; review and critique. Atsuo Nakahara: study concept and design; acquisition of data; review. Hiromasa Matsuno: study concept and design; acquisition of data; review. All authors have seen and approved the manuscript.

References

1. Goldstein DS. Orthostatic hypotension as an early finding in Parkinson disease. *Clin Auton Res*. 2006; 6: 46-54.
2. Goldstein DS, Holmes C, Sharabi Y, Wu T. Survival in synucleinopathies: A prospective cohort study. *Neurology*. 2015; 85: 1554-1561.
3. Perlmutter LC, Sarda G, Casavant V, Mosnaim AD. A review of orthostatic blood pressure regulation and its association with mood and cognition. *Clin Auton Res*. 2012; 22: 99-107.
4. De Pablo-Fernandez E, Tur C, Revesz T, Lees AJ, Holton JJ, Warner TT. Association of autonomic dysfunction with disease progression and survival in Parkinson disease. *JAMA Neurol*. 2017;74: 970-976.
5. Allcock LM, Kenny RA, Mosimann UP, Tordoff S, Wesnes KA, Hildreth AJ, Burn DJ. Orthostatic hypotension in Parkinson's disease: association with cognitive decline? *Int J Geriatr Psychiatry*. 2006;21: 778-783.
6. Hohler AD, Zuzuárregui JR, Katz DI, Depiero TJ, Hehl CL, Leonard A, Allen V, Dentino J, Gardner M, Phenix H, Saint-Hilaire M, Ellis T. Differences in motor and cognitive function in patients with Parkinson's disease with and without orthostatic hypotension. *Int J Neurosci*. 2012; 122: 233-236.
7. McDonald C, Newton JL, Burn DJ. Orthostatic hypotension and cognitive impairment in Parkinson's disease: Causation or association? *Mov Disord*. 2016; 31: 937-946.
8. Guo H, Tabara Y, Igase M, Yamamoto M, Ochi N, Kido T, Uetani E, Taguchi K, Miki T, Kohara K. Abnormal nocturnal blood pressure profile is associated with mild cognitive impairment in the elderly: the J-SHIPP study. *Hypertens Res*. 2010; 33: 32-36.
9. Komori T, Eguchi K, Saito T, Nishimura Y, Hoshide S, Kario K. Riser blood pressure pattern is associated with mild cognitive impairment in heart failure patients, *Am J Hypertens*. 2016;29: 194-201.
10. Alperovitch A, Blachier M, Soumaré A, Ritchie K, Dartigues JF, Richard-Harston S, Tzourio C. Blood pressure variability and risk of dementia in an elderly cohort, the Three-City Study. *Alzheimers Dement*. 2014; 10: S330-S337.
11. Kim JS, Oh YS, Lee KS, Kim YI, Yang DW, Goldstein DS. Association of cognitive dysfunction with neurocirculatory abnormalities in early Parkinson disease. *Neurology*. 2012; 79: 1323-1331.
12. Svenningsson P, Westman E, Ballard C, Aarsland D. Cognitive impairment in patients with Parkinson's disease: diagnosis, biomarkers, and treatment. *Lancet Neurol*. 2012; 11: 697-707.
13. Tanaka R, Shimo Y, Yamashiro K, Ogawa T, Nishioka K, Oyama G, Umemura A, Hattori N. Association between abnormal nocturnal blood pressure profile and dementia in Parkinson's disease. *Parkinsonism Relat Disord*. 2018;46: 24-29.
14. Akaogi Y, Asahina M, Yamanaka Y, Koyama Y, Hattori T. Sudomotor, skin vasomotor, and cardiovascular reflexes in 3 clinical forms of Lewy body disease. *Neurology*. 2009;73: 59-65.

15. Benarroch EE, Schmeichel AM, Low PA, Boeve BF, Sandroni P, Parisi JE. Involvement of medullary regions controlling sympathetic output in Lewy body disease. *Brain*. 2005;128: 338-344.
16. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease. A clinicopathological study of 100 cases. *J Neurol Neurosurgery Psychiatry*. 1992; 55: 181-184.
17. McKeith IG, Boeve BF, Dickson DW, Halliday G, Taylor JP, Weintraub D, Aarsland D, et al. Diagnosis and management of dementia with Lewy bodies: Fourth consensus report of the DLB Consortium. *Neurology*. 2017;89: 88-100.
18. Spiegel J, Hellwig D, Samnick S, Jost W, Möllers MO, Fassbender K, Kirsch CM, Dillmann U. Striatal FP-CIT uptake differs in the subtypes of early Parkinson's disease. *J Neural Transm*. 2007; 114: 331-335.
19. Oka H, Toyoda C, Yogo M, Mochio S. Cardiovascular dysautonomia in de novo Parkinson's disease without orthostatic hypotension. *Eur J Neurol*. 2011;18: 286-292.
20. Oka H, Toyoda C, Yogo M, Mochio S. Olfactory dysfunction and cardiovascular dysautonomia in Parkinson's disease. *J Neurol*. 2010; 257: 969-976.
21. Ogihara H, Kobayashi M, Nishida K, Kitano M, Takeuchi K. Applicability of the cross-culturally modified University of Pennsylvania Smell Identification Test in a Japanese population. *Am J Rhinol Allergy*. 2011; 25: 404-410.
22. Hashimoto Y, Fukazawa K, Fujii M, Takayasu S, Muto T, Saito S, Takashima Y, Sakagami M. Usefulness of the odor stick identification test for Japanese patients with olfactory dysfunction. *Chem Senses*. 2004;29: 565-571.
23. Kaufmann H. Consensus statement on the definition of orthostatic hypotension, pure autonomic failure and multiple system atrophy. *Clin Auton Res*. 1996; 6:125-126.
24. Umehara T, Matsuno H, Toyoda C, Oka H. Clinical characteristics of supine hypertension in de novo Parkinson disease. *Clin Auton Res*. 2016; 26: 15-21.
25. Fanciulli A, Jordan J, Biaggioni I, Calandra-Buonaura G, Cheshire WP, Cortelli P, Eschlboeck S, et al. Consensus statement on the definition of neurogenic supine hypertension in cardiovascular autonomic failure by the American Autonomic Society (AAS) and the European Federation of Autonomic Societies (EFAS): endorsed by the European Academy of Neurology (EAN) and the European Society of Hypertension (ESH). *Clin Auton Res*. 2018;28:355-362.
26. Manabe Y, Fujii D, Kono S, Sakai Y, Tanaka T, Narai H, Omori N, Imai Y, Abe K. Systemic blood pressure profile correlates with cardiac ¹²³I-MIBG uptake in patients with Parkinson's disease. *J Neurol Sci*. 2011;307: 153-156.
27. Oka H, Nakahara A, Umehara T. Rotigotine improves abnormal circadian rhythm of blood pressure in Parkinson's Disease. *Eur Neurol*. 2018;79:281-286.
28. Umehara T, Toyoda C, Oka H. Postprandial hypotension in de novo Parkinson's disease: A comparison with orthostatic hypotension. *Parkinsonism Relat Disord*. 2014;20: 573-577.

29. Umehara T, Nakahara A, Matsuno H, Toyoda C, Oka H. Predictors of postprandial hypotension in elderly patients with de novo Parkinson's disease. *J Neural Transm (Vienna)*. 2016;123: 1331-1339.
30. Umehara T, Oka H, Nakahara A, Matsuno H, Toyoda C. High norepinephrergic orthostatic hypotension in early Parkinson's disease. *Parkinsonism Relat Disord*. 2018;55: 97-102.
31. Bohnen NI, Müller ML, Kotagal V, Koeppe RA, Kilbourn MA, Albin RL, Frey KA. Olfactory dysfunction, central cholinergic integrity and cognitive impairment in Parkinson's disease. *Brain*. 2010;133: 1747-1754.
32. Baba T, Kikuchi A, Hirayama K, Nishio Y, Hosokai Y, Kanno S, Hasegawa T, Sugeno N, Konno M, Suzuki K, Takahashi S, Fukuda H, Aoki M, Itoyama Y, Mori E, Takeda A. Severe olfactory dysfunction is a prodromal symptom of dementia associated with Parkinson's disease: a 3 year longitudinal study. *Brain*. 2012;135: 161-169.
33. Berganzo K, Diez-Arrola B, Tijero B, Somme J, Lezcano E, Llorens V, Ugarriza I, Ciordia R, Gómez-Esteban JC, Zarranz JJ. Nocturnal hypertension and dysautonomia in patients with Parkinson's disease: are they related? *J Neurol*. 2013; 260: 1752-1756.
34. Milazzo V, Di Stefano C, Vallelonga F, Sobrero G, Zibetti M, Romagnolo A, Merola A, Milan A, Espay AJ, Lopiano L, Veglio F, Maule S. Reverse blood pressure dipping as marker of dysautonomia in Parkinson disease. *Parkinsonism Relat Disord*. 2018; 56: 82-87.
35. Braak H, Tredici KD, Rüb U, de Vos RAI, Jansen Steur ENH, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging*. 2003; 24:197-211.
36. Kövari E, Gold G, Herrmann F, Canuto A, Hof PR, Bouras C, Giannakopoulos P. Lewy body densities in the entorhinal and anterior cingulate cortex predict cognitive deficits in Parkinson's disease. *Acta Neuropathol*. 2003;106: 83-88.
37. Matsui H, Udaka F, Miyoshi T, Hara N, Tamura A, Oda M, Kubori T, Nishinaka K, Kameyama M. Three-dimensional stereotactic surface projection study of orthostatic hypotension and brain perfusion image in Parkinson's disease. *Acta Neurol Scand*. 2005;112: 36-41.
38. Del Tredici K, Braak H. Dysfunction of the locus coeruleus norepinephrine system and related circuitry in Parkinson's disease related dementia. *J Neurol Neurosurg Psychiatry*. 2013; 84: 774-783.
39. Benarroch EE. Locus coeruleus. *Cell Tissue Res*. 2018; 373:221-232.
40. Gall AJ, Todd WD, Blumberg MS. Development of SCN connectivity and the circadian control of arousal: a diminishing role for humoral factors? *PLoS One*. 2012; 7: e45338.
41. Schmidt R, Ropele S, Enzinger C, Petrovic K, Smith S, Schmidt H, Matthews PM, Fazekas F. White matter lesion progression, brain atrophy, and cognitive decline: the Austrian stroke prevention study. *Ann Neurol*. 2005;58: 610-616.
42. Van Den Heuvel DMJ, Ten Dam VH, De Craen AJM, Admiraal-Behloul F, Olofsen H, Bollen EL, Jolles J, Murray HM, Blauw GJ, Westendorp RG, van Buchem MA. Increase in periventricular white matter hyperintensities parallels decline in mental processing speed in a non-demented elderly population. *J Neurol Neurosurg Psychiatry*. 2006;77: 149-153.

43. Walker L, McAleese KE, Thomas AJ, Johnson M, Martin-Ruiz C, Parker C, et al. Neuropathologically mixed Alzheimer's and Lewy body disease: burden of pathological protein aggregates differs between clinical phenotypes. *Acta Neuropathol.* 2015; 129: 729-748.
44. Hepp DH, Vergoossen DL, Huisman E, Lemstra AW; Netherlands Brain Bank, Berendse HW, et al. Distribution and load of amyloid- β pathology in Parkinson Disease and Dementia with Lewy Bodies. *J Neuropathol Exp Neurol.* 2016; 75: 936-945.
45. Walker L, Stefanis L, Attems J. Clinical and neuropathological differences between Parkinson's disease, Parkinson's disease dementia, and dementia with Lewy bodies - current issues and future directions. *J Neurochem.* 2019; Mar 20. doi: 10.1111/jnc. 14698. [Epub ahead of print]
46. Kim JS, Park HE, Oh YS, Song IU, Yang DW, Park JW, et al. ^{123}I -MIBG myocardial scintigraphy and neurocirculatory abnormalities in patients with dementia with Lewy bodies and Alzheimer's disease. *J Neurol Sci.* 2015; 357: 173-177.

Tables

Table 1. Comparison of clinical characteristics in patients with de novo DLB and PD

Item	PD (N=75)	DLB (N=24)	p value ^a
Age (years)	72.2±9.4	80.6±5.0	0.001
Gender (male/female)	27/48	18/6	0.001
Symptom duration (years)	1.6±1.6	1.9±1.7	0.296
UPDRS motor score	20.3±10.9	21.3±10.8	0.236
Subtype (akinetic-rigid/tremor-dominant/mixed)	45/30	19/5	0.087
Olfaction	4.1±2.8	2.4±2.2	0.010
Mini-mental state examination	27.7±2.2	20.0±4.7	0.001
Frontal assessment battery	15.1±2.8	10.2±3.4	0.001
MIBG H/M ratio delay	1.44±0.33	1.22±0.39	0.001
BP fall in head-up tilt-table test (HUT) (mmHg)	23.6±20.0	48.5±23.6	0.001
Prevalence of OH (-/+)	34/31	1/23	0.001
Norepinephrine at rest in supine position in HUT (µg/ml)	217.9±88.2	172.8±82.5	0.009
Nocturnal fall in BP in ABPM (%)	8.5±9.4	-4.4±12.6	0.001
ABPM type (dipper/non dipper or riser type)	37/38	4/20	0.005
BP fall in postprandial hypotension (mmHg)	18.4±13.1	32.8±17.2	0.001
Postprandial hypotension (-/+)	42/33	6/18	0.008
BP in supine hypertension (mmHg)	134.8±21.5	147.5±19.2	0.011
Supine hypertension (-/+)	45/30	10/14	0.116
Neurogenic supine hypertension (-/+)	56/19	12/13	0.013
Constipation (-/+)	25/50	1/23	0.008

^a Wilcoxon rank sum test and Lepage analysis for continuous variables and χ^2 test for binary variables.

UPDRS: Unified Parkinson's Disease Rating Scale, MIBG: ¹²³I-metaiodobenzylguanidine, H/M: ratio of the average pixel count in the heart (H) to that in the mediastinum (M), ABPM: 24-hour ambulatory blood pressure monitoring

Table 2. Relationships of MMSE with clinical parameters in PD

Clinical parameter	Parameter Estimate	Standardp-value ^a	
		Error	
Age (years)	-0.053	0.031	0.099
Gender (male/female)	0.260	0.594	0.664
Symptom duration (years)	-0.208	0.162	0.204
UPDRS motor score	-0.017	0.026	0.515
Subtype (akinetic-rigid/tremor-dominant/mixed)	-0.906	0.528	0.091
Olfaction	-0.159	0.101	0.121
MIBG H/M ratio delay	-0.0310	0.837	0.971
BP fall in head-up tilt-table test (HUT) (mmHg)	-0.013	0.013	0.330
Norepinephrine at rest in supine position in HUT (µg/ml)	0.005	0.003	0.095
Nocturnal fall in BP in ABPM (%)	0.066	0.029	0.028
BP fall in postprandial hypotension (mmHg)	0.028	0.020	0.169
BP in supine hypertension (mmHg)	-0.180	0.013	0.165
Constipation	-0.018	0.622	0.519

^a Analysis were performed by multiple regression analysis.

Abbreviations are as shown in Table 1.

Table 3. Relationships of FAB with clinical parameter in PD

Clinical parameter	Parameter Estimate	Standardp-value ^a	
		Error	
Age (years)	-0.098	0.036	0.008
Gender (male/female)	-0.867	0.667	0.200
Symptom duration (years)	-0.028	0.183	0.880
UPDRS motor score	-0.012	0.02904	0.686
Subtype (akinetic-rigid/tremor-dominant/mixed)	-0.551	0.595	0.358
Olfaction	-0.002	0.114	0.990
MIBG H/M ratio delay	-0.118	0.944	0.901
BP fall in head-up tilt-table test (HUT) (mmHg)	0.018	0.015	0.225
Norepinephrine at rest in supine position in HUT (µg/ml)	0.002	0.003	0.650
Nocturnal fall in BP in ABPM (%)	0.070	0.033	0.040
BP fall in postprandial hypotension (mmHg)	0.002	0.022	0.920
BP in supine hypertension (mmHg)	-0.043	0.014	0.004
Constipation	-0.199	0.701	0.778

^a Analysis were performed by multiple regression analysis.

Abbreviations are as shown in Table 1.

Table 4. Relationships of MMSE with clinical parameters in DLB

Clinical parameter	Parameter Estimate	Standardp-value ^a	
		Error	
Age (years)	-0.435	0.255	0.118
Gender (male/female)	-3.063	3.454	0.396
Symptom duration (years)	0.959	0.807	0.262
UPDRS motor score	-0.176	0.128	0.198
Subtype (akinetic-rigid/tremor-dominant/mixed)	1.492	3.664	0.693
Olfaction	0.566	0.651	0.405
MIBG H/M ratio delay	1.136	3.243	0.733
BP fall in head-up tilt-table test (HUT) (mmHg)	0.039	0.049	0.448
Norepinephrine at rest in supine position in HUT (µg/ml)	-0.024	0.019	0.245
Nocturnal fall in BP in ABPM (%)	0.095	0.143	0.524
BP fall in postprandial hypotension (mmHg)	0.110	0.100	0.298
BP in supine hypertension (mmHg)	-0.010	0.072	0.895
Constipation	-6.900	5.725	0.256

^a Analysis were performed by multiple regression analysis.

Abbreviations are as shown in Table 1.

Table 5. Relationships of FAB with clinical parameters in DLB

Clinical parameter	Parameter Estimate	Standardp-value ^a	
		Error	
Age (years)	-0.326	0.188	0.114
Gender (male/female)	-1.791	2.551	0.499
Symptom duration (years)	0.221	0.596	0.719
UPDRS motor score	-0.169	0.094	0.104
Subtype (akinetic-rigid/tremor-dominant/mixed)	0.081	2.706	0.976
Olfaction	-0.177	0.481	0.721
MIBG H/M ratio delay	1.731	2.395	0.486
BP fall in head-up tilt-table test (HUT) (mmHg)	0.006	0.036	0.868
Norepinephrine at rest in supine position in HUT (µg/ml)	0.003	0.014	0.829
Nocturnal fall in BP in ABPM (%)	0.016	0.106	0.885
BP fall in postprandial hypotension (mmHg)	-0.016	0.074	0.830
BP in supine hypertension (mmHg)	-0.009	0.053	0.875
Constipation	1.464	4.227	0.736

^a Analysis were performed by multiple regression analysis.

Abbreviations are as shown in Table 1.