

Compliance with spectacle use in children with refractive errors- a systematic review and meta-analysis

Nonita Dhirar

Post Graduate Institute of Medical Education and Research

Sankalp Dudeja

Post Graduate Institute of Medical Education and Research

Mona Duggal (✉ monaduggal2@gmail.com)

Post Graduate Institute of Medical Education and Research <https://orcid.org/0000-0002-9404-2871>

Parul Chawla Gupta

Post Graduate Institute of Medical Education and Research, Chandigarh

Nishant Jaiswal

Post Graduate Institute of Medical Education and Research, Chandigarh

Meenu Singh

Post Graduate Institute of Medical Education and Research, Chandigarh

Jagat Ram

Post Graduate Institute of Medical Education and Research, Chandigarh

Research article

Keywords: Compliance, spectacles, refractive error, children

Posted Date: October 24th, 2019

DOI: <https://doi.org/10.21203/rs.2.16388/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background Primary objective of this review was to measure compliance with spectacle use in children with refractive errors. Secondary objective was to understand the reasons for non-compliance.

Methods The databases searched were Ovid, EMBASE, CINAHL and Pubmed. All studies up to March, 2018 were included. The search terms were- ((((((Compliance [Title/Abstract]) OR Adherence[Title/Abstract]) OR Compliant[Title/Abstract]) OR Adherent[Title/Abstract])) AND (((Spectacle[Title/Abstract]) OR Spectacles[Title/Abstract]) OR Eye Glasses[Title/Abstract])) AND (((Child[Title/Abstract]) OR Children[Title/Abstract]) OR Adolescent[Title/Abstract]) OR Adolescents[Title/Abstract])). Two researchers independently searched the databases and initial screening obtained 33 articles. The PRISMA guidelines were followed for conducting and writing the systematic review. Two reviewers assessed data quality independently using the Quality Assessment tool for systematic reviews of observational studies (QATSO). Poor quality studies were those, which had a score of less than 33% on the QATSO tool. Sensitivity analysis was done to determine if poor quality studies effected compliance. Galbraith plot was used to investigate statistical heterogeneity amongst studies. A random effects model was used to pool compliance.

Results Twenty-three studies were included in the review, of which 20 were included in the quantitative analysis. All the studies were cross sectional. The overall compliance with spectacle use was 40.14% (95% CI- 32.78-47.50). The compliance varied from 9.84% (95% CI=2.36-17.31) to 78.57% (95% CI=68.96-88.18). The compliance derived in sensitivity analysis was 40.09%. Reasons for non-compliance were broken/lost spectacles, forgetfulness, and parental disapproval.

Conclusion Appropriate remedial measures such as health education and strengthening vision care services will be required to address poor compliance with spectacle use among children.

Background

Refractive errors are a major cause of morbidity globally. Recent data shows that uncorrected refractive error is among the leading of moderate or severe vision impairment in the global population in 2015¹. It is estimated that in developing countries, 7% to 31% of childhood blindness and visual impairment is avoidable². In the age group of 5–15 years, nearly 12.8 million (0.97% global prevalence) children are visually impaired due to uncorrected or unsatisfactorily corrected refractive errors³. The projected cost of uncorrected refractive error (RE) described as direct and indirect loss of world productivity is 269 billion international dollars (I\$) (US\$ 202 billion), and the projected cost of addressing the issue is US\$ 28 billion over 5 years^{4–5}. RE in children due to myopia, amblyogenic hyperopia, astigmatism and anisometropia can present with blindness and visual impairment and warrants appropriate treatment^{5–6}. Visual disturbances in children lead to poor academic growth, injuries, reduced social participation, and functional impairment⁷. Correction of visual impairment with spectacles is the most cost-effective intervention for improving eye care and thus the productivity and functionality of children. Spectacles have a quality of being simple to use, non-invasive and inexpensive. However, the benefit of these visual aids depends on the compliance by end users.

REs have been given highest priority in various eye health programs globally. Very few studies are available worldwide to look into the factors determining compliance with spectacle use^{8–27}. Studies have shown that the compliance with spectacle use in children with visual impairment due to REs is only one third or less^{11,16,17,19,24}. Compliance remained low even when the spectacles were provided for free, and poorer rates were observed in older children^{14,18,21,23,24} and children residing in rural areas^{13,15}. Poor follow up after school-based screenings, broken spectacles, loss, forgetfulness^{9,11,13–15,17–20,23,24,27} parental and children's perceptions^{8,11,13,15,16,18,22–24}, peer pressure^{9,11,14,18,19,22,24}, safety concerns and the patient's self-esteem are few of the reasons cited for poor compliance. The health programs across countries have incorporated provision of free spectacles to school children as a component for prevention of blindness, but further measures of ensuring that the free spectacles are used are mostly missing.

Variable rates of compliance worldwide suggest that augmented attention is warranted, including investment in development and assessment of spectacle compliance interventions to assist in reducing complications associated with non-wear of spectacles. Based on literature review, we hypothesized that compliance with spectacle use would be low in children. The primary objective of the present review is to study the compliance with spectacle use in children with REs and to arrive at a summary measure of the rate of compliance by pooling data from various studies. The second objective was to assess the reasons for non-compliance in children with RE.

Methods

This systematic review was conducted in accordance with the Meta- analysis of Observational Studies in Epidemiology²⁸ (MOOSE) guidelines and the Preferred Reporting Items for Systematic Reviews and Meta-analyses standard (PRISMA)²⁹.

Literature Search

An extensive search was conducted on published literature and efforts were made to acquire information about the unpublished literature from various sources. The searches were performed on 28th May, 2017 and were updated on 31st March, 2018 on the following databases: Ovid, EMBASE, CINAHL and Pubmed. A thorough search of title and abstract (tiab) using the key words was performed. We did MeSH terms screening along with use of Boolean operators. The search terms were- ((((((Compliance [Title/Abstract]) OR Adherence[Title/Abstract]) OR Compliant[Title/Abstract]) OR Adherent[Title/Abstract])) AND (((Spectacle[Title/Abstract]) OR Spectacles[Title/Abstract]) OR Eye Glasses[Title/Abstract])) AND (((Child[Title/Abstract]) OR Children[Title/Abstract]) OR Adolescents[Title/Abstract]))).

Adolescent[Title/Abstract] OR Adolescents[Title/Abstract]). Reference lists of articles retrieved in the initial step were screened for pertinent studies. Efforts were made to contact authors for articles, which could not be obtained. We also searched additional platforms like Google search for non-indexed studies.

Inclusion Criteria

Observational and experimental study designs were included in the systematic review. These included cross sectional, case control and cohort study designs. Experimental study designs were also explored. Studies assessing compliance with spectacle use in children with REs, published in English language, with one or more of the key words in the title or abstract were included in the review. Compliance was defined as regular use of glasses prescribed for refractive errors including myopia, hypermetropia and astigmatism, assessed either by observation or by interviewing the children. Studies irrelevant to the objective of this review e.g. in children suffering from other eye disorders, conducted in adults, not published in English language were excluded from the review. The studies in which raw data were missing or unclear were excluded from quantitative analysis. We did not include conference proceedings. The participants comprised of children of both sexes with REs.

Data extraction

The studies were independently reviewed by two researchers (N.D, S.D), performed a thorough search of the databases and screened titles and abstracts based on the research question, and population and outcome in terms of compliance with spectacle use. Compliance was defined as regular use of glasses prescribed for refractive errors including myopia, hypermetropia and astigmatism, assessed either by observation or by interviewing the children. Based on the initial screening, full-text articles were obtained. Duplicates were removed in the initial stages. The third investigator (M.D) solved any disagreement in the selection of studies. Two reviewers conducted quality assessment independently using the QATSO tool used in previous studies.³⁰⁻³². No significant difference was found in individual assessments of the reviewers. This study has been registered with PROSPERO, with registration number CRD42017068190.

Data analysis

The data were entered separately from the included studies in a pre-designed and piloted format that recorded the information about author name, country of study, date of publication, study design, number of children, duration of follow up, type of study, percentage compliance and reasons for non-compliance.

After extraction, all related data were entered into Microsoft Excel for compilation. The data were analyzed with STATA MP 12 v11.³³. Pooled compliance estimate for spectacle use in children was generated using a forest plot. Compliance rates were calculated from raw proportions or percentages reported in the selected studies. The raw proportions/percentages were pooled using a random-effects model and pooled estimates and the 95% Confidence intervals (CI) were calculated. Correlation coefficient was calculated between per capita GDP expenditure and percentage expenditure on health with percentage compliance. In addition compliance was measured on basis of setting of the study i.e, whether the children were provided spectacles in screening settings in the field or in clinical setting. Sensitivity analysis was done to assess the effect of poor-quality studies on the overall compliance with spectacle use. Galbraith plot was used to investigate the statistical heterogeneity amongst the studies.

Results

Study Characteristics

A total of 2452 articles were retrieved, of which 33 articles met the study criteria for inclusion (Figure 1). After construing the full text, we finally included 23 studies for the review (Table 1). We did not include data for pooled compliance estimate from three studies^{8,26,34} as the proportion of children compliant with spectacle use was not available. However, descriptive data from these studies was included.

All the 23 studies included were cross-sectional in nature. The studies are geographically diverse and covered 14 countries. Most (34.7%) of the studies were from South-East Asian region^{8,10-12,15,19,23}, followed by Eastern Mediterranean region^{9,13,18,25} (17.4%). More number (39.1%) of studies was from lower-middle income country (LMIC) groups. The per capita GDP of the countries ranged from 835 USD for Nepal to 59532 USD for USA. Percentage expenditure on health ranged from 2.69 for Pakistan to 16.84 for USA. In most studies, compliance with spectacles was defined if the child was wearing spectacles at the time of surprise visit by the investigators or had the spectacles in bag (65.2%). For a few studies compliance was defined by taking interviews of the children and asking questions regarding their patterns of spectacle use (34.8%). The prescription patterns varied across studies. Some studies used the cut off of >0.5 Diopters for defining myopia and assessing compliance while a few defined myopia with a cut off of >1 Diopters. Similarly, for hyperopia the cut offs ranged from +1.0 D to 2.5 Diopters. A few studies defined the prescriptions and cutoffs in terms of visual acuity and included children with visual acuity <6/9 to <6/12 across studies.

Population characteristics

The total number of children studied was 7859. The children enrolled were from all age groups ranging from pre school to end of school. The period of follow up ranged from 'no follow up' to follow up of 18 months. The variation in ages was vast and contributed to significant heterogeneity.

Quality Assessment

The studies were assessed for the methodological quality based on the tool developed by Wong WC et al^{31,32}. Close to half (43.5%) of the studies were of good quality while 9 studies (39.1%) were of satisfactory quality and 4 studies (17.4%) were poor quality studies with score <30%. The tool used for quality assessment is as shown in eTable 1 (supplementary file). The results of quality assessment of studies are attached as a supplementary file (eTable 2).

Compliance with Spectacle use:

The overall compliance with spectacle use was 40.14% (95% CI- 32.78–47.50) for 20 studies. The compliance varied from 9.84%(95% CI = 2.36–17.31) to 78.57 %(95% CI = 68.96–88.18). Four studies had extreme values and their confidence intervals did not overlap those of other studies. Two of these studies reported very low compliance^{16,24} while 2 reported very high compliance^{23,25} with spectacle use. A forest plot of the studies and the compliance rate is as shown in Figure 2. The compliance across almost all income groups was poor. The correlation coefficient for relationship between per capita GDP and percentage compliance was -0.051 , indicating a decrease in compliance with an increase in per capita income. However, this was not statistically significant ($p = 0.830$). Similarly, the correlation coefficient for relation between percentage expenditure on health and percentage compliance was -0.238 , indicating that an increase in percentage expenditure on health was associated with a decrease in compliance. This relationship also, was not statistically significant ($p = 0.312$). The prescription cut offs used for measuring compliance had an effect on the compliance pattern across studies. Based on the setting of the study the compliance was pooled for screening vs clinical care. It was observed that the compliance was 45.84% in the setting of clinical care and 39.33 in the setting of screening (eFig.1). This difference was not statistically significant. The method of measurement of compliance also varied across groups. A few studies used observation/ surprise visit as a measurement method while others used interviews/questionnaires for measurement of compliance. We grouped and analyzed the studies by compliance measure. The pooled compliance where observation was the method of measurement, was 39.24% while the pooled compliance where interview was the method of measurement of compliance, the pooled compliance was 43.23% (eFig.2). This difference was not statistically significant, indicating that the measurement method did not significantly alter the compliance. A few studies that used a higher prescription cut off, reported better compliance as compared to studies that had lower cut offs for measurement of compliance, indicating lesser compliance at lower refractive errors. However most studies did not report any specific cut offs for prescriptions and some studies took cut offs using visual acuity, hence the pooled effect could not be observed.

Sensitivity Analysis and Investigation for heterogeneity

Sensitivity analysis was done to know whether the poor quality studies^{10,12,13,17} had any effect on the overall compliance. The overall compliance with spectacle use did not vary significantly and was 40.09% when these studies were removed from the analysis. The forest plot excluding these studies is as shown in Figure 3. The Galbraith plot (eFig.3) has shown that the studies had significant heterogeneity. The heterogeneity could be attributed to varying sample sizes and prescription cut offs (degree of refractive errors).

Reasons for Non-compliance with Spectacle Use

The most commonly cited reasons for non-compliance with spectacle use in the studies were broken^{9–12,14,15,17–20,23,24,27}/lost spectacles^{9–12,14,15,17–20,24,27}, forgetfulness^{9–11,14,15,17–19,24,27} and parental disapproval^{8,9,11,12,15,16,18,23,24,34}. These were followed by headache^{10–12,15,17,23,24,27}, teasing by peers^{10,12,19} and dislike for spectacles^{9,14,16–23}. Other reasons mentioned in a few studies were use only when required^{10,12} unclear vision^{11,12,22,27,34}, unattractive frames/poor appearance^{8,34}, fear of injuries^{8,34}, lack of affordability^{9,12,16}, uncomfortable spectacles^{18,34} and negative attitude of the society^{8,15,21} (eFig.4). The reasons for non-compliance were pooled for different studies, and is as shown in the forest plots (Figure. 4(a-d)). The reasons were broadly classified into personal factors, social factors, visual problems and breakage/loss/forgetfulness. The non-compliance due to personal factors was 25.78% (fig 4a), for social factors it was 13.18% (fig 4b), visual problems/headache it was 5.47% (fig 4c) and for breakage/loss/forgetfulness it was 23.34% (fig 4d) of the total. The results clearly show that personal factors and breakage/loss/forgetfulness were the most commonly cited reasons for noncompliance.

Discussion

Compliance with spectacle use is an important aspect in eye health of children. This systematic review and meta-analysis was conducted with a purpose of detecting the overall compliance rate with spectacle use in children prescribed spectacles and to identify the reasons for non-compliance. Twenty-three studies were selected after thorough literature search and of these 20 studies were included in the quantitative analysis (meta-analysis). The overall pooled estimate shows that the compliance with spectacle use is considerably low among children 40.14% (95% CI- 32.78–47.50) as assessed from 20 studies, though the studies were heterogeneous. The most commonly reported reasons for non-compliance were broken glasses, lost spectacles, forgetfulness and parental disapproval complaints of visual disturbances and headache.

Majority of the studies were from South East Asia Region (SEAR) and lower middle-income countries. There was a dearth of studies from low-income countries, which could have altered the results and improved representativeness. The estimated compliance rate with spectacle was less than half (40.14%). Sub optimal compliance is a point of concern and can lead to progression of refractive errors. Negative, non significant correlation of percentage compliance with per capita GDP and percentage expenditure on health, suggests that apart from pecuniary factors, psychosocial factors may be a contributor to compliance. It was observed that the setting (screening vs clinical care) and the method of assessing compliance (interview vs observation) did not have significant effect on the compliance. Although, pooled compliance in the clinical care setting where problem was identified by the child or family compared to vision screening setting provided at school was not significant, but the results point to of better compliance in clinical care setting, thus indicating that they are more likely to be aware and are motivated to use the spectacles. Personal, behavioral and cultural factors influence children's compliance with spectacle use. The rate was less than half for most countries except a few which showed extremes on the lower and higher side. Compliance rate was overlapping between

almost all studies except for two studies which reported a very high compliance^{23,25} and two that reported low compliance^{16,24}. In the study by Khandekar *et al*²³ (2008), the compliance may be high for two reasons. First, the sample size was small (77), that may have been a contributor to the difference in the compliance value from other studies. Second, the cut off for spectacle compliance was >0.75D error, hence free glasses were given to children with higher refractive error, which itself is a factor for good compliance. In the study by Khandekar *et al*²⁵(2002), the students were being assessed for compliance at regular intervals, thus improving the compliance rate. On the other hand, the reasons reported by Megbeylian *et al*¹⁶ (2013) for poor compliance were lack of affordability and deep-rooted customs /traditions. In Nigeria, the expenditure on health as percentage of GDP and the HDI are low, which may be other contributory factors to affordability. The prescription cut offs for spectacle compliance assessment also varied across studies, though most studies did not report specific cut offs, a few showed that the compliance was poorer in children who had lower refractive errors as compared to children with higher refractive errors. Increased severity of refractive error warrants a stricter compliance to spectacle use as it hinders daily activities due to poor visibility. Some studies^{14,16,19,24,27} reported better compliance in myopia as compared to hypermetropia. Recent study by Mc Cormick I et al (2018)³⁵ also reported better compliance with higher refractive error in their study on determinants of compliance to spectacle use. Due to different definitions and cut offs we could not find an effect of these factors on the compliance as some studies have mentioned the prescription cut offs in terms of diopters while others have measured it in terms of visual acuity. In addition the available data was also not uniform to be pooled together.

Identifying reasons for non-compliance with spectacle use is important for understanding the social determinants for intervention. The most commonly reported reasons for non-compliance were broken glasses^{9-12,14,15,17-20,23,24,27}, forgetfulness^{9-11,14,15,17-19,24,27}, loss of spectacles^{9-12,14,15,17-20,24,27} and parental disapproval^{8,9,11,12,15,16,18,23,24,34}. (Fig. 4 a- 4d) Addressing these issues by generating awareness is imperative. The outcomes indicate more of socio cultural factors as major contributors to poor compliance, that are commonly seen in Upper middle income countries. Any habit, if inculcated in the early years of life is bound to show results in adulthood. Most of the reasons identified for poor compliance are modifiable and are due to carelessness and poor encouragement of children. Breakage, loss and forgetfulness are intervention points that can bring substantial difference in the compliance rate. Parental disapproval is a significant contributor to child behavior. Parents usually stop their children from wearing spectacles, as they are concerned with the societal opinions. In many low and middle income countries, spectacles are considered a sign of weakness and their use hinders the process of finding a suitable match for the children when they reach adulthood. Many parents are seeking LASIK as a procedure for permanent removal of eyeglasses by surgical correction. Factors related to visual problems and headache can be addressed by modification of the prescription glasses and appropriate correction till comfort is achieved.

Compliance with spectacle use in children is an often-neglected issue. Unless social and perceptual barriers are overcome, the families will not access the financial and logistical assistance available to seek eye care for school-aged children. Behavior change communication (BCC) targeted at education and behavior change of parents, so they encourage children to use spectacles, is advocated. Another point of action could be school health programs, which should focus on incorporating the component of ensuring compliance through follow-up, apart from screening of children for refractive errors. Some potential actions that we think may address the poor compliance found in our study include provision of spectacles to children at zero cost, promotion of school vision screening programs by government, involvement of teachers in the process of identifying children not compliant and ensuring remedial actions. Parental education and support are key pillars to strengthen the interventions. It is unlikely that uni-dimensional intervention approaches to increase follow up and spectacle adherence in the context of refractive errors (e.g., free spectacles) will be adequate to achieve sustained improvement in treatment outcomes among school children. Positive reinforcement is essential at both the school and household levels. Generating awareness and glamourizing spectacles by using lightweight, unbreakable and trendy frames will promote their acceptance, especially in adolescent age groups. Lastly, the role of eye care practitioners is imperative in early identification, diagnosis and treatment of refractive errors among children, so as to curb the problem at a very nascent stage.

There are a number of strengths of this study. First, the present review is a first systematic review on spectacle compliance in children. No systematic review or meta-analysis has been conducted previously on this topic. Efforts have been made to include all the available studies on the topic. Secondly, no time restriction was imposed and we have obtained studies for all years. There are a few limitations of the study as well. Despite all our efforts to extract maximum number of studies, we may have missed relevant studies in unpublished literature (publication bias). Also, the number of studies obtained was mostly from middle-income countries and a clearer picture of influential factors from high income and low-income countries would not be made very clear.

Conclusion

In conclusion, this systematic review and meta-analysis has thrown light upon compliance with spectacle use and reasons poor compliance with spectacles which are a low cost intervention for refractive errors management in children. This issue needs to be addressed through behavioral motivation of children, parents and the society. This review can help stakeholders and program managers in defining health care interventions to improve compliance with spectacle use.

Abbreviations

Abbreviation	Full form
RE	Refractive Error
MOOSE	Meta- analysis of Observational Studies in Epidemiology
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-analyses
CI	Confidence Interval
LIC	Low Income Country
LMIC	Lower Middle Income Country
UMIC	Upper Middle Income Country
HIC	High Income Country
SEAR	South East Asia Region
GDP	Gross Domestic Product
HDI	Human Development Index
BCC	Behavior Change Communication

Declarations

Ethics approval and consent to participate: Not Applicable

Consent for publication: Not Applicable

Competing interests: The authors declare that they have no competing interests.

Data Availability: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Funding: None

Authors' contributions

Conceptualization: ND, SD, MD

Data extraction: ND, SD, NJ, MD

Formal analysis: ND, SD, NJ, MD, PCG

Software: NJ,ND

Supervision: MD,PCG,MS,JR

Writing original draft: ND,SD.

Writing, review & editing: MD,PCG,NJ,MS,JR

Approval of final version: ND,SD,MD,NJ,PCG,MS,JR.

Acknowledgement: ICMR Centre for Advanced Research in Evidence Based Child

Health, Advanced Pediatrics Centre, PGIMER for providing the support in the form of

Software to carryout analysis and in searching the literature.

References

1. Flaxman SR, Bourne RRA, et al. Global causes of blindness and distance vision impairment 1990–2020: a systematic review and meta-analysis. *Lancet Glob Health* 2017;5(12):e1221-e34.
2. Kong L, Fry M, Al-Samarraie M, et al. An update on progress and the changing epidemiology of causes of childhood blindness worldwide. *J AAPOS*, 16(6), 501–507.
3. Global magnitude of visual impairment caused by uncorrected refractive errors in 2004. *Bulletin WHO* 2008;86(1):63–70
4. WHO. Blindness avoided for millions of children [Internet]. Available from http://www.who.int/features/2009/preventing_blindness/photo_story/en/ Accessed June 27,2017

5. Smith TST, Frick KD, Holden BA, Fricke TR, Naidoo KS. Potential lost productivity resulting from the global burden of uncorrected refractive error. *Bull World Health Organ.* 2009;87(6):431–437.
6. Fricke TR, Holden BA, Wilson DA et al. Global cost of correcting vision impairment from uncorrected refractive error. *Bull World Health Organ.* 2012;90(10):728–738.
7. Holden BA, Fricke TR, Wilson DA et al. Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. *Ophthalmology* [Internet]. 2016;123(5):1036–1042. Available from: <http://dx.doi.org/10.1016/j.ophtha.2016.01.006>. Accessed on June 27, 2017.
8. Narayanan A, Kumar S, Ramani KK. Spectacle Compliance among Adolescents: A Qualitative Study from Southern India. *Optom vis sci.*2017;94(5):582–587.
9. Anwar I, Waqar S, Altaf A. Spectacle Wear Among School Going Children in District Rawalpindi, Pakistan. *Int J Ophthalmol Vis Sci.* 2017;2(1):1–4.
10. Bhatt NK, Rathi M, Dhull CS, Sachdeva S, Phogat J. Spectacle compliance amongst school children of Rohtak, Harayan, India. *Int J Community Med Public Health.* 2017;4(3):734–737.
11. Bhandari G, Pradhan S, Shrestha M, Bassett K. Visual System Eye Glasses Compliance among Children Undergoing School Visual Acuity Screening in Nepal. *Adv Ophthal visual sys* 2016;5(3):1–5.
12. Kumar MR, Mallika O. U. Study of refractive Errors, Amblyopia and Compliance of Spectacles in School Children. *J Med Sci Clin Res.*2017;5(4):1–8.
13. Al-shamarti SA. Factors Associated With spectacle-wear compliance among hypermetropic pre-school children in Al-diawaniya Governorate, Iraq. *AL oadisiva medical journal*2015;11(20):170–175.
14. Barria von-Bischhoffshausen F, Muñoz B, Riquelme A, Ormeño MJ, Silva JC. Spectacle-Wear Compliance in School Children in Concepción Chile. *Ophthalmic Epidemiol.*2014;21(6):362–369.
15. Pavithra M, Hamsa L, Madhukumar S. Factors associated with spectacle-wear compliance among school children of 7–15 years in South India. *Int J Med Public Heal* 2014;4(2):146.
16. Megbelayin EO. Barriers to uptake of prescribed refractive spectacles amongst Nigerian students. 2013;1(May):71–77.
17. Turcin L-A, Jompan A. Compliance of the Students To the Refractive Errors Correction With Glasses. *Stud Univ Vasile Goldiș Ser Științele Vietii.* 2013;23(3):257–263.
18. Aldebasi YH. A descriptive study on compliance of spectacle-wear in children of primary schools at Qassim Province, Saudi Arabia. *Int J Health Sci (Qassim).* 2013;7(3):291–299.
19. Gogate P, Mukhopadhyaya D, Mahadik A et al. Spectacle compliance amongst rural secondary school children in Pune district, India. *Indian J Ophthalmol.* 2013;61(1):8–12
20. Messer DH, Mitchell GL, Twelker JD, Crescioni M. Spectacle wear in children given spectacles through a school-based program. *Optom Vis Sci.* 2012;89(1):19–26.
21. Keay L, Zeng Y, Munoz B, He M, Friedman DS. Predictors of early acceptance of free spectacles provided to junior high school students in China. *Arch Ophthalmol.* 2010;128(10):1328–34.
22. Odedra N, Wedner SH, Shigongo ZS, Nyalali K, Gilbert C. Barriers to Spectacle Use in Tanzanian Secondary School Students. *Ophthalmic Epidemiol.* 2008;15(6):410–417.
23. Khandekar R, Sudhan A, Jain BK, Tripathy R, Singh V. Compliance with spectacle wear and its determinants in school students in Central India. *Asian J Ophthalmol.* 2008;10(3):174–177.
24. Castanon Holguin AM, Congdon N, Patel Net al. Factors associated with spectacle-wear compliance in school-aged Mexican children. *Investig Ophthalmol Vis Sci.* 2006;47(3):925–928.
25. Khandekar R, Mohammed AJ, Raisi A Al. Compliance of spectacle wearand its determinants among schoolchildren of Dhakhiliya region of Oman:A descriptive study. *Squ J Sci Res med sci.*2002;4:39–43.
26. Horwood AM. Compliance with first time spectacle wear in children under eight years of age. *Eye (Lond).*1998;12:173–178.
27. Congdon N, Zheng M, Sharma A et al. Prevalence and determinants of spectacle nonwear among rural Chinese secondary schoolchildren: the Xichang Pediatric Refractive Error Study Report 3. *Arch Ophthalmol.*2008;126(12):1717–1723.
28. Stroup DF, Berlin JA, Morton SC et al. Meta-analysis of Observational Studies. *JAMA* 2000; 283(15):2008–2012
29. Moher D, Liberati A, Tetzlaff J, Altman DG, Group TP. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLOS Med.*2009;6(7):e1000097.
30. Wong WCW, Cheung CSK, Hart GJ. Development of a quality assessment tool for systematic reviews of observational studies (QATSO) of HIV prevalence in men having sex with men and associated risk behaviours. *Emerg Themes Epidemiol.*2008;4:3–6.
31. Roman N V, Frantz JM. The prevalence of intimate partner violence in the family : a systematic review of the implications for adolescents in Africa. *Fam Pract.* 2013:256–65.
32. Louw QA, Morris LD, Grimmer-somers K. The Prevalence of low back pain in Africa : a systematic review. *BMC Musculoskel Disord.*2007;14:1–14.
33. STATA: Data analysis and statistical software [Internet]. Texas,USA: StataCorp LLC; Available from: <http://www.stata.com>.Accessed on 29th June,2017.
34. Li L, Lam J, Lu Y et al. Attitudes of students, parents, and teachers toward glasses use in rural China. *Arch Ophthalmol.* 2010;128(6):759–65
35. McCormick I, Morjaria P, Mactaggart I, et al.Spectacle Compliance and Its Determinants in a School Vision Screening Pilot in Botswana. *Ophthalmic Epidemiol.*2019;26(2):109–116

Tables

Table 1: Characteristics of selected studies included in systematic review and meta-analysis (N=23)

S. No	Authors	Country	Age	Year	Study Design	WHO Region ^a	World Bank Income ^b	Per capita GDP (2017 in \$)	HDI	Expenditure on Health (% of GDP)	Compliance Measure	Type of prescription	Du of I up
1.	Narayanan A et al	India	13-17 years	2017	Cross Sectional (Qualitative)	SEAR	LMIC	1940	0.640	3.89	Questionnaire/ Interview based	No specific type	
2.	Bhatt NK et al	India	6-15 years	2017	Cross Sectional	SEAR	LMIC	1940	0.640	3.89	Wearing spectacles on surprise visit	No specific type	3 m
3.	Anwar I et al	Pakistan	11-16 years	2017	Cross Sectional	EMR	LMIC	1548	0.562	2.69	Questionnaire/ Interview based	Myopia, hypermetropia, astigmatism and compound	8-12 months
4.	Kumar MR et al	India	6-17 years	2017	Cross Sectional	SEAR	LMIC	1940	0.640	3.89	Wearing spectacles on surprise visit	Children with visual acuity <6/9	6 m
5.	Bhandari G et al	Nepal	5-16 years	2016	Cross Sectional	SEAR	LIC	835	0.574	6.15	Wearing spectacles on surprise visit or in bag	No specific type	No up
6.	Sumana M et al	India	9-16 years	2015	Cross sectional	SEAR	LMIC	1940	0.640	3.89	Wearing spectacles on surprise visit	No specific type	6 m
7.	Al Shamarti S et al	Iraq	Pre-school	2015	Cross Sectional	EMR	UMIC	5166	0.685	3.40	Questionnaire /Interview based	No specific type	Not me
8.	Von-Bischhoffs-hausen FB et al	Chile	4-19 years	2014	Cross Sectional	AMR	HIC	15346	0.843	8.07	Wearing spectacles on surprise visit	Visual Acuity 20/40 in either eye, myopia 0.75 diopters (D), 1.5D of astigmatism, or hyperopia.	12 m
9.	Pavithra MB et al	India	7-15 years	2014	Cross Sectional	SEAR	LMIC	1940	0.640	3.89	Wearing spectacles on surprise visit	Children with myopia ≥ -0.5 spherical equivalent diopters in one or both eyes, hypermetropia $\geq +1.00$ spherical equivalent diopters in one or both eyes and astigmatism ≥ 1.00 D	3 m
10.	Gogate P et al	India	8-16 years	2014	Cross Sectional	SEAR	LMIC	1940	0.640	3.89	Wearing spectacles on surprise visit followed by interview	myopia ≥ -0.50 D or hyperopia $\geq +1.00$ D	6-12 months
11.	Aldebasi YH et al	Saudi Arabia	7-13 years	2013	Cross Sectional	EMR	HIC	20761	0.853	5.83	Wearing spectacles on surprise visit or in bag	A child with myopia of more than -0.50 Dioptre (D) or hypermetropia of more than +1.00D are prescribed spectacles	6 m
12.	Megbelayin EO et al	Nigeria	9-21 years	2013	Cross Sectional (mixed methods)	AFR	LMIC	1969	0.532	3.56	Wearing spectacles on surprise visit	VA <6/9 in at least one eye	No up
13.	Turcin LA et al	Romania	6-11 years	2013	Cross Sectional	EUR	UMIC	10814	0.811	4.95	Wearing spectacles on surprise visit or in bag	Myopia (≥ -1.00 DS in one or both eyes) Hyperopia	12 m

												($\geq +1.50$ DS in one or both eyes) Astigmatism (values ≥ 1.00 D)	
14.	Messer DH <i>et al</i>	USA	8-14 years	2012	Cross Sectional (mixed methods)	AMR	HIC	59532	0.924	16.84	Wearing spectacles on surprise visit	Myopia ≥ -0.75 D, astigmatism ≥ 1.00 DC, and anisometropia ≥ 1 D and hyperopia $\geq +2.50$ D	3 m
15.	Keay L <i>et al</i>	China	12-15 years	2010	Cross Sectional	WPR	UMIC	8827	0.752	5.32	Wearing spectacles on surprise visit or in bag	No specific type	1 m
16.	Li L <i>et al</i>	China	14-18 years	2010	Cross sectional (Qualitative)	WPR	UMIC	8827	0.752	5.32	Questionnaire/interview based	No specific type	
17.	Khandekar R <i>et al</i>	India	School going	2008	Cross Sectional	SEAR	LMIC	1940	0.640	3.89	Questionnaire/Interview based	Myopia (>0.75 D spherical unilateral or bilateral)	3-4 months
18.	Congdon N <i>et al</i>	China	11.4-17.1 years	2008	Cross Sectional	WPR	UMIC	8827	0.752	5.32	Questionnaire/Interview based	Uncorrected VA of 6/12 or worse and a 25% random sample of subjects with VA better than 6/12	No up
19.	Li L <i>et al</i>	China	Average 14.7 years	2008	Cross Sectional	WPR	UMIC	8827	0.752	5.32	Questionnaire/Interview based	All children with presenting VA 6/12 in either eye and children with spectacles improving vision to 6/12	3 m
20.	Odedra N <i>et al</i>	Tanzania	Average 14 years	2008	Cross Sectional (mixed methods)	AFR	LIC	936	0.538	6.12	Wearing spectacles on surprise visit	VA worse than 6/12 in either eye or hyperopia of 2 diopters (D) or more	3 m
21.	Holguin AM <i>et al</i>	Mexico	6-18 years	2006	Cross Sectional	AMR	UMIC	8903	0.774	5.86	Wearing spectacles on surprise visit	Children with visual acuity 6/12 or less	4-12 months
22.	Khandekar R <i>et al</i>	Oman	6-17 years	2002	Cross Sectional	EMR	HIC	15668	0.821	3.83	Wearing spectacles on surprise visit	No specific group	12 m
23.	Horwood AM <i>et al</i>	UK	1-7 years	1998	Cross Sectional	EUR	HIC	39720	0.922	9.88	Questionnaire/interview based	No specific group	Not measured

^aSEAR: South East Asia Region; EMR: Eastern Mediterranean Region, WPR: Western pacific region; AFR: African region; AMR: Region of the Americas

^bUMIC: Upper middle income country; LMIC: lower middle income country; LIC: Lower income country; HIC: High income country

Empty cells depict missing information.

VA: Visual acuity

Figures




Figure 1

Flow of selection of studies for the review




Figure 2

Compliance to Spectacle use (%) as reported by selected studies (N=20)




Figure 3

Sensitivity analysis showing compliance to spectacle use (%) after removal of poor quality studies (N=16)




Figure 4

Reasons for non compliance to spectacle use

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryfiles1.docx](#)