

# Analysis of ADAM9 Regulation and Function in Vestibular Schwannoma Primary Cells

**Anja Nattmann**

Universitätsklinikum Würzburg Neurochirurgie

**Maria Breun** (✉ [Breun\\_M@ukw.de](mailto:Breun_M@ukw.de))

Universitätsklinikum Würzburg Neurochirurgie

**Camelia Monoranu**

Universitätsklinikum Würzburg Pathologie

**Cordula Matthies**

Universitätsklinikum Würzburg Neurochirurgie

**Ralf-Ingo Ernestus**

Universitätsklinikum Würzburg Neurochirurgie

**Mario Löhr**

Universitätsklinikum Würzburg Neurochirurgie

**Carsten Hagemann**

Universitätsklinikum Würzburg Neurochirurgie

---

## Research note

**Keywords:** Vestibular schwannoma, pathogenesis, ADAM9, knock down, Integrin, Immunofluorescence double staining, Merlin, primary cell culture

**DOI:** <https://doi.org/10.21203/rs.3.rs-71416/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Objective.** Recently, we described a disintegrin and metalloproteinase 9 (ADAM9) overexpression by Schwann cells of vestibular schwannoma (VS) and suggested that it might be a marker for VS tumor growth and invasiveness. This research note provides additional data utilizing a small cohort of VS primary cultures and tissue samples. We examined whether reconstitution of Merlin expression in VS cells regulates ADAM9 protein expression and performed lentiviral ADAM9 knock down to investigate possible effects on VS cells numbers. Moreover, the co-localization of ADAM9 and Integrins  $\alpha 6$  and  $\alpha 2\beta 1$ , respectively, was examined by immunofluorescence double staining.

**Results.** ADAM9 expression was confirmed in VS but not regulated by Merlin. However, ADAM9 knock down led to significant 2.38 fold reduced cell numbers in VS primary cell cultures ( $p < 0.001$ ). While ADAM9 and Integrin  $\alpha 2\beta 1$  were co-localized in only 22% (2 of 9) of VS, ADAM9 and Integrin  $\alpha 6$  were co-localized in 91% (10 of 11) of VS. Therefore, we provide first observations on possible regulatory functions of ADAM9 expression in VS.

## Introduction

Vestibular schwannoma (VS) are benign tumors emerging from Schwann cells of the vestibular part of the 8th cranial nerve [1]. The hallmark mutation is loss of Merlin function, a 4.1 protein/ezrin/radixin/moesin protein (FERM) connecting the cytoskeleton with the cell membrane. Merlin is activated by the cells' attachment to the extracellular matrix (ECM) and intercellular adhesion [2]. Recently, we showed that a disintegrin and metalloproteinase 9 (ADAM9) mRNA-expression is 8.8 times higher in VS compared to healthy vestibular nerves and we suggested that ADAM9 inhibition may be of significance for VS pathogenesis and potential medical treatment [3]. Furthermore, we discussed whether ADAM9 might interact with the VS cell-matrix [3], since it is known to promote tumor invasion by such interactions [4]. In particular, this enzyme liberates membrane-bound proteins by an enzymatic modification called "shedding" and therefore plays a role in tumor growth and invasiveness [4].

This research note builds on our recent findings by describing three new experiments utilizing a small cohort of VS primary cultures and tissue samples. We examined whether reconstitution of Merlin expression in VS cells regulates ADAM9 protein expression. We performed an ADAM9 mRNA knock down and investigated its influence on VS primary cell number. Finally, we investigated co-localization of ADAM9 with Integrin  $\alpha 6$  and Integrin  $\alpha 2\beta 1$ , respectively, by immunofluorescence double staining of VS tumor samples.

## Material And Methods

### *Tissue samples and cell culture*

Tissue of 24 VS, surgically resected in the Department of Neurosurgery of the University Hospital Würzburg, was collected from January 2018 until July 2019. Half of each sample was embedded in

paraffin for immunohistochemistry, the other half was processed for primary cell culture as described elsewhere [5]. Briefly, the fresh tumor samples were washed three times with ice-cold HBSS +/+ (Gibco, Carlsbad, CA, USA), shred into 0.2 × 0.2 cm small pieces and digested with 1.7 mg/ml dispase (PluriSTEM, Millipore, Massachusetts, USA) and 1.5 mg/ml collagenase (Sigma, Munich, Germany) in 6 ml VS-medium (DMEM supplemented with 10% heat-inactivated fetal bovine serum, 1% penicillin/streptomycin (all from Gibco, Carlsbad, CA, USA), 0.35% Glucose (B. Braun, Melsungen, Germany), 8 nM Heregulin (Stemcell Technologies, Cologne, Germany), 0.5 mM IBMX and 0.25 µg/ml insulin (both from Sigma, Munich, Germany)) for 17 h at 37°C and 5% CO<sub>2</sub>. Afterwards, the suspension was centrifuged for 5 min at 300 ×g, the supernatant was discharged and the pellet resuspended in 12 ml VS-medium, before plating 2 ml into each well of laminated 6-well-plates [5]. The cells were cultured in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C.

### ***Merlin overexpression and ADAM9 knock down***

95,000 cells per well of the primary cell cultures were treated with 2 ml VS-medium containing 8 µg/ml protamine (Sigma, Munich, Germany) and 100 µl NF2 transcript variant 1 (NM\_000268) Human mGFP Tagged ORF Clone Particle RC205883L2 (Merlin overexpression), 57 µl ADAM9 Human shRNA Lentiviral Particle TL314947VB (Locus ID 8754) (ADAM9 knock down) or 120 µl Lenti-shRNA Control Particles TR30021V (scrambled control) (all from OriGene, Rockville, MD, USA) and incubated for 2 (Merlin overexpression) or 3 days (shRNA). On the next day the medium was replaced by 2 ml fresh VS-medium, the Merlin overexpressing cells were photographically documented and lysed as described below. The shRNA transfected cells were cultured for another 24 h. Their medium was replaced by 2 ml VS-medium containing 1 µg/ml puromycin (Gibco, Carlsbad, CA, USA) on day 5, which was exchanged every 2 days by DMEM. At day 11 five fields of view of each well were photographed through the 10× magnifying objective of the DMI 3000B fluorescence microscope and the DFC 450C camera (Leica, Wetzlar, Germany) in bright field with 35 ms exposure time and for green fluorescence using filter L5 ET and an exposure time of 900 ms. Cells were counted supported by usage of the program ImageJ (National Institutes of Health (NIH), Bethesda, MD, USA; <http://imagej.nih.gov/ij/>).

### ***Western-blot***

After washing the cells twice with phosphate buffered saline (PBS; Biochrom, Berlin, Germany), they were lysed with 100 µl lysis buffer (10 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM ethylenediaminetetraacetic acid (EDTA), 1 mM ethylene glycerol-bis (β-aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA), 1% Triton X-100, 0.5% IGEPAL CA-630, 1 mM phenylmethanesulfonylfluoride (PMSF), 10 µg/ml leupeptin and 23 µg/ml aprotinin (all from Sigma, Munich, Germany)). The extracted protein was measured with the Qubit 2.0 Fluorometer (Thermo Fisher Scientific, Waltham, MA, USA). Polyacrylamid gel electrophoresis (PAGE) was performed as described [6] and the gel blotted for 7 min using the iBlot system (Thermo Fisher Scientific, Waltham, MA, USA) set to program 3. The membrane was blocked in TBST (Sigma, Munich, Germany) containing 5% non-fat milk powder (Roth, Karlsruhe, Germany) and probed with antibodies as described previously [6]. Antibodies NF2 B-12 sc-55575 (Santa Cruz Biotechnology, Dallas, TX, USA),

ADAM9 ab186833 (Abcam, Cambridge, UK), and anti- $\alpha$ -tubulin T6557 (Sigma, Munich, Germany) were utilized diluted in TBST at 1:200, 1:1500, and 1:5000, respectively. Goat anti-rabbit IgG H&L (HRP) ab6721 (Abcam, Cambridge, UK), anti-mouse IgG HRP NA931 (GE Healthcare, Freiburg, Germany) and anti-mouse m-IgGk BP-HRP sc516102 (Santa Cruz Biotechnology, Dallas, TX, USA) antibodies were used as secondary antibodies at a 1:1000 dilution in TBST. The ECL Western Blotting Analysis System (Amersham, Freiburg, Germany) was used to visualize the antibody labeled proteins.

### ***Immunofluorescence double-staining***

Immunofluorescence staining of 3  $\mu$ m thick formalin-fixed paraffin sections has been described previously [6, 7]. However, blocking with 10 % goat serum (Life Technologies, Waltham, MA, USA) was performed for 2 h prior to an incubation of the slides with antibody dilution buffer (DCS, Jena, Germany) containing antibodies anti-ADAM9 ab186833 (1:100) in combination with anti-Integrin  $\alpha$ 2 $\beta$ 1 [16B4] ab30483 (1:50) or anti-Integrin  $\alpha$ 6 [MP 4F10] ab20142 (1:50) (all from Abcam, Cambridge, UK) at 4°C overnight. Protein expression was visualized by 1 h incubation using Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (Alexa Fluor Plus 488 (A32732) and 555 (A32732); Thermo Fisher Scientific, Waltham, MA, USA), both diluted 1:400. Slides were mounted using Fluoroshield mounting medium, containing DAPI (Abcam, Cambridge, UK) and photographed with the Leica microscope DMI 3000B and the DFC 450C camera, using three different filters (Filtercubes A, L5 ET, and TXR ET) with exposure times of 77 ms, 2.5 s and 1.5 s, respectively, at both 10 and 40 $\times$  objective magnification.

### ***Statistical analysis***

Statistical analysis was performed with GraphPad Prism 6 software (GraphPad Software, La Jolla, CA, USA) to determine significance using unpaired t-tests.  $p < 0.05$  was considered to be statistically significant.

## **Results**

All analyzed VS were tested for ADAM9 protein expression by Western-blotting (Fig. 1a) and immunofluorescence (Fig. 2). Without exception, high amounts of ADAM9 were detectable in all VS. Since Merlin loss is a hallmark of VS development [8], we wondered whether the ADAM9 expression was due to lacking regulation by Merlin. Therefore, we restored Merlin expression by lentiviral transfection (Fig. 1b) and checked for any alteration in ADAM9 expression utilizing Western-blot. ADAM9 expression levels, however, did not significantly differ in cells with or without Merlin expression (Fig. 1c), ruling out Merlin-mediated ADAM9 regulation. Next, we asked whether ADAM9 expression might be involved in the proliferation of VS cells. Therefore, a lentiviral shRNA-mediated ADAM9 knock down was performed. It led to a statistically significant 2.38 fold reduction of VS cell numbers in comparison to the scrambled controls in primary cell cultures ( $p < 0.001$ ) (Fig. 1d).

ADAM9 is a cell-matrix modifying enzyme, reported to interact with Integrin  $\alpha$ 6 $\beta$ 1 and thereby regulating cellular motility [9]. This integrin is also expressed by VS cells [9] and therefore could be associated with

ADAM9 in VS. Indeed, Integrin  $\alpha 6$  expression could be found in 10 of 11 investigated VS (91%) and was co-localized with ADAM9 (Fig. 2). This co-localization was mainly found in connective tissue (Fig. 2a, b), near blood vessels (Fig. 2c) and enriched along the tumor capsule (Fig. 2e-f). In contrast, Integrin  $\alpha 2\beta 1$  was only weakly expressed in the analyzed VS (Fig. 2g-h). Strong expression with scattered ADAM9 co-localization was observed in merely two of 9 VS (22%) (Fig. 2g).

## Discussion

We were the first to describe ADAM9 overexpression in VS, suggesting it might be a marker for tumor growth and invasiveness [3]. However, our study was descriptive in nature and we only could speculate on ADAM9 regulation and function. Here, we performed first experiments to address these questions. We confirmed expression of ADAM9 in the analyzed VS, showed that Merlin overexpression does not regulate expression of the 47 kDa ADAM9 isoform, but that an ADAM9 knock down resulted in reduced VS cell numbers. Furthermore, we demonstrated ADAM9 co-localization with Integrin  $\alpha 6$  in 91% of the analyzed VS, but with Integrin  $\alpha 2\beta 1$  only in 22% of the cases.

Merlin modulates cell-cell interactions and cell migration. It is a tumor suppressor gene, with loss of function mutations in VS [10], leading to overexpression of Neuregulin 1 (NDRG1), which in turn activates ERK- and AKT-signaling pathways in VS, promoting VS cell proliferation [11–13]. Since we observed overexpression of ADAM9 in VS, we wondered whether this may be due to the loss of Merlin in these cells and restored Merlin expression. However, our data suggest that there is no causal link between loss of Merlin and overexpression of at least the proteolytically-processed, membrane-bound 47 kDa isoform of ADAM9 [4]. However, the knock down of ADAM9 led to a reduction of VS cell numbers. ADAM9 inhibition had already been shown to reduce migration and invasion of human glioma cell lines [14]. Therefore, our data might be a first hint that ADAM9 plays a similar role in VS and its expression might induce progression or possibly reduce cell death in these benign tumors.

ADAM9 is a cell membrane spanning protein connecting the cytoskeleton with the ECM or neighboring cells. Its cysteine-rich domain binds to proteoglycans and its disintegrin domain to integrins [4]. Metalloproteinases of the ADAM-family digest proteins of the ECM, allowing tumor cells to detach from the tissue and thereby promoting their migration and invasion [4, 15] Thus, highest ADAM9 expression could be detected in areas of liver metastases with high invasive growth [4]. Integrins mediate attachment of cells to the basal lamina [16]. Integrin  $\alpha 6$  and Integrin  $\alpha 2\beta 1$  are both known substrates of ADAM9. Interaction of ADAM9 and Integrin  $\alpha 2\beta 1$  has been described for liver metastases [4] and of ADAM9 and Integrin  $\alpha 6\beta 1$  for fibroblasts [9]. However, their association had not yet been shown for VS. Since we found co-localization of ADAM9 and Integrin  $\alpha 2\beta 1$  in only 22% of the analyzed VS, we conclude that this is not the main substrate of ADAM9 in VS. On the other hand, Integrin  $\alpha 6$  is detectable in Schwann cells [17]. Its precursor has higher expression levels in VS cells in comparison to Schwann cells [18] and we found co-localization in 91% of the VS with highest levels in connective tissue and near blood vessels. ADAM9 knock-out mice display a significant reduction of retinal neovascularization, whereas ADAM9 was highly expressed in pathological retinal blood vessels of wild type mice [19]. It has been suggested

that hypoxia induces ADAM9 expression and this enhanced expression in turn leads to release of pro-angiogenic factors [19]. Our data may support such assumption and could indicate that VS may require neoangiogenesis for their growth, although they are generally considered to be weakly vascularized tumors [20].

Our new data supplement our previous observations on ADAM9 expression in VS, are a first glimpse on possible regulatory functions and provide a rationale for more in-depth future investigations.

## Limitations

VS are slowly growing, benign tumors. Thus, it is a challenge to establish primary cell cultures [5] and even more difficult to achieve high transfection rates. For this reason, only a small number of experiments could be performed for protein knock down and overexpression, limiting generalizability of these experiments. Several ADAM9 isoforms have been described [4, 21, 22]. However, the antibody used in this study was specific for the 47 kDa isoform only. Therefore, we cannot conclude on a possible Merlin mediated regulation of other ADAM9 isoforms. The immunofluorescence double staining indicates co-localization of ADAM9 and Integrin  $\alpha 6$  or  $\alpha 2\beta 1$ , which does not necessarily mean that there is direct protein-protein interaction.

## Abbreviations

ADAM9: a disintegrin and metalloproteinase 9; DAPI: 4',6-diamidin-2-phenylindol; DMEM: Dulbeccos's modified Eagle's medium; ECM: extracellular matrix; EDTA : ethylenediaminetetraacetic acid; EGTA: ethylene glycerol-bis ( $\beta$ -aminoethylether)-N,N,N',N'-tetraacetic acid; FERM: 4.1 protein/ezrin/radixin/moesin protein ; IBMX: 3-isobutyl-1-methylxanthin; NDRG1: Neuregulin 1; PAGE: polyacrylamid gel electrophoresis; PBS: phosphate buffered saline; PMSF: phenylmethanesulfonylfluoride; TBST: Tris-buffered saline Tween 20; VS: vestibular schwannoma.

## Declarations

### Acknowledgements

We thank Siglinde Kühnel, Elisabeth Karl and Katharina Grüning (Department of Neurosurgery, University Hospital Würzburg, Germany) for excellent technical assistance. We are grateful to Ellaine Salvador (Department of Neurosurgery, University Hospital Würzburg, Germany) for language editing.

### Author's contributions

All authors contributed to the study conception and design. The study was supervised by MB, ML and CH. Samples were collected by MB and CM, experiments performed by AN and data analyzed by AN, MB and CH. CMM analyzed the histological tumor tissue samples. AN and CH wrote the draft of the manuscript

which was substantially revised by MB, CM, R-IE and ML. All authors read and approved the final manuscript. AN and MB contributed equally to this paper. MB is the corresponding author.

## **Funding**

This work was funded by the Interdisciplinary Center of Clinical Research (IZKF) Würzburg (Z-2/72). This publication was supported by the Open Access Publication Fund of the University of Würzburg.

## **Availability of data and material**

The datasets generated and analyzed during the current study are included in this published article and its supplementary information files. Raw data are available from the corresponding author on reasonable request.

## **Ethics approval and consent to participate**

The patients were treated at the Department of Neurosurgery, University Hospital Würzburg. They gave their written informed consent for participating in the study, which was conducted in accordance with the declaration of Helsinki and as approved by the institutional review board of the University of Würzburg in July 2016 (#145/16).

## **Consent for publication**

Not applicable.

## **Competing interests**

The authors declare that they have no competing interests.

## **Author details**

<sup>#</sup>Anja Nattmann and Maria Breun contributed equally to this paper.

<sup>1</sup>Department of Neurosurgery, University Hospital Würzburg, Josef-Schneider-Str. 11, D-97080 Würzburg, Germany. <sup>2</sup>Department of Neuropathology, Institute of Pathology, University of Würzburg, D-97080 Würzburg, Germany.


## **References**

1. Tatagiba M, Ebner FH. Kleinhirnbrückenwinkelprozesse. In: Moskopp D, Wassmann H, editors. Neurochirurgie. 2nd ed. Stuttgart: Schattauer GmbH; 2015. pp. 596–608.
2. Cooper J, Giancotti FG. Molecular insights into NF2/Merlin tumor suppressor function. FEBS Lett. 2014;588:2743–52. doi:10.1016/j.febslet.2014.04.001.

3. Breun M, Schwerdtfeger A, Martellotta DD, Kessler AF, Monoranu CM, Matthies C, et al. ADAM9: A novel player in vestibular schwannoma pathogenesis. *Oncol Lett.* 2020;19:1856–64. doi:10.3892/ol.2020.11299.
4. Klein T, Bischoff R. Active metalloproteases of the A Disintegrin and Metalloprotease (ADAM) family: biological function and structure. *J Proteome Res.* 2011;10:17–33. doi:10.1021/pr100556z.
5. Breun M, Martellotta DD, Leberle A, Nietzer S, Baur F, Ernestus R-I, et al. 3D in vitro test system for vestibular schwannoma. *J Neurosci Methods.* 2020;336:108633. doi:10.1016/j.jneumeth.2020.108633.
6. Breun M, Schwerdtfeger A, Martellotta DD, Kessler AF, Perez JM, Monoranu CM, et al. CXCR4: A new player in vestibular schwannoma pathogenesis. *Oncotarget.* 2018;9:9940–50. doi:10.18632/oncotarget.24119.
7. Feldheim J, Kessler AF, Schmitt D, Salvador E, Monoranu CM, Feldheim JJ, et al. Ribosomal Protein S27/Metallopanstimulin-1 (RPS27) in Glioma-A New Disease Biomarker? *Cancers (Basel)* 2020. doi:10.3390/cancers12051085.
8. Hanemann CO. Magic but treatable? Tumours due to loss of Merlin. *Brain.* 2008;131:606–15. doi:10.1093/brain/awm249.
9. Nath D, Slocombe PM, Webster A, Stephens PE, Docherty AJ, Murphy G. Meltrin  $\gamma$  (ADAM-9) mediates cellular adhesion through  $\alpha 6\beta 1$  integrin, leading to a marked induction of fibroblast cell motility. *J Cell Sci.* 2000;113:2319–28.
10. Sughrue MIE, Yeung AH, Rutkowski MJ, Cheung SW, Parsa AT. Molecular biology of familial and sporadic vestibular schwannomas: implications for novel therapeutics. *J Neurosurg.* 2011;114:359–66. doi:10.3171/2009.10.JNS091135.
11. Ammoun S, Flaiz C, Ristic N, Schuldt J, Hanemann CO. Dissecting and targeting the growth factor-dependent and growth factor-independent extracellular signal-regulated kinase pathway in human schwannoma. *Cancer Res.* 2008;68:5236–45. doi:10.1158/0008-5472.CAN-07-5849.
12. Carroll SL. Molecular mechanisms promoting the pathogenesis of Schwann cell neoplasms. *Acta Neuropathol.* 2012;123:321–48. doi:10.1007/s00401-011-0928-6.
13. Schulze KMM, Hanemann CO, Müller HW, Hanenberg H. Transduction of wild-type merlin into human schwannoma cells decreases schwannoma cell growth and induces apoptosis. *Hum Mol Genet.* 2002;11:69–76. doi:10.1093/hmg/11.1.69.
14. Huang C-F, Yang S-F, Chiou H-L, Hsu W-H, Hsu J-C, Liu C-J, et al. Licochalcone A inhibits the invasive potential of human glioma cells by targeting the MEK/ERK and ADAM9 signaling pathways. *Food Funct.* 2018;9:6196–204. doi:10.1039/c8fo01643g.
15. Duffy MJ, McKiernan E, O'Donovan N, McGowan PM. Role of ADAMs in cancer formation and progression. *Clin Cancer Res.* 2009;15:1140–4. doi:10.1158/1078-0432.CCR-08-1585.
16. Lüllmann-Rauch R. Binde- und Stützgewebe. In: Lüllmann-Rauch R, editor. *Taschenlehrbuch Histologie.* 4th ed. Stuttgart: Thieme; 2012. pp. 119–73.

17. Hogervorst F, Admiraal LG, Niessen C, Kuikman I, Janssen H, Daams H, et al. Biochemical characterization and tissue distribution of the A and B variants of the integrin  $\alpha 6$  subunit. *J Cell Biol.* 1993;121:179–91. doi:10.1083/jcb.121.1.179.
18. Hanemann CO, Bartelt-Kirbach B, Diebold R, Kämpchen K, Langmesser S, Utermark T. Differential gene expression between human schwannoma and control Schwann cells. *Neuropathol Appl Neurobiol.* 2006;32:605–14. doi:10.1111/j.1365-2990.2006.00769.x.
19. Guaiquil V, Swendeman S, Yoshida T, Chavala S, Campochiaro PA, Blobel CP. ADAM9 is involved in pathological retinal neovascularization. *Mol Cell Biol.* 2009;29:2694–703. doi:10.1128/MCB.01460-08.
20. Brieger J, Bedavanija A, Lehr H-A, Maurer J, Mann WJ. Expression of angiogenic growth factors in acoustic neurinoma. *Acta Otolaryngol.* 2003;123:1040–5. doi:10.1080/00016480310005101.
21. Mazzocca A, Coppari R, Franco R de, Cho J-Y, Libermann TA, Pinzani M, et al. A Secreted Form of ADAM9 Promotes Carcinoma Invasion through Tumor-Stromal Interactions. *Cancer Res.* 2005;65:4728–38. doi:10.1158/0008-5472.CAN-04-4449.
22. Fry JL, Toker A. Secreted and membrane-bound isoforms of protease ADAM9 have opposing effects on breast cancer cell migration. *Cancer Res.* 2010;70:8187–98. doi:10.1158/0008-5472.CAN-09-4231.


## Figures



**Figure 1**

ADAM9 was not regulated by Merlin, but involved in controlling VS primary cell proliferation. a Western blot of ADAM9 expression in four representative VS. b Merlin protein expression in VS primary cells after lentiviral transfection. The immunofluorescent images compare successfully transfected green fluorescent cells (left) with the same field of view showing all cells (right) to estimate the transfection rate. Western blot of MCF7 breast cancer cells (MCF7), VS primary cells transfected with Merlin (Merlin) and untransfected VS primary cells (control). Shown is one representative experiment of  $n = 3$ . c ADAM9 protein expression in the same cells as shown in (b). d Lentiviral mediated shRNA knock down of ADAM9. Green fluorescent ADAM9 knock down (left) and scrambled transfected control cells (right). One representative experiment of  $n = 4$  is shown. Quantification of the cell numbers.  $\beta$ -Tubulin served as

loading control in all Western-blot experiments. Blots were cropped for better clarity. The full length blots are presented in Supplementary Figure S1.



**Figure 2**

Co-localization of ADAM9 with Integrin  $\alpha 6$  and Integrin  $\alpha 2\beta 1$  in VS tumor samples. a-f Immunofluorescence double-staining of ADAM9 (red) and Integrin  $\alpha 6$  (green). a-b Co-localization (yellow) was mainly found in connective tissue and c near blood vessels. d One VS did not express any detectable

Integrin  $\alpha 6$ . e and f demonstrate enriched co-localization along the tumor capsule. g-h Immunofluorescence double-staining of ADAM9 (red) and Integrin  $\alpha 2\beta 1$  (green). g Integrin  $\alpha 2\beta 1$  VS with ADAM9 co-localization. h Integrin  $\alpha 2\beta 1$  negative VS. DAPI = blue. Representative images of n = 20.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SUPPLEMENTARYINFORMATION.docx](#)
- [SupplementaryFigure1.docx](#)