

TCONS_00230836 silencing restores stearic acid-induced β -cell dysfunction through alleviating endoplasmic reticulum stress via a PERK/eIF2 α -dependent pathway rather than apoptosis

Rui Guo

Harbin Medical University

Yunjin Zhang

Harbin Medical University

Yue Yu

Harbin Medical University

Shenghan Su

Harbin Medical University

Qingrui Zhao

Harbin Medical University

Xia Chu

Harbin Medical University

Huimin Lu (✉ lhm_519@sina.com)

Harbin Medical University <https://orcid.org/0000-0002-4972-0899>

Changhao Sun

Harbin Medical University

Research

Keywords: Stearic acid, Pancreatic β cell, LncRNA, Endoplasmic reticulum stress

Posted Date: September 9th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-71491/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Chronic exposure of pancreatic β cells to high levels of stearic acid (C18:0) leads to impaired insulin secretion, which accelerates the progression of type 2 diabetes mellitus (T2DM). Recently, long noncoding RNAs (lncRNAs) were found to participate in saturated fatty acid-induced metabolism dysfunction. However, their contribution to stearic acid-induced β -cell dysfunction remains largely unknown. This study evaluated the possible role of the lncRNA TCONS_00230836 in stearic acid-stimulated lipotoxicity to β cells.

Method: Using high-throughput RNA-sequencing, TCONS_00230836 was screened out as being exclusively differentially expressed in stearic acid-treated mouse β -TC6 cells. Co-expression network was constructed to reveal the potential mRNAs targeted for lncRNA TCONS_00230836. Changes in this lncRNA's and candidate mRNAs' levels were further assessed by real-time PCR in stearic acid-treated β -TC6 cells and islets of mice fed a high-stearic-acid diet (HSD). The localization of TCONS_00230836 was detected by fluorescent in situ hybridization. The endogenous lncRNA TCONS_00230836 in β -TC6 cells was abrogated by its Smart Silencer.

Results: The lncRNA TCONS_00230836 was enriched in mouse islets and mainly localized in the cytoplasm. Its expression was significantly increased in stearic acid-treated β -TC6 cells and HSD-fed mouse islets. Knockdown of TCONS_00230836 apparently restored stearic acid-impaired GSIS through alleviating endoplasmic reticulum stress via a PERK/eIF2 α -dependent pathway. However, stearic acid-induced β -cell apoptosis was not obviously recovered.

Conclusion: Our findings suggest the involvement of the lncRNA TCONS_00230836 in stearic acid-induced β -cell dysfunction, which provides novel insight into stearic acid-induced lipotoxicity to β cells. Anti-lncRNA TCONS_00230836 might be a new therapeutic strategy for alleviating stearic acid-induced β -cell dysfunction in the progression of T2DM.

Introduction

Chronic exposure to elevated saturated fatty acids (SFAs) of pancreatic β cells is a key trigger of impaired insulin secretion, which is one of the most important characteristics of type 2 diabetes mellitus (T2DM). Evidence accumulated to date has mainly focused on palmitic acid, with little attention being paid to stearic acid-induced worsening of β -cell function. In our previous studies and other research, it has been demonstrated that, whether in the postprandial serum of T2DM patients or in the fasting serum of high-fat-diet-fed mice [1–4], only stearic acid levels were profoundly increased. Meanwhile, a long-term high level of stearic acid exhibited a stronger destructive effect on β cells than that of palmitic acid [2, 5]. Although endoplasmic reticulum (ER) stress [2, 6] and apoptosis [5, 7] are accepted as major contributors to stearic acid-induced β -cell dysfunction, the molecular mechanisms involved in this remain largely unclear.

Long noncoding RNAs (lncRNAs) [8], generally defined as RNA molecules > 200 nucleotides in length without evident protein-coding capacity, are well known to be involved in diverse gene-regulatory mechanisms such as transcription, imprinting, splicing, protein degradation, and epigenetic marks on chromatin [9, 10], and their dysregulation has been implicated in many diseases, especially in several metabolic diseases [11, 12]. Recently, a number of lncRNAs have been reported to participate in SFA-induced metabolic dysfunction in liver [13], adipocytes [14, 15], skeletal muscle cells [16], and pancreatic β cells [17, 18]. However, there are no data regarding the possible contribution of lncRNAs to stearic acid-induced β -cell dysfunction.

Using RNA-sequencing and qRT-PCR, we showed for the first time that the lncRNA TCONS_00230836 was differentially expressed exclusively in stearic acid-treated β -TC6 cells, compared with both palmitic acid and control groups. We investigated the effect of the lncRNA TCONS_00230836 on stearic acid-induced β -cell dysfunction by loss-of-function approaches in a cellular model of lipotoxicity. Our experimental results suggested that this lncRNA is a detrimental factor to pancreatic β cells in the presence of stearic acid and that its knockdown could mitigate stearic acid-stimulated impairment of insulin secretion, which would provide a potential target for the prevention and treatment of T2DM.

Material And Methods

Chemicals.

Stock solutions of stearic acid and palmitic acid (Sigma, St. Louis, MO, USA) supplemented with BSA (3 mmol/L fatty acids:1.5 mmol/L BSA) were prepared by dissolving them in ethanol and saponifying them with sodium hydroxide, as described previously [19]. After the sodium salt had dried, it was resuspended in saline and then heated at 80 °C until completely dissolved. Next, 20% (w/v) BSA was added and the mixture was stirred at 50 °C for 4 h. The complex was then sterilized and aliquoted.

Cell culture.

The pancreatic β -TC6 cells and AML12 cells were obtained from Shanghai Academy of Chinese Sciences Cell Library and cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco/Life Technologies, Carlsbad, CA, USA), supplemented with 15% fetal bovine serum (FBS; Gibco/Life Technologies), 50 μ g/mL streptomycin, 50 IU/mL penicillin (Gibco/Life Technologies), and 1.5 g/L NaHCO₃. Mouse islets were isolated after collagenase P (Cat. 11213873001; Roche Molecular Biochemicals, Germany) digestion of the pancreas by ductal injection and filtered with hardware cloth (600 μ m). Next, islets were purified with different concentrations of Ficoll 400 (Cat. 17-0300-10; Pharmacia). Finally, islets were incubated in RPMI 1640 (Gibco) containing 10% FBS, 0.11 g/L sodium pyruvate, and 100 units/mL penicillin-streptomycin. The β -TC6 cells and islets were incubated with 400 μ mol/L stearic or palmitic acid for 24 h. AML 12 cells were also treated with 400 μ mol/L stearic acid for 24 h.

RNA sequencing and data analysis.

The lncRNA and mRNA high-throughput sequencing was performed by Genewiz (Suzhou, China), processing nine samples from pancreatic β -TC6 cells (3 control, 3 stearic acid, and 3 palmitic acid samples). Differential expression analysis was performed using the DESeq Bioconductor package in R by first transforming the raw count data to \log_2 counts per million reads using the “voom” function ($P < 0.05$).

Construction of gene co-expression network.

The Pearson correlation coefficients (PCC) were performed for lncRNA TCONS_00230836-mRNA pairs, of which the $PCC \geq 0.950$, or $PCC \leq -0.950$, $P < 0.05$ were selected to construct the gene co-expression network using Cytoscape software. Each gene corresponds to a node in the network.

Animal experiments.

Twenty male C57BL/6J mice (6 weeks old) were purchased from Beijing Vital River Laboratory Animal Technology Company (Beijing, China). After adaptive feeding for 1 week, these mice were randomly divided into control and high-stearic-acid diet groups ($n = 10$ per group), according to their body weights. The control diet (1025) and high-stearic-acid diet (HSD) (H10060) were obtained from Beijing HFK Bioscience Co. Ltd. (Beijing, China) (Additional file 1). After 20 weeks of feeding the mice, islets and blood samples were collected for biochemical analysis. Fasting (12 hours) serum glucose, total cholesterol, triacylglycerol, high-density-lipoprotein cholesterol, and low-density-lipoprotein cholesterol levels were calculated using an automatic analyzer (Hitachi-7100; Hitachi, Tokyo, Japan), kits for which were purchased from Biosino Biotechnology Co. (Beijing, China). Serum insulin level was measured using a mouse/rat insulin ELISA kit (Linco Research, St. Charles, MO, USA).

Serum nonesterified fatty acid profile measurement.

Fasting serum nonesterified fatty acids were transformed to fatty acid methyl esters, as described in our previous studies [4, 20]. Gas chromatography–mass spectrometry analysis was performed using a TRACE gas chromatograph with a Polaris Q mass spectrometer (Thermo Finnigan, San Jose, CA, USA). Separation was obtained on a J&W DB-WAX capillary column (30-m, 0.25-mm I.D., 0.25- μ m film thickness; Agilent J&W Scientific, Folsom, CA, USA). Heptadecanoic acid (C17:0) was used as an internal standard.

Cell viability measurements.

Cell viability was measured by the assessment of lactate dehydrogenase (LDH) release and CCK 8 assay. For measuring LDH, the culture medium was collected and characterized using LDH assay kit (Thermo Fisher Scientific Inc.). For the CCK 8 assay, we used the Cell Counting Kit 8 (CCK-8, C0038; Beyotime Biotechnology, Shanghai, China). The β -TC6 cells were seeded into each well of a 96-well plate and 100 μ L of CCK-8 mixed reagents were added to each well. After 1 h of incubation at 37 °C, absorbance was determined at 450 nm using a microplate reader (SpectraMax M2; Molecular Devices, San Jose, CA, USA).

Apoptosis assay.

The β -TC6 cells were collected and stained with fluorescein isothiocyanate (FITC)-annexin V and propidium iodide (PI), in accordance with the manufacturer's instructions (Lot#AB116F; Absin, Shanghai, China). The apoptosis rate was analyzed by flow cytometry (LSR Fortessa; BD Biosciences, San Jose, CA, USA).

Fluorescent in situ hybridization (FISH).

The β -TC6 cells were seeded into a 24-well plate with sterile slides overnight. Cells were rinsed using 1 \times PBS for 5 min and fixed in 4% formaldehyde at room temperature for 10 min. The β -TC6 cells were washed three times with 1 \times PBS for 5 min each. Next, the cells were permeabilized in 1 \times PBS containing 0.5% Triton X-100 for 5 min at 4 $^{\circ}$ C. After washing the cells with 1 \times PBS, they were blocked in Blocking Solution and Pre-hybridization (1:99) mixed solution at 37 $^{\circ}$ C for half an hour. Then, in the dark, discarding the mixed solution, the β -TC6 cells were incubated with a hybridized mixture containing lncRNA Probe Mix or U6/18S at 37 $^{\circ}$ C overnight. On the second day, the β -TC6 cells were washed at 42 $^{\circ}$ C with different concentrations of SSC solution, which were also protected from the light. After immersing the β -TC6 cells in 1 \times PBS for 5 min, they were stained with DAPI staining solution at room temperature for 10 min and then washed with 1 \times PBS. The slides taken from the 24-well plate were observed with a laser confocal microscope. The whole process was protected from light to prevent quenching. The locked nucleic acid-modified oligonucleotide probe targeting lncRNA TCONS_00230836 and the FISH Kit were purchased from Ribobio Co. Ltd. (Guangzhou, China).

Transfection procedures.

Smart Silencer for TCONS_00230836 and its negative control (Ribobio Co. Ltd.) were transfected into β -TC6 cells using C10511-05 riboFectTM CP Transfection Kit (Ribobio Co. Ltd.) for 24 h, in accordance with the manufacturer's instructions. The sequences of lncRNA Smart Silencer for mouse TCONS_00230836 are displayed in Table 1, including three siRNA and three antisense oligonucleotides.

Table 1
The sequences of lncRNA Smart Silencer for mouse TCONS_00230836

Gene	sequences
siRNA-1 target sequence	CCTTAATCCCAACCCTCAA
siRNA-2 target sequence	TAGACATAGCCACATGAAA
siRNA-3 target sequence	CCCAATAGTTAATGACAGA
antisense oligonucleotides target sequence-1	GCTGTCAAAAAGGAATCACA
antisense oligonucleotides target sequence-2	GGAAATGCAGTGTAGTAGAA
antisense oligonucleotides target sequence-3	ATGGGCCACCCCTCTAAGAT

Quantitative real-time polymerase chain reaction (qRT-PCR).

Total RNA was extracted from β -TC6 cells and mouse islets using TRIzol reagent (Invitrogen), in accordance with the manufacturer's protocol. The qRT-PCR was performed with SYBR Green PCR Master Mix (Applied Biosystems, Foster City, CA, USA), with β -actin as an internal control. All primers used in this study were synthesized by Sangon Biotech Co. Ltd. (Shanghai, China), the sequences of which are listed in Table 2.

Table 2
Primer sequences used for qRT-PCR

Gene		sequences (5' to 3')
TCONS_00230836	Forward	AACCACAGGCTTCGGGATAGTC
	Reverse	CAAGCATAACAGGCGGGAAGTC
ENSMUST00000074898	Forward	ACCTTAAACGACGAGAAGCAATGG
	Reverse	AGCCAGACACGTAGCCCACACG
ENSMUST00000086399	Forward	TGGTCACCGTTGTGATCCC
	Reverse	TGAGGTCCTCCTTGCATG
ENSMUST00000050785	Forward	CTGTCTGTCTGCCACTCCAT
	Reverse	GCTGGCCAAATAAGAAGAGG
ENSMUST00000124100	Forward	ACAAACCGCAGAGAAACAAA
	Reverse	CCTCCCTCTGCCCTAGTATG
ENSMUST00000092822	Forward	GCAGACCCAGTCGTTGTAC
	Reverse	AAGCCTGTGGCACAACATC
ENSMUST00000021620	Forward	GGCTGAAGACAGTTGTGGAA
	Reverse	GGGTAAATCTTGCCCTTTCA
ENSMUST00000073388	Forward	GAGTCACTTGTCCGCAGCTGTC
	Reverse	TCGCTGTCAGTTAAGTCCAGG
ENSMUST00000159720	Forward	CGCTATTGTCTTCTTGATGGAC
	Reverse	CTTCAACAGTTTCCCTGAGTTG
ENSMUST00000131456	Forward	TGATGAGTAGCGTAAAGTACCC
	Reverse	ATATGAGGCATCGTCAGGTAAG
ENSMUST00000185596	Forward	ATCTCTTTGCCTTCCTCAATCA
	Reverse	GTTTTGATCAGCTCATTACAGT
β-actin	Forward	GGTCAGAAGGACTCCTATGTGG
	Reverse	TGTCGTCCCAGTTGGTAACA

Western blotting.

Cells were gently washed three times in 1 × PBS and then 50 µL of intermediate RIPA Lysis Buffer (Beyotime) was added per 24-well plate. Protein concentrations were detected using bicinchoninic acid (BCA) protein assay kits (Cat. No. P0010; Beyotime). Protein samples (approximately 80 µg) were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred to polyvinylidene fluoride membranes. The primary antibodies used were as follows: Phospho-PERK (3179S, CST, dilution of 1:1000), PERK (3192, CST, dilution of 1:1000), Phospho-eIF2α (9721, CST, dilution of 1:1000), eIF2α (9722, CST, dilution of 1:1000), Caspase-3 (9662, CST, dilution of 1:1000), Parp-1 (9532, CST, dilution of 1:1000), Bcl-2 (4223, CST, dilution of 1:1000), ATF-6 (65880, CST, dilution of 1:1000), IRE1α (3294, CST, dilution of 1:1000), and β-actin (sc-130657, Santa Cruz, dilution of 1:800). The secondary antibody was anti-rabbit (s3738, Promega, dilution of 1:7500) alkaline phosphatase-conjugated antibody. Protein coloration was performed using the Stabilized Substrate for Alkaline Phosphatase (S3841, Promega) and was screened by the FluorChem R system (ProteinSimple, San Jose, CA, USA).

Glucose-stimulated insulin secretion assay.

To assess glucose-stimulated insulin secretion (GSIS), β-TC6 cells were incubated in secretion buffer [NaCl 129, KCl 4.8, MgSO₄ 1.2, KH₂PO₄ 1.2, CaCl₂ 2.5, NaHCO₃ 5.0, and HEPES 10 (all mmol/L) supplemented with 1 mg/mL bovine serum albumin, adjusted to pH 7.4] for an additional 60 min with 2.8 or 20 mmol/L glucose [2]. After collecting the supernatant for insulin measurement, the β-TC6 cells were lysed in intermediate RIPA Lysis Buffer (Beyotime) for later evaluation of the total protein content with BCA protein assay reagent kit (Cat. No. P0010, Beyotime) following the manual. Insulin levels in cell culture medium were measured using a mouse/rat insulin ELISA kit (Linco Research) and standardized by every milligram protein per hour in each well.

Statistics.

All statistical data were analyzed with SPSS software, version 21.0 (SPSS Inc., Chicago, IL, USA), and are reported as mean ± SEM. Two-tailed Student's *t*-test was used to analyze differences between two groups, and one-way ANOVA followed by Student–Newman–Keuls test was used to test differences among three or more groups. *P*-values < 0.05 were considered statistically significant.

Results

Animal model characteristics.

We successfully established a mouse model to mimic the increased proportions of stearic acid *in vivo*, as shown by a significant increase in circulating stearic acid in HSD-fed mice (Additional file 2). The metabolic characteristics of these animals are displayed in Additional file 3.

Upregulation of lncRNA TCONS_00230836 expression in stearic acid-treated β-TC6 cells and islets of mice fed a high-stearic-acid diet.

The results obtained by high-throughput RNA-sequencing demonstrated that the lncRNA TCONS_00230836, an intergenic lncRNA located at 65803111 to 65809116 of chromosome 10, was markedly elevated with a 1.328 log₂ fold change (*P* value: 5.062E-07) in β-TC6 cells exposed to stearic acid and a 0.270 log₂ fold change (*P* value: 0.327) in the palmitic group, compared with the level in the normal group (Fig. 1a). A similar increase in lncRNA TCONS_00230836 level was also revealed in both stearic acid-treated β-TC6 cells and islets of mice fed a high-stearic-acid diet by qRT-PCR. The expression level of the lncRNA TCONS_00230836 was increased by 3.744-fold in stearic acid-treated β-TC6 cells (Fig. 1b) and 5.115-fold in the islets of mice from the high-stearic-acid group (Fig. 1c), compared with that in the control group.

Differential expression of lncRNA TCONS_00230836 in various tissues and cell lines of mouse.

qRT-PCR results verified that the level of the lncRNA TCONS_00230836 was significantly differentially expressed in various tissues as well as cell lines those are closely related to high-fat-diet induced metabolic disorders. Among them, the expression of this lncRNA was highest in the islets and lowest in the skeletal muscle of HSD-fed mice (Fig. 2a). However, no matching sequence was detected in peritesticular and perirenal adipose tissues (data not shown). Meanwhile, the expression of the lncRNA TCONS_00230836 was apparently higher in β-TC6 cells than that in AML12 cells (Fig. 2b). In addition to islets and β-TC6 cells, after the treatment with a high concentration of stearic acid, the lncRNA TCONS_00230836 level was also increased in brown adipose, whereas it was decreased in liver and skeletal muscle of HSD-fed mice and AML12 cells (Additional file 4).

lncRNA TCONS_00230836 is mainly localized in cytoplasm.

The confocal images of FISH showed that the lncRNA TCONS_00230836 is mainly distributed in the cytoplasm of β-TC6 cells (Fig. 3).

Knockdown of lncRNA TCONS_00230836 reversed stearic acid-induced impaired insulin secretion in islets and β-TC6 cells.

Both in islets and β-TC6 cells (Fig. 4a and 4b), transfection of the lncRNA TCONS_00230836 Smart Silencer for 24 h efficiently reduced the stearic acid-increased intracellular lncRNA TCONS_00230836 levels. Meanwhile, similar results were consistently observed in β-TC6 cells as indicated by FISH (Fig. 4c). Then, as shown in Fig. 4d and 4e, stearic acid markedly decreased GSIS in islets and β-TC6 cells, separately. After transfection of the lncRNA TCONS_00230836 Smart Silencer into pancreatic islets and β-TC6 cells, stearic acid-induced GSIS impairment was significantly alleviated. However, no significant alteration was observed in GSIS when the lncRNA TCONS_00230836 Smart Silencer was transfected alone in the absence of stearic acid.

Inhibition of lncRNA TCONS_00230836 alleviated stearic acid-induced ER stress rather than apoptosis in β-TC6 cells and mice islets.

Stearic acid apparently suppressed the cell survival rate and induced cell death in β -TC6 cells. Silencing the lncRNA TCONS_00230836 did not significantly block the stearic acid-stimulated cell viability reduction (Fig. 5a) and cell death (Fig. 5b). Meanwhile, the annexin V-FITC apoptosis assay results showed that lncRNA TCONS_00230836 inhibition did not reduce the stearic acid-increased percentage of apoptotic cells (Fig. 5c). Besides, the elevated protein levels of cleaved-Parp1 and cleaved-Caspase3, and the decreased levels of B-cell CLL/lymphoma 2 in the stearic acid group were not rescued by the lncRNA TCONS_00230836 Smart Silencer (Fig. 5d). However, the expression of the ER stress-related proteins Phospho-PERK and Phospho-eIF2 α in stearic acid-treated β -TC6 cells was significantly decreased after inhibition of the lncRNA TCONS_00230836 (Fig. 6a), whereas stearic acid-increased protein levels of ATF-6 and IRE1 α were not suppressed after knockdown of TCONS_00230836 (Fig. 6b). Moreover, transfection of the lncRNA TCONS_00230836 Smart Silencer alone did not alter apoptosis and ER stress in β -TC6 cells. In addition, similar results were consistently observed in mice islets (Fig. 7a-c, Fig. 8a, b).

Effect of lncRNA TCONS_00230836 on the expressions of candidate ER stress-related mRNAs.

In order to reveal the potential mRNAs targeted for lncRNA TCONS_00230836 in stearic acid-treated β -cell ER stress, the lncRNA TCONS_00230836-mRNAs co-expression network was constructed. 12 up-regulated mRNAs are positively associated with lncRNA TCONS_00230836 (Fig. 9a), and 15 down-regulated mRNAs are negatively related to this lncRNA (Fig. 9b). Among these mRNAs, five stearic acid-increased mRNAs levels (positively related) (ENSMUST00000074898 [*Hp*], ENSMUST00000159720 [*Alpk1*], ENSMUST00000086399 [*Icam1*], ENSMUST00000131456 [*Serp1*], ENSMUST00000050785 [*Lcn2*]) were effectively reversed by lncRNA TCONS_00230836 Smart Silencer. However, after transfection of lncRNA TCONS_00230836 Smart Silencer alone without stearic acid treatment, only *Alpk1*, *Icam1* and *Serp1* were significantly reduced (Fig. 9c). As for negatively related mRNAs, the expressions of five stearic acid-decreased mRNAs (ENSMUST00000124100 [*Prn*], ENSMUST00000092822 [*Bcas3*], ENSMUST00000021620 [*Otub2*], ENSMUST00000073388 [*Afmid*], ENSMUST00000185596 [*Srgap2*]) were apparently rescued after knocking down lncRNA TCONS_00230836. Meanwhile, these five mRNAs were significantly up-regulated after silencing lncRNA TCONS_00230836 in the absence of stearic acid (Fig. 9d). The log₂ fold change of these top five up- and down-regulated differentially expressed mRNAs were listed in Additional file 5.

Discussion

The prevalence of T2DM has been increasing worldwide, with an estimated 336 million people currently affected [21]. It is well established that exposing pancreatic β cells to an elevated level of stearic acid causes their dysfunction, which is far advanced by the time diabetes is diagnosed clinically [1–3, 5]. Therefore, the optimal management and prevention of T2DM should aim to ameliorate β -cell dysfunction at an early stage. However, the effective preventive targets are limited and still need to be comprehensively explored. The present study elucidated the pathophysiological role of lncRNA in stearic acid-treated β cells. Our findings demonstrated that the lncRNA TCONS_00230836 was significantly increased in both stearic acid-induced β -TC6 cells and the islets of mice fed a high-stearic-acid diet.

Inhibition of the lncRNA TCONS_00230836 effectively protected against stearic acid-induced β -cell dysfunction. Interestingly, after knocking down the lncRNA TCONS_00230836, we found that stearic acid-stimulated β -cell ER stress was apparently ameliorated via a PERK/eIF2 α -dependent pathway, but no obvious improvement of stearic acid-induced apoptosis was observed.

Although a number of studies have preliminarily explored the mechanisms underlying stearic acid-induced pancreatic β -cell damage, the exact molecular targets and associated pathways still need to be established. Recently, accumulating evidence has suggested that noncoding RNAs play an important role in the β -cell injury caused by the long-term consumption of a high-fat diet, such as microRNAs [2, 5], circRNAs [22, 23], and lncRNAs [17, 24, 25]. Nevertheless, few studies have focused on the role of lncRNAs in β -cell dysfunction upon exposure to elevated levels of SFAs, especially stearic acid, on which no related research has previously been published. In our study, we used high-throughput sequencing and bioinformatic technology to select five lncRNAs that were differentially expressed specifically in stearic acid-treated β -TC6 cells, compared with their levels in both control and palmitic acid groups (data not shown). Among the known lncRNAs, the lncRNA TCONS_00230836 displayed the largest fold increase, with an increase of 1.328 log₂ fold change in RNA sequencing results, 3.744-fold in stearic acid-induced β -TC6 cells, and 5.115-fold in islets of mice fed a high-stearic-acid diet, as revealed by qRT-PCR.

It has been well clarified that lncRNAs have low sequence conservation and high tissue specificity [26]. Meanwhile, the expression of lncRNAs typically varies more between tissues than for other noncoding RNAs [27–29]. To investigate the difference of sequence conservation and differential expression of the lncRNA TCONS_00230836, we tested it in different tissues (islet, liver, skeletal muscle, brown adipose, peritesticular, and perirenal adipose tissues) and cell lines (β -TC6 cells and AML12 cells) which are closely related to T2DM. Our data indicated that there was a highly similar sequence of lncRNA TCONS_00230836 (similarity > 98%) in liver, brown adipose, skeletal muscle, as well as the AML12 cell line, but not in peritesticular and perirenal adipose tissues, compared with that in islets and β -TC6 cells. Moreover, the level of this lncRNA was highest in islets and lowest in skeletal muscle of high-stearic-acid-fed mice, which suggested that this lncRNA may be specific to and enriched in islets. This is the first key finding of our study.

To demonstrate the role of the lncRNA TCONS_00230836 in stearic acid-mediated lipotoxicity in β cells, endogenous TCONS_00230836 in β -TC6 cells was abrogated by the application of its Smart Silencer, a highly effective inhibitor that specifically targets lncRNA TCONS_00230836 expressed in both the cytoplasm and the nucleus. We found that stearic acid-impaired GSIS was largely restored after silencing TCONS_00230836, which implied that a TCONS_00230836-mediated mechanism operated in the stearic acid-induced lipotoxicity of β cells. However, how does the lncRNA TCONS_00230836 exert its influence and which process was improved in stearic acid-induced pancreatic β -cell injury?

ER stress and apoptosis are very important pathophysiological perturbations and are closely associated with β -cell lipotoxicity in the early stage of T2DM; they are not only independent of each other, but also have a close cause–effect relationship [30–34]. To reveal which process was affected by the lncRNA

TCONS_00230836, we detected the alterations of ER stress and apoptosis induced by stearic acid after transfection of the Smart Silencer for the lncRNA TCONS_00230836. Our data indicated that transfection of the lncRNA TCONS_00230836 effectively ameliorated stearic acid-stimulated ER stress via a PERK/eIF2 α -dependent pathway. As we know, ER stress can transduce apoptotic signals that may eventually result in apoptosis. However, unexpectedly, stearic acid-induced β -cell apoptosis was not significantly recovered after knocking down lncRNA TCONS_00230836 expression. Therefore, we speculate that there may be some other key regulatory factors involved in lncRNA TCONS_00230836-mediated β -cell ER stress. This is another important finding of our study.

In an effort to reveal the potential target for lncRNA TCONS_00230836 in stearic acid-induced ER stress, we analyzed the mRNAs profile by high-throughput screens in our previous study [35] and constructed the co-expression network between lncRNA TCONS_00230836 and targeted mRNAs which are closely related to ER stress. Then top five up-regulated mRNAs and another top five down-regulated mRNAs were selected for further identification. In the positive-relation group, qPCR results for the mRNA expressions indicated that *Alpk1*, *Icam1* and *Serping1* are probably involved in the process of stearic acid-induced β -cell ER stress associated with lncRNA TCONS_00230836. While in negative-relation group, *Prn*, *Bcas3*, *Otub2*, *Afmid* and *Srgap2* might play a critical role in stearic acid- β -cell ER stress closely related to lncRNA TCONS_00230836. Among them, *Afmid* has been reported that they could lead to impaired glucose tolerance in type 2 diabetes [36]. Meanwhile, *Otub2* has been shown to act through the inhibition of NF- κ B signaling in type 1 diabetes [37, 38]. Further experiments should be performed to explore the precise regulatory relationship between these candidate mRNAs and lncRNA TCONS_00230836 as well as PERK/eIF2 α -dependent signaling in stearic acid-treated β cells.

Our understanding of lncRNAs in stearic acid-induced lipotoxicity to β cells is still in its infancy and several questions remain unanswered. First, we did not assess the role of the lncRNA TCONS_00230836 in stearic acid-induced impairment of insulin secretion in an *in vivo* experiment. Second, we all know that lncRNAs exert their function through diverse regulatory mechanisms, for example, modulating translation, promoting mRNA degradation, or acting as miRNA sponges. Therefore, further experiments still need to be performed to improve our understanding of the regulatory mechanisms of the lncRNA TCONS_00230836 in stearic acid-induced lipotoxicity to pancreatic β cells.

Conclusions

We provide evidence that stearic acid can stimulate upregulation of the lncRNA TCONS_00230836 in stearic acid-treated pancreatic β cells. It appears that this lncRNA is specific to and enriched in pancreatic β cells. Suppression of this lncRNA could effectively restore stearic acid-induced impairment of insulin secretion through the amelioration of ER stress rather than apoptosis via PERK/eIF2 α -dependent signaling. These findings provide novel insight into stearic acid-induced lipotoxicity and the progression of T2DM, whereby lowering the level of the lncRNA TCONS_00230836 might be an effective strategy for improving lipotoxicity in β cells.

Declarations

Ethics approval and consent to participate

All animal procedures were ethically approved by Institutional Animal Care and Use Committee of Harbin Medical University, and animals were maintained in line with the guidelines of the Animal Experimental Center of Harbin Medical University.

Consent for publication

Not applicable.

Availability of data and material

All of the data are available with reasonable request from the corresponding authors.

Competing interests

The authors declare that there is no conflict of interest.

Funding

This work was supported by the Excellent Youth Foundation of Heilongjiang Province of China (YQ2020H033); University Nursing Program for Young Scholars with Creative Talents in Heilongjiang Province (UNPYSCT-201705), National Natural Science Foundation of China (No. 81773424).

Authors' contributions

H.L. and C.S. conceived and designed the experiments. R.G., Y.Z., Y.Y., S.S. and Q.Z. performed the experiments. R.G. analyzed data. H.L. and R.G. wrote the manuscript. H.L., R.G. and X.C. reviewed and edited the manuscript. We thank Liwen Bianji, Edanz Group China (www.liwenbianji.cn/ac), for editing the English text of a draft of this manuscript.

Acknowledgments

Not applicable.

References

1. Chu X, Liu L, Na L, et al. Sterol regulatory element-binding protein-1c mediates increase of postprandial stearic acid, a potential target for improving insulin resistance, in hyperlipidemia.

- Diabetes. 2013;62:561–71.
2. Lu H, Hao L, Li S, et al. Elevated circulating stearic acid leads to a major lipotoxic effect on mouse pancreatic beta cells in hyperlipidaemia via a miR-34a-5p-mediated PERK/p53-dependent pathway. *Diabetologia*. 2016;59:1247–57.
 3. Zhao L, Ni Y, Yu H, et al. Serum stearic acid/palmitic acid ratio as a potential predictor of diabetes remission after Roux-en-Y gastric bypass in obesity. *FASEB J*. 2017;31:1449–60.
 4. Liu L, Li Y, Guan C, et al. Free fatty acid metabolic profile and biomarkers of isolated post-challenge diabetes and type 2 diabetes mellitus based on GC-MS and multivariate statistical analysis. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2010;878:2817–25.
 5. Guo R, Yu Y, Zhang Y, et al. Overexpression of miR-297b-5p protects against stearic acid-induced pancreatic beta-cell apoptosis by targeting LATS2. *Am J Physiol Endocrinol Metab*. 2020;318:E430–9.
 6. Nemcova-Furstova V, Balusikova K, Sramek J, James RF, Kovar J. Caspase-2 and JNK activated by saturated fatty acids are not involved in apoptosis induction but modulate ER stress in human pancreatic beta-cells. *Cell Physiol Biochem*. 2013;31:277–89.
 7. Nemcova-Furstova V, Balusikova K, Halada P, Pavlikova N, Sramek J, Kovar J. Stearate-Induced Apoptosis in Human Pancreatic beta-Cells is Associated with Changes in Membrane Protein Expression and These Changes are Inhibited by Oleate. *Proteomics Clin Appl*. 2019;13:e1800104.
 8. Sarropoulos I, Marin R, Cardoso-Moreira M, Kaessmann H. Developmental dynamics of lncRNAs across mammalian organs and species. *Nature*. 2019;571:510–4.
 9. Ferre F, Colantoni A, Helmer-Citterich M. Revealing protein-lncRNA interaction. *Brief Bioinform*. 2016;17:106–16.
 10. Zhang X, Wang W, Zhu W, et al. Mechanisms and Functions of Long Non-Coding RNAs at Multiple Regulatory Levels. *Int J Mol Sci*. 2019;20.
 11. Jin F, Wang N, Zhu Y, et al. Downregulation of Long Noncoding RNA Gas5 Affects Cell Cycle and Insulin Secretion in Mouse Pancreatic beta Cells. *Cell Physiol Biochem*. 2017;43:2062–73.
 12. Bai X, Geng J, Li X, et al. Long Noncoding RNA LINC01619 Regulates MicroRNA-27a/Forkhead Box Protein O1 and Endoplasmic Reticulum Stress-Mediated Podocyte Injury in Diabetic Nephropathy. *Antioxid Redox Signal*. 2018;29:355–76.
 13. Zhang Q, Wang J, Li H, et al. LncRNA Gm12664-001 ameliorates nonalcoholic fatty liver through modulating miR-295-5p and CAV1 expression. *Nutr Metab (Lond)*. 2020;17:13.
 14. Muret K, Desert C, Lagoutte L, et al. Long noncoding RNAs in lipid metabolism: literature review and conservation analysis across species. *BMC Genom*. 2019;20:882.
 15. Degirmenci U, Li J, Lim YC, et al. Silencing an insulin-induced lncRNA, lncASIR, impairs the transcriptional response to insulin signalling in adipocytes. *Sci Rep*. 2019;9:5608.
 16. Han M, You L, Wu Y, et al. RNA-sequencing analysis reveals the potential contribution of lncRNAs in palmitic acid-induced insulin resistance of skeletal muscle cells. *Biosci Rep*. 2020;40.

17. Kong X, Liu CX, Wang GD, et al. LncRNA LEGLTBC Functions as a ceRNA to Antagonize the Effects of miR-34a on the Downregulation of SIRT1 in Glucolipotoxicity-Induced INS-1 Beta Cell Oxidative Stress and Apoptosis. *Oxid Med Cell Longev*. 2019;2019:4010764.
18. Arnes L, Akerman I, Balderes DA, Ferrer J, Sussel L. *betalinc1* encodes a long noncoding RNA that regulates islet beta-cell formation and function. *Genes Dev*. 2016;30:502–7.
19. Goldstein JL, Basu SK, Brown MS. Receptor-mediated endocytosis of low-density lipoprotein in cultured cells. *Methods Enzymol*. 1983;98:241–60.
20. Guo F, Huang C, Liao X, et al. Beneficial effects of mangiferin on hyperlipidemia in high-fat-fed hamsters. *Mol Nutr Food Res*. 2011;55:1809–18.
21. Ashcroft FM, Rorsman P. Diabetes mellitus and the beta cell: the last ten years. *Cell*. 2012;148:1160–71.
22. Stoll L, Sobel J, Rodriguez-Trejo A, et al. Circular RNAs as novel regulators of beta-cell functions in normal and disease conditions. *Mol Metab*. 2018;9:69–83.
23. Bai C, Yang W, Lu Y, Wei W, Li Z, Zhang L. Identification of Circular RNAs Regulating Islet beta-Cell Autophagy in Type 2 Diabetes Mellitus. *Biomed Res Int*. 2019;2019:4128315.
24. Ruan Y, Lin N, Ma Q, et al. Circulating LncRNAs Analysis in Patients with Type 2 Diabetes Reveals Novel Genes Influencing Glucose Metabolism and Islet beta-Cell Function. *Cell Physiol Biochem*. 2018;46:335–50.
25. Knoll M, Lodish HF, Sun L. Long non-coding RNAs as regulators of the endocrine system. *Nat Rev Endocrinol*. 2015;11:151–60.
26. Nagano T, Fraser P. No-nonsense functions for long noncoding RNAs. *Cell*. 2011;145:178–81.
27. Cabili MN, Trapnell C, Goff L, et al. Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev*. 2011;25:1915–27.
28. Derrien T, Johnson R, Bussotti G, et al. The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res*. 2012;22:1775–89.
29. Pauli A, Valen E, Lin MF, et al. Systematic identification of long noncoding RNAs expressed during zebrafish embryogenesis. *Genome Res*. 2012;22:577–91.
30. Vong CT, Tseng HHL, Kwan YW, Lee SM, Hoi MPM. Novel protective effect of O-1602 and abnormal cannabidiol, GPR55 agonists, on ER stress-induced apoptosis in pancreatic beta-cells. *Biomed Pharmacother*. 2019;111:1176–86.
31. Ljubkovic M, Gressette M, Bulat C, et al. Disturbed Fatty Acid Oxidation, Endoplasmic Reticulum Stress, and Apoptosis in Left Ventricle of Patients With Type 2 Diabetes. *Diabetes*. 2019;68:1924–33.
32. Kharroubi I, Ladriere L, Cardozo AK, Dogusan Z, Cnop M, Eizirik DL. Free fatty acids and cytokines induce pancreatic beta-cell apoptosis by different mechanisms: role of nuclear factor-kappaB and endoplasmic reticulum stress. *Endocrinology*. 2004;145:5087–96.
33. Wang J, Yang X, Zhang J. Bridges between mitochondrial oxidative stress, ER stress and mTOR signaling in pancreatic beta cells. *Cell Signal*. 2016;28:1099–104.

34. Tomita T. Apoptosis in pancreatic beta-islet cells in Type 2 diabetes. *Bosn J Basic Med Sci.* 2016;16:162–79.
35. Yu Y, Guo R, Zhang Y, et al. miRNA-mRNA profile and regulatory network in stearic acid-treated beta-cell dysfunction. *J Endocrinol.* 2020;246:13–27.
36. Hugill AJ, Stewart ME, Yon MA, et al. Loss of arylformamidase with reduced thymidine kinase expression leads to impaired glucose tolerance. *Biol Open.* 2015;4:1367–75.
37. Ma Y, Sun Y. miR-29a-3p inhibits growth, proliferation, and invasion of papillary thyroid carcinoma by suppressing NF-kappaB signaling via direct targeting of OTUB2. *Cancer Manag Res.* 2019;11:13–23.
38. Beck A, Vinik Y, Shatz-Azoulay H, et al. Otubain 2 is a novel promoter of beta cell survival as revealed by siRNA high-throughput screens of human pancreatic islets. *Diabetologia.* 2013;56:1317–26.

Figures




Figure 1

LncRNA TCONS_00230836 upregulation in stearic acid-treated mice β -TC6 cells and islets fed a high-stearic-acid diet. (a) The studied lncRNA TCONS_00230836 with fold changes, p values, and genomic locations in the presence of stearic acid by RNA sequencing. (n=5) (b) qRT-PCR results verified that the

level of the lncRNA TCONS_00230836 was elevated in stearic acid-treated β -TC6 cells. (n=4) (c) The expression of the lncRNA TCONS_00230836 was also increased in islets from mice fed a high-stearic-acid diet, as revealed by qRT-PCR. Ctrl, control group; SA, stearic acid; HSD, high-stearic-acid diet. **P < 0.01, versus the Ctrl group.




Figure 2

Identification of the lncRNA TCONS_00230836 in mice various tissues and multiple cell lines, by qRT-PCR. (a) Differential expression of the lncRNA TCONS_00230836 in islet, brown adipose, liver, and skeletal muscle of mice fed a chow diet. (b) Alteration of the lncRNA TCONS_00230836 level in β -TC6 cells and AML12 cells under non-stearic acid treatment. n = 5 mice per group; n = 4 independent cell cultures. **P < 0.01 versus the Islet/ β -TC6 group.




Figure 3

Confocal images of FISH for localization of the lncRNA TCONS_00230836 (red) in β -TC6 cells. Nuclei were stained with DAPI (blue). 18S, probe for 18S rRNA, was used for cytoplasmic localization. U6, probe for U6 snRNA, was taken as a representative of nuclear localization. Scale bar, 10 μ m.




Figure 4

Inhibition of the TCONS_00230836 alleviated stearic acid-induced insulin secretion reduction in β -TC6 cells and islets. (a) The expression of the IncRNA TCONS_00230836 in islets under high-stearic acid diet. (b) The IncRNA TCONS_00230836 level in β -TC6 cells under stearic acid treatment. (c) Representative images of FISH in β -TC6 cells showing the change of IncRNA TCONS_00230836 expression after transfection of the IncRNA TCONS_00230836 Smart Silencer in the presence of stearic acid. Scale bar, 10 μ m. (d) Effect of the IncRNA TCONS_00230836 Smart Silencer on GSIS in islets isolated from mice in the presence of stearic acid. (e) Restoration of stearic acid-induced glucose-stimulated insulin secretion (GSIS) impairment after transfection of the IncRNA TCONS_00230836 Smart Silencer in β -TC6 cells. Ctrl, control group; SA, stearic acid; si-Inc836, Smart Silencer for IncRNA TCONS_00230836. * $P < 0.05$, ** $P <$

0.01 versus the control group; ^{*}P < 0.05, ^{**}P < 0.01 versus the stearic acid group. n =5 independent animal experiments per group n = 4 independent cell cultures per group.

Fig. 5





Figure 5

The lncRNA TCONS_00230836 did not significantly reverse stearic acid-induced β -TC6 cell apoptosis. (a, b) Effect of the Smart Silencer for TCONS_00230836 on stearic acid-induced cell viability by CCK8 assay and cell death by LDH measurement, separately. (c) FCM revealed that the number of apoptotic cells was not altered after TCONS_00230836 inhibition under stearic acid treatment. (d) No significant changes in the expression of cleaved-caspase3, cleaved-Parp1, and Bcl-2 after inhibition of the lncRNA TCONS_00230836 by western blot analysis. NS, not significant; SA, stearic acid; si-Inc836, Smart Silencer for lncRNA TCONS_00230836. n = 3 independent cell cultures per group. ^{*}P < 0.05, ^{**}P < 0.01 versus the control group; ^AP < 0.05, ^{AA}P < 0.01 versus the stearic acid group.

Fig. 6**Figure 6**

Knockdown of the lncRNA TCONS_00230836 apparently relieved stearic acid-induced ER stress-related proteins in β -TC6 cells. (a) Effect of the lncRNA TCONS_00230836 Smart Silencer on the expression of Phospho-PERK and Phospho-eIF2 α proteins in stearic acid-treated β -TC6 cells. (b) Western blot analysis showed the alteration of ATF-6 and IRE1 α protein levels after transfection of the TCONS_00230836 Smart Silencer in β -TC6 cells exposed to stearic acid. NS, not significant; SA, stearic acid; si-Inc836, transfection of the Smart Silencer for the lncRNA TCONS_00230836. * $P < 0.05$ versus the control group; ^ $P < 0.05$ versus the stearic acid group. $n = 3$ independent cell cultures in each group.

Fig. 7





Figure 7

Silencing lncRNA TCONS_00230836 did not markedly alter islet apoptosis exposure to stearic acid. (a, b) Assessment of cell viability in HSD mice islets treated with Smart Silencer for TCONS_00230836 by CCK8 assay and cell death by LDH measurement, separately. (c) Western blot analysis verifying the changes in the expression of cleaved-caspase3, cleaved-Parp1, and Bcl-2 after inhibition of the lncRNA TCONS_00230836. NS, not significant; SA, stearic acid; si-Inc836, Smart Silencer for lncRNA TCONS_00230836. n = 5 independent animal experiments per group. *P < 0.05, **P < 0.01 versus the control group; ^P < 0.05, ^^P < 0.01 versus the SA group.

Fig. 8**Figure 8**

Inhibition of the lncRNA TCONS_00230836 apparently recovered ER stress-related proteins in stearic acid-treated mice islets. (a) The expression of Phospho-PERK and Phospho-eIF2 α proteins in mice islet under high-stearic-acid treatment by the lncRNA TCONS_00230836 Smart Silencer. (b) The alteration of ATF-6 and IRE1 α protein levels after transfection of the TCONS_00230836 Smart Silencer in islet exposed to high level of stearic-acid. NS, not significant; SA, stearic-acid; si-lnc836, transfection of the Smart Silencer for the lncRNA TCONS_00230836. * $P < 0.05$ versus the control group; $\Delta P < 0.05$ versus SA group. $n = 5$ independent animal experiments per group.




Figure 9

Alterations of candidate ER stress-related mRNAs expressions after transfection of lncRNA TCONS_00230836 Smart Silencer. (a, b) The positive and negative co-expression network of lncRNA TCONS_00230836 and target mRNAs specifically in stearic acid-treated β -TC6 cells. Red circle nodes represent up-regulated lncRNA TCONS_00230836. Genes labeled with yellow and blue squares refer to up-regulated and down-regulated mRNAs, respectively. Red lines represent positive correlation and green lines refer to negative correlation. (c) Effect of lncRNA TCONS_00230836 Smart Silencer on stearic acid-increased ER stress-related mRNAs expressions. (d) Changes in the down-regulated ER stress-related mRNAs after silencing lncRNA TCONS_00230836. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, versus the ctrl group; ^ $P < 0.05$, ^^ $P < 0.01$ versus SA group.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile5.docx](#)

- [Additionalfile4.docx](#)
- [Additionalfile3.docx](#)
- [Additionalfile2.docx](#)
- [Additionalfile1.docx](#)