

Necrotizing Funisitis As an Intrauterine Manifestation of Cryopyrin-Associated Periodic Syndrome: A Case Report and Review of The Literature

Kyoko Yokoi (✉ kyoyo0410@yahoo.co.jp)

Komaki City Hospital <https://orcid.org/0000-0002-5972-1131>

Sachiko Minamiguchi

Kyoto University Graduate School of Medicine Faculty of Medicine: Kyoto Daigaku Daigakuin Igaku Kenkyuka Igakubu

Yoshitaka Honda

Kyoto University Graduate School of Medicine

Mizuho Kobayashi

Nagoya West Medical Center

Satoru Kobayashi

Nagoya West Medical Center

Ryuta Nishikomori

Kurume University School of Medicine

Case Report

Keywords: cryopyrin associated periodic syndrome, neonatal-onset multisystem inflammatory disease/ chronic infantile neurologic cutaneous and articular syndrome, funisitis, preterm

DOI: <https://doi.org/10.21203/rs.3.rs-76666/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Cryopyrin-associated periodic syndrome (CAPS) is a life-long, autoinflammatory disease associated with a gain-of-function mutation in the nucleotide-binding domain, leucine-rich repeat family, pyrin domain containing 3 (*NLRP3*) gene, which result in uncontrolled production of IL-1 β and chronic inflammation. Chronic infantile neurologic cutaneous and articular (CINCA) syndrome/neonatal-Onset multisystem inflammatory disease (NOMID) is the most severe form of CAPS. Although the first symptoms may be presented at birth, there are few reports on the involvement of the placenta and umbilical cord in the disease. Therefore, we present herein a preterm case of CINCA/NOMID syndrome and confirms intrauterine-onset inflammation with conclusive evidence by using fetal and placental histopathological examination.

Case presentation: The female patient was born at 33weeks of gestation by emergency caesarean section and weighted at 1,514 g. The most common manifestations of CINCA/NOMID syndrome including recurrent fever, urticarial rash, and ventriculomegaly due to aseptic meningitis were presented. She also exhibited atypical symptoms such as severe hepatosplenomegaly with cholestasis. The genetic analysis of *NLRP3* revealed a heterozygous c.1698C>A (p.Phe566Leu) mutation, and she was diagnosed with CINCA/NOMID syndrome. Further, a histopathological examination revealed necrotizing funisitis, mainly inflammation of the umbilical artery, along with focal neutrophilic and lymphocytic villitis.

Conclusion: The necrotizing funisitis, which only involved the artery, was an unusual observation for chorioamnionitis. These evidences suggest that foetal inflammation, probably due to overproduction of IL-1 β , caused tissue damage in utero, and the first symptom of a newborn with CINCA/NOMID.

Background

Cryopyrin-associated periodic syndrome (CAPS) is a life-long, autoinflammatory disease associated with a gain-of-function mutation in the nucleotide-binding domain, leucine-rich repeat family, pyrin domain containing 3 (*NLRP3*) gene, which codes for cryopyrin [1]. Cryopyrin is a vital part of the inflammasome complex, which activates caspase-1 and causes subsequent production of interleukin-1 beta (IL-1 β). Abnormalities in cryopyrin cause uncontrolled production of IL-1 β , resulting in chronic inflammation and tissue damage in CAPS [2, 3]. CAPS comprise of three diseases with specific clinical features and disease activity; the mildest form of which is familial cold auto-inflammatory syndrome, Muckle-Wells syndrome is the moderate form, and the most severe form is chronic infantile neurologic cutaneous and articular (CINCA)/neonatal-onset multisystem inflammatory disease (NOMID) syndrome. All three forms are caused by the same gene, *NLRP3* [4, 5]. The prevalence of CINCA/NOMID syndrome is estimated to be 1 in 1,000,000, and approximately 40 people were diagnosed with CINCA/NOMID syndrome in Japan [6]. Although the first symptoms of CINCA/NOMID syndrome may be present at birth, there are few reports on the involvement of the placenta and umbilical cord in the disease [7]. Therefore, we present herein a preterm case of CINCA/NOMID that was complicated with the necrotizing funisitis which only involved the artery, which was confirmed by using fetal and placental histopathological examination.

Case Presentation

The female patient was born at 33 weeks and 5 days of gestation by emergency cesarean section following presentation of pregnancy complication including polyhydramnios, spontaneous preterm labor, and elevated maternal C-reactive protein (CRP). Her parents were nonconsanguineous, and family history was unremarkable. Apgar scores were 5 at 1 min, and 7 at 5 min, with birthweight of 1,514 g (-1.8 SD), length of 42 cm (-0.74 SD), and head circumference of 28.3 cm (-1.29 SD). She presented with severe hepatosplenomegaly and mild ventriculomegaly at birth, as well as urticarial rash on the body and limbs after a few hours of birth. The blood examination at birth revealed CRP at 6.4 mg/dL, white blood cell count (WBC) of 34650 / μ L, red blood cell count of 323×10^4 / μ L, hemoglobin 12.7 g/dL and platelet 7×10^4 / μ L. In addition, the examination of the cerebrospinal fluid (CSF) showed a WBC cell count of 667 / μ L (polymorphonuclear: mononuclear 4:1), glucose at 21 mg/dL, and protein at 191 mg/dL. No pathogenic bacteria were isolated from the patient, her CRP continued to increase to 15 mg/dL despite intensive antibiotics treatment. The neonatal-onset rash and aseptic meningitis, combined with persistent elevation of CRP suggested the diagnosis of auto-inflammatory disease, more specifically CAPS. Urinary excretion of mevalonic acid was not elevated, and the possibility of hyper IgD syndrome was excluded. After obtaining an informed consent from her parents, we performed genetic analysis of *NLRP3* on her whole blood, which revealed heterozygous *NLRP3* c.1698C > A (p.Phe566Leu) variant that has been reported as disease-causing in infefers database (<https://infefers.umai-montpellier.fr/web/search.php?n=4>) [8]. Based on the clinical manifestations and genetic analysis, she was finally diagnosed with CINCA/NOMID.

Meanwhile, the hepatosplenomegaly with cholestasis gradually deteriorated to a total bilirubin of 5.4 mg/dL and direct bilirubin of 3.5 mg/dL, aspartate aminotransferase of 147 IU/L and alanine aminotransferase of 83 IU/L. Contrast-enhanced computed tomography of the abdomen was normal except severe hepatosplenomegaly. Specific IgM antibody serology tests against toxoplasmosis, cytomegalovirus, rubella virus, syphilis, and herpes simplex virus were negative. The plasma analysis of amino acids by high performance liquid chromatography was not characteristic of neonatal intrahepatic cholestasis caused by citrin deficiency such as elevated citrulline, tyrosine, phenylalanine, methionine and threonine. Bone marrow examination was performed, and no malignancy nor hemophagocytosis were observed. Oral ursodeoxycholic acid, fat-soluble vitamins and medium chain fatty acids containing milk (MCT milk) were started at 16 days of age, however, these treatments had no remarkable effect on her cholestasis.

At 70 days of age, after diagnosis was confirmed genetically, treatment with recombinant anti-human IL-1 β monoclonal antibody (canakinumab) was started at a dose of 2.5 mg/kg. The rash immediately disappeared, and inflammatory markers slightly decreased after the first administration. Cholestasis slowly improved and ursodeoxycholic acid, fat-soluble vitamins and MCT milk were discontinued at the age of 4 months. We finally increased the dose of canakinumab stepwise up to 7.8 mg/kg every 4–6 weeks, and no further complications from the use of canakinumab were noted since the age of 7 months.

The pathological findings of the placenta and umbilical cord

The umbilical cord was found to be edematous and contained yellow discoloration on the surface. Moreover, the placenta was friable, and weighted 615 g, which is a very heavy placenta, greater than 90th percentile for the gestational age. The histopathological examination revealed necrotizing funisitis, namely inflammation of the umbilical cord (Fig.1A), and along with focal neutrophilic and lymphocytic villitis (Fig.1B). The neutrophils were not infiltrated in the chorioamnion. While inflammatory cells presented ring-shaped infiltration on the surface of one of the umbilical arteries and within the Wharton's jelly, there was no inflammatory response in umbilical vein (Fig.1A). These inflammatory cells were mainly neutrophils and macrophages, demonstrated by immunohistochemical positive for myeloperoxidase (Fig.1C), a portion of which shows CD68 positivity.

Discussion

CINCA/NOMID syndrome is the most severe form of CAPS, which is characterized by neonatal onset and presents with recurrent fever and urticarial rash. Without appropriate treatments, chronic inflammation continues, causing neurological manifestations, arthritis, arthropathy, and uveitis [9, 10]. In Japan, only canakinumab has been approved since 2011 for the treatment of all three forms of CAPS [11]. The definite diagnosis for CINCA/NOMID relies mainly on genetic analysis of *NLRP3*, which has more than 210 reported variants [8]. Further, approximately 28 ~ 35% of all the CINCA/NOMID patients carry an *NLRP3* mutation in somatic mosaicism state [12, 13], which causes patients to present with milder neurologic symptoms compared to those with germline mutation of the same variant. However, no additional differences in symptoms was observed in age at disease onset, skin symptoms and joint involvement, whereas, the p.Phe566Leu variant was previously reported as a CAPS disease-causing mutation within somatic mosaicism cohort [14]. Our patient was found to have p.Phe566Leu variant as a heterozygous germline mutation, which might explain the most severe phenotype including continuous inflammatory biological markers and atypical symptoms.

Furthermore, hepatosplenomegaly with normal liver functions was occasionally described in CINCA/NOMID patients, and liver biopsy demonstrated non-specific presentation for chronic inflammation in the parenchyma and subcapsular area in such a case [7]. However, the patients with cholestasis were only reported by Paccaud et al [14]. Their patient was identified as heterozygous for the p.Glu567Lys mutation and had many similar characteristics to our case. First, both cases were born at preterm 33 weeks of gestation, complicated with polyhydramnios. Second, hepatosplenomegaly with cholestasis was initially severe and gradually improved in parallel to decreasing inflammatory reaction. Following detailed examination, our patient did not have any other disease associated with cholestasis, and her symptoms gradually improved as CRP decreased. The observed chronic inflammation caused the liver damage in preterm neonate, resulting in hepatosplenomegaly and cholestasis. It is of great interest whether the treatment with canakinumab is able to subside the occurrence of complications on hepatobiliary system.

In addition, p.Glu567Lys mutation is the most frequently identified mutation among the somatic mosaicism cohort [15]. The coincidence might indicate that the cholestasis might occur in very severe

inflammatory conditions due to gene dosages effect of germline mutations which can cause CINCA/NOMID even in the somatic mosaicism. This hypothesis remains to be evaluated by further accumulating the CINCA/NOMID neonatal cases, especially focusing on their neonatal manifestations.

The preterm, intrauterine growth retardation, polyhydramnios, and placental and umbilical abnormalities were often presented in CINCA/NOMID syndrome. It has been reported that histological examination of the placenta from CINCA/NOMID syndrome has revealed vascular thrombosis, microcalcification and polymorphonuclear cell infiltrates [7, 14]. Table 1 describes intrauterine manifestation of CINCA/NOMID syndrome patients in previous reports. Though two polyhydramnios have been reported, making our case the third, it was not possible to perform the detailed search for polyhydramnios due to an emergency cesarean section.

Table 1
Intrauterine manifestation of CINCA/NOMID patients in previous reports.

Intrauterine manifestation	Reference	The number of cases	Mutation	Commentary
Abnormal amount of amniotic fluid	Sandra G, et al (1983) [16]	1	N.D.	Polyhydramnios
	Prieur AM, et al (1987) [7]	2	N.D.	Oligohydramnios
	Caroli F, et al (2006) [17]	1	M406I	Oligohydramnios
	Paccaud Y, et al (2014) [14]	1	E567K	Polyhydramnios
	Our case	1	F566L	Polyhydramnios
Histological findings of placenta and umbilical cord	Prieur AM, et al (1987) [7]	1	N.D.	Placenta with thrombosis and calcification Small villi and umbilical cord with polymorphonuclear infiltration
	Paccaud Y, et al (2014) [14]	1	E567K	Umbilical cord infection
	Madison E, et al (2019) [22]	1	N.D.	Necrotizing funisitis
	Our case	1	F566L	Neutrophilic and lymphocytic villitis Necrotizing umbilical arteritis
N.D.: not described				

Necrotizing funisitis, which is often associated with increased rates of stillbirth, perinatal infection, and preterm delivery, is sometimes accompanied with congenital syphilis [18]. In our case, the possibility of congenital syphilis was excluded because of the absence of the specific IgM antibodies. Acute chorioamnionitis provides evidence of a maternal host response, whereas funisitis represents a foetal inflammatory response [19]. These inflammatory responses are characterized by the infiltration of neutrophils and release of cytokines [19, 20]. Moreover, funisitis is often associated with inflammation of the chorioamnion due to ascending infection. The inflammation of the umbilical vessels typically begins in the vein (phlebitis) and is followed by involvement of the arteries (arteritis) and the Wharton's jelly [21]. Since our case was not complicated with severe chorioamnionitis, it is likely that the necrotizing funisitis that only involved the artery, was not secondary to chorioamnionitis, but due to foetal origin inflammation. These evidences suggest that foetal inflammation, probably due to overproduction of IL-1 β by CINCA/NOMID, might cause necrotizing funisitis in utero. According to previous reports, all histopathological findings revealed funisitis or inflammation of umbilical cords, which might be quite usual manifestation in CINCA/NOMID syndrome (Table 1).

Moreover, our patient was complicated with villitis, namely inflammation of chorioamniotic villi. Some villitis is caused by an infectious agent, however, most chronic inflammatory lesions are of unknown aetiology [23] [24]. Excessive foetal IL-1 β might have injured the villi to induce a maternal inflammatory response, causing the elevated maternal CRP before the emergency caesarean section.

Conclusion

We emphasize that the histopathological examination of placenta and umbilical cord can be one of the clues indicating intrauterine-onset inflammation and the first symptom of a newborn with CINCA/NOMID.

List Of Abbreviations

CAPS: cryopyrin-associated periodic syndrome, *NLRP3*: nucleotide-binding domain, leucine-rich repeat family, pyrin domain containing 3, IL-1 β : interleukin-1 beta, CINCA/ NOMID : chronic infantile neurologic cutaneous and articular syndrome/neonatal-onset multisystem inflammatory disease, CRP: c-reactive protein, WBC: white blood cell, MCT milk: medium chain fatty acids containing milk,

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: The patient's parents gave informed consent for publication.

Availability of data and materials: The data are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: This study received no funding.

Authors' contributions: KY contributed medical care for this patient and drafted the manuscript with support from SK and RN. HY provided the genetic analysis. KM and SM undertook the histopathological examination. RN supervised the patient's diagnosis and treatment. All authors read and approved the final manuscript.

Acknowledgements

We are grateful to Yosuke Shigematsu (Department of Pediatrics, Faculty of Medical Sciences, University of Fukui) for the measurement of mevalonic acid.

Authors' information (optional): Not applicable

References

1. Hoffman HM, Mueller JL, Broide DH, Wanderer AA, Kolodner RD. Mutation of a new gene encoding a putative pyrin-like protein causes familial cold autoinflammatory syndrome and Muckle-Wells syndrome. *Nat Genet.* 2001; 29(3):301-5. doi:1038/ng756.
2. Neven B, Callebaut I, Prieur AM, Feldmann J, Bodemer C, Lepore L, et al. Molecular basis of the spectral expression of CIAS1 mutations associated with phagocytic cell-mediated autoinflammatory disorders CINCA/NOMID, MWS, and FCU. *Blood.* 2004; 103:2809–2815. doi: 1182/blood-2003-07-2531.
3. Gattorno M, Tassi S, Carta S, Delfino L, Ferlito F, Pelagatti MA, et al. Pattern of interleukin-1 β secretion in response to lipopolysaccharide and ATP before and after interleukin-1 blockade in patients with CIAS1 mutations. *Arthritis Rheum.* 2007; 56:3138–3148. doi: 1002/art.22842.
4. Hentgen V, Despert V, Leprêtre AC, Cuisset L, Chevrant-Breton J, Jégo P, et al. Intrafamilial variable phenotypic expression of a CIAS1 mutation: from Muckle-Wells to chronic infantile neurological cutaneous and articular syndrome. *J Rheumatol.* 2005; 32(4): 747-51.
5. Feldmann J, Prieur AM, Quartier P, Berquin P, Certain S, Cortis E, et al. Chronic infantile neurological cutaneous and articular syndrome is caused by mutations in CIAS1, a gene highly expressed in polymorphonuclear cells and chondrocytes. *Am J Hum Genet.* 2002; 71(1): 198–203. doi: 1086/341357.
6. Nakamichi S, Origuchi T, Fukui S, Yoda A, Matsubara H, Nagaura Y, et al. A Rare Case of Cryopyrin-associated Periodic Syndrome in an Elderly Woman with NLRP3 and MEFV Mutations. *Intern Med.* 2019; 58(7): 1017-1022. doi: 10.2169/internalmedicine.1401-18.
7. Prieur AM, Griscelli C, Lampert F, Truckenbrodt H, Guggenheim MA, Lovell DJ, et al. A chronic, infantile, neurological, cutaneous and articular (CINCA) syndrome. A specific entity analysed in 30 patients. *Scand J Rheumatol Suppl.* 1987; 66:57-68. doi:3109/03009748709102523.

8. The registry of hereditary auto-inflammatory disorders mutations.
<http://fmf.igh.cnrs.fr/ISSAID/infevers/>
9. Hoffman HM. Hereditary immunologic disorders caused by pyrin and cryopyrin. *Curr Allergy Asthma Rep.* 2007; 7(5): 323-30. doi: [1007/s11882-007-0049-4](https://doi.org/10.1007/s11882-007-0049-4).
10. Cuisset L, Jeru I, Dumont B, Fabre A, Cochet E, Le Bozec J, et al. Mutations in the autoinflammatory cryopyrin-associated periodic syndrome gene: epidemiological study and lessons from eight years of genetic analysis in France. *Ann Rheum Dis.* 2011; 70 3: 495-9. doi: [1136/ard.2010.138420](https://doi.org/10.1136/ard.2010.138420).
11. Yokota S, Imagawa T, Nishikomori R, Takada H, Abrams K, Lheritier K, et al. Long-term safety and efficacy of canakinumab in cryopyrin-associated periodic syndrome: results from an open-label, phase III pivotal study in Japanese patients. *Clin Exp Rheumatol.* 2017; 35 Suppl 108 6: 19-26.
12. Tanaka N, Izawa K, Saito MK, Sakuma M, Oshima K, Ohara O, et al. High incidence of NLRP3 somatic mosaicism in patients with chronic infantile neurologic, cutaneous, articular syndrome: results of an International Multicenter Collaborative Study. *Arthritis Rheum* 2011; 63 11: 3625-32. doi:[1002/art.30512](https://doi.org/10.1002/art.30512).
13. Izawa K, Hijikata A, Tanaka N, Kawai T, Saito MK, Goldbach-Mansky R, et al. Detection of base substitution-type somatic mosaicism of the NLRP3 gene with >99.9% statistical confidence by massively parallel sequencing. *DNA Res.* 2012; 19: 143-52. doi: [10.1093/dnares/dsr047](https://doi.org/10.1093/dnares/dsr047).
14. Paccaud Y, Berthet G, Von Scheven-Gete A, Vaudaux B, Mivelaz Y, Hofer M, et al. Neonatal treatment of CINCA syndrome. *Pediatr Rheumatol Online J.* 2014; 12:52. doi:[1186/1546-0096-12-52](https://doi.org/10.1186/1546-0096-12-52).
15. Labrousse M, Kevorkian-Verguet C, Boursier G, Rowczenio D, et al. Mosaicism in autoinflammatory disease: Cryopyrin-associated periodic syndrome (CAPS) and beyond. A systemic review. *Crit Rev Clin Lab Sci.* 2018 Sep;55(6):432-442. doi:[10.1080/10408363.2018.1488805](https://doi.org/10.1080/10408363.2018.1488805).
16. Hassink SG, Goldsmith DP. Neonatal onset multisystem inflammatory disease. *Arthritis Rheum.* 1983 May;26(5):668-73. doi: [10.1002/art.1780260515](https://doi.org/10.1002/art.1780260515)
17. Caroli F, Pontillo A, D'Osualdo A, Travan L, Ceccherini I, et al. Clinical and genetic characterization of Italian patients affected by CINCA syndrome. *Rheumatology (Oxford).* 2007 Mar;46(3):473-8. doi: [10.1093/rheumatology/kel269](https://doi.org/10.1093/rheumatology/kel269).
18. Guarner J, Southwick K, Greer P, Barlett J, Santander A, Blanco S, et al., Testing umbilical cords for funisitis due to *Treponema pallidum* infection, Bolivia. *Emerg Infect Dis.* 2000; 6(5): 487–492. doi: [3201/eid0605.000507](https://doi.org/10.3201/eid0605.000507).
19. Kim CJ, Romero R, Chaemsathong P, Chaiyasit N, Yoon BH, Kim YM. Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. *Am J Obstet Gynecol.* 2015; 213(4 Suppl): S29-52. doi:[1016/j.ajog.2015.08.040](https://doi.org/10.1016/j.ajog.2015.08.040).
20. Kim CJ, Yoon BH, Park SS, Kim MH, Chi JG. Acute funisitis of preterm but not term placentas is associated with severe fetal inflammatory response. *Hum Pathol.* 2001; 32(6): 623-9. doi: [1053/hupa.2001.24992](https://doi.org/10.1053/hupa.2001.24992).

21. Redline RW, Faye-Petersen O, Heller D, et al. Amniotic infection syndrome: nosology and reproducibility of placental reaction patterns. *Pediatr Dev Pathol*. 2003; 6(5): 435-48. doi:1007/s10024-003-7070-y.
22. Kerley ME, Ahmed S, Ramasubramanian A. Bilateral Chorioretinitis From NOMID syndrome. *J Pediatr Ophthalmol Strabismus*. 2019 May 22;56(3):204. doi:10.3928/01913913-20190220-01.
23. Redline RW. Villitis of unknown etiology: noninfectious chronic villitis in the placenta. *Hum pathol*. 2007; 38 (10): 1439-46. doi: 1016/j.humpath.2007.05.025.
24. Kim CJ, Romero R, Chaemsaihong P, Kim JS. Chronic inflammation of the placenta: definition, classification, pathogenesis, and clinical significance. *Am J Obstet Gynecol*. 2015; 213(4 Suppl): S53-69. doi:1016/j.ajog.2015.08.041.

Figures




Figure 1

Microscopic findings of the placenta and umbilical cord. a. Necrotizing funisitis: the umbilical cord showed ring-shaped infiltration of inflammatory cells around the artery (arrow and inset), but not in the vein (arrow head). (Hematoxylin an eosin (HE) staining). b. Focal neutrophilic and lymphocytic villitis: destructive inflammation of chorionic villus (arrow), followed by mild collapse of villous vessels (HE staining). c. Immunohistochemical staining of myeloperoxidase: the inflammation around the umbilical artery and perivascular Wharton's jelly caused mainly by macrophages and neutrophils.