

Inflammation drives synucleinopathy propagation

Tae-Kyung Kim^{1,7#}, Eun-Jin Bae^{1,#}, Byung Chul Jung^{1,#}, Minsun Choi¹, Soo Jean Shin¹, Jeong Tae Kim¹, Min Kyo Jung², Ayse Ulusoy³, Mi-Young Song⁴, Jun Sung Lee¹, He-Jin Lee^{5,6}, Donato A. Di Monte³ and Seung-Jae Lee^{1,*}

¹Department of Biomedical Sciences and Neuroscience Research Institute, Seoul National University College of Medicine, Seoul 03080, Korea

²Neural circuits research group, Korea Brain Research Institute, Daegu 41068, Korea

³German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany

⁴Department of Biomedical Science and Technology, Konkuk University, Seoul 143-701, Korea

⁵Department of Anatomy, Konkuk University, Seoul 05029, Korea

⁶IBST, Konkuk University, Seoul 05029, Korea

⁷Department of Exercise Physiology and Sport Science Institute, Korea National Sport University, Seoul 05541, Korea

*Corresponding author: Department of Biomedical Sciences, Seoul National University College of Medicine, 103 Daehak-ro, Jongro-gu, Seoul 03080, Korea, Tel: +82-2-3668-7037, Fax: +82-2-447-5683, Email: sjlee66@snu.ac.kr

#These authors contributed equally to this work.

Description of Additional Supplementary Files

File Name: Supplementary Table 2

Description: The list of differentially expressed genes (DEGs) in the mice treated with WT-syn and V40G-syn relative to those of PBS-treated mice. (\log_2 [fold change] ≥ 0.5 , $p < 0.05$).

File Name: Supplementary Table 3

Description: The list of common DEGs in the mice treated with WT-syn and V40G-syn relative to those of PBS-treated mice. Related to Figure 3. (\log_2 [fold change] ≥ 0.5 , $p < 0.05$).