

Prevalence of microhematuria in renal colic and urolithiasis: a systematic review and meta-analysis

CURRENT STATUS: UNDER REVIEW

BMC Urology  BMC Series

Bruno Minotti
Kantonsspital Sankt Gallen

✉ bruno.minotti@kssg.ch *Corresponding Author*
ORCID: <https://orcid.org/0000-0002-4820-9968>

Giorgio Treglia
Repubblica e Cantone Ticino Ente Ospedaliero Cantonale

Maria Rosa Pascale
Repubblica e Cantone Ticino Ente Ospedaliero Cantonale

Samuele Ceruti
Clinica Luganese

Laura Cantini
Repubblica e Cantone Ticino Ente Ospedaliero Cantonale

Luciano Anselmi
Repubblica e Cantone Ticino Ente Ospedaliero Cantonale

Andrea Saporito
Repubblica e Cantone Ticino Ente Ospedaliero Cantonale

DOI:

10.21203/rs.2.17200/v1

SUBJECT AREAS

Urology & Nephrology

KEYWORDS

renal colic, urolithiasis, microhematuria, stone score

Abstract

Background This systematic review and meta-analysis aims to investigate the prevalence of microhematuria in patients presenting with suspected acute renal colic and/or confirmed urolithiasis at the emergency department.

Methods A comprehensive literature search was conducted to find relevant data on prevalence of microhematuria in patients with suspected acute renal colic and/or confirmed urolithiasis. Data from each study regarding study design, patient characteristics and prevalence of microhematuria were retrieved. A random effect-model was used for the pooled analyses.

Results Forty-nine articles including 15'860 patients were selected through the literature search. The pooled microhematuria prevalence was 77% (95%CI: 73-80%) and 84% (95%CI: 80-87%) for suspected acute renal colic and confirmed urolithiasis, respectively. This proportion was much higher when the dipstick was used as diagnostic test (80% and 90% for acute renal colic and urolithiasis, respectively) compared to the microscopic urinalysis (74% and 78% for acute renal colic and urolithiasis, respectively).

Conclusions This meta-analysis revealed a high prevalence of microhematuria in patients with acute renal colic (77%), including those with confirmed urolithiasis (84%). Intending this prevalence as sensitivity, we reached moderate values, which make microhematuria alone a poor diagnostic test for acute renal colic or urolithiasis. Microhematuria could possibly still important to assess the risk in patients with renal colic.

1. Background

Renal colic is caused by the presence of stones in the urinary tract and it is characterized by sudden onset of severe loin pain, radiating to the flank, groin, and testes or *labia majora* [1]. Incidence amounts to 240 per 100'000 persons [2] with a prevalence up to 10%; men are commonly more affected than women with a ratio of 3-2:1 [3]. Lifetime risk is up to 19% in men and 9% in women [4], varying depending on geographic location and increasing constantly over last years [5]. Guidelines for the diagnostic pathway suggest assessing (micro)hematuria, while the gold standard of imaging is unenhanced multi-detector computed tomography (MDCT) [1]. As diagnostic tool the STONE Score

was developed and validated; this score includes parameters as sex, duration of pain prior to presentation, race, nausea, vomiting and microhematuria [6]. Microhematuria prevalence in suspected renal colic has been studied in several trials, ranging from 55% [7] to 93% [8, 9]. In order to better understand the difference existing in prevalence range, we performed a meta-analysis of studies dealing with microhematuria by suspected acute renal colic and/or confirmed urolithiasis.

2. Methods

This systematic review and meta-analysis conforms to the statement on Preferred Reporting Items for Systematic reviews and Meta-Analyses [10].

2.1 Search strategy

A literature search of the electronic PubMed/MEDLINE database and Cochrane Central Register of Controlled Trials (CENTRAL), without language restriction, was carried out from inception to October 11, 2018. A search algorithm was established using a combination of the following terms: A) renal colic AND urolithiasis (Problem), B) urinalysis (Intervention), C) microhematuria (Outcome). The final search query is reported in Appendix 1. Reference lists of the retrieved articles were also screened for additional studies.

2.2 Eligibility criteria

We included in this systematic review and meta-analysis studies which filled the following inclusion criteria: a) original article published in peer-reviewed journal; b) studies including adults only; c) patients presenting with acute renal colic at the emergency department; d) studies reporting data on microhematuria.

Exclusion criteria were: a) articles not within the field of interest of this review; b) review articles, letters or editorials; c) case reports or case series (less than 10 patients included); d) articles with possible patient data overlap.

2.3 Study selection

Titles and abstracts of the retrieved studies were independently reviewed by two researchers (MP, GT), applying the inclusion and exclusion criteria mentioned above. Articles were rejected if they were clearly ineligible. The full texts of the potentially eligible articles were reviewed independently by the

same researchers to confirm or exclude their eligibility for inclusion. Disagreements were resolved in a consensus meeting.

2.4 Data extraction

For each included study, one author (MP) manually extracted data relevant to the review aims using a customized form. Information regarding basic study data (authors, year of publication, country of origin, type of study), patient characteristics (number of patients, mean age, gender), methods (microhematuria test, microhematuria definition) and outcomes (number of patients with microhematuria, microhematuria prevalence) were retrieved. The number of patients with microhematuria and microhematuria prevalence were also extracted for patients with confirmed urolithiasis, where available. Diagnostic methods for detection of stones were also retrieved. One other author (GT) independently checked all extracted data.

2.5 Outcome measures

The primary outcome was the percentage of microhematuria among patients presenting with suspected acute renal colic at the emergency department. The secondary outcome was the percentage of microhematuria among patients presenting with acute renal colic and confirmed urolithiasis at the emergency department.

2.6 Quality assessment

The overall quality of the studies included in the systematic review was critically appraised based on the revised “Quality Assessment of Diagnostic Accuracy Studies” tool (QUADAS-2). This tool comprises four domains: patient selection, index test, reference standard, and flow and timing. Each domain was assessed in terms of risk of bias, and the first three domains were also assessed in terms of concerns regarding applicability.

2.7 Statistical analysis

Microhematuria prevalence was defined as the ratio between the number of patients with suspected acute renal colic with microhematuria detected by urinalysis or dipstick and the total number of patients with suspected acute renal colic who underwent the analysis. This proportion was calculated also for patients presenting with acute renal colic and confirmed urolithiasis.

Pooled analyses of the proportion of microhematuria detected by urinalysis or dipstick were performed using data retrieved from the selected studies. When microhematuria was assessed using both urinalysis and dipstick, the test with the better outcome was chosen. Subgroup analyses taking into account the microhematuria test were planned.

A random-effects model was used for statistical pooling of the data, taking into account the heterogeneity between studies. The different weight of each study in the pooled analysis was related to the different sample size. Pooled data were presented with their respective 95% confidence interval (95%CI) values, and data were displayed using plots.

Heterogeneity was estimated by using the I-square index (I^2), which describes the percentage of variation across studies that is due to heterogeneity rather than chance [11].

Publication bias was assessed through the Egger's test [12].

Statistical analyses were performed using the StatsDirect software version 3 (StatsDirect Ltd., Cambridge, UK).

3. Results

3.1 Literature search

The literature search from PubMed/MEDLINE and Cochrane CENTRAL databases yielded a total of 1,377 records. After reviewing titles and abstracts, 77 were selected as potentially eligible articles. The full text was retrieved for all. Following eligibility's assessment, 31 articles did not meet the inclusion criteria and were excluded from the systematic review. Within the selected articles, screening of the reference lists allowed to add 3 additional records. Finally, 49 studies [7-9, 13-58] including 15'860 patients were identified as potentially relevant and were selected for the systematic review and meta-analysis. All of the included studies except two [30, 50] were published in English. These studies covered the period from inception to October 11, 2018. Search results and articles' selection are displayed in a PRISMA flow chart (Figure 1).

3.2 Selected studies

The characteristics of selected studies are reported in Table 1. The studies were conducted in different countries worldwide (Europe, North America, Asia, Africa). The sample size of the included

trials ranged from 32 to 2,218 adults presenting to the emergency department or urology clinic with acute renal colic. Most of the studies were observational with a prospective (19) or retrospective (29) or mixed (1) design.

Microhematuria was tested by urinalysis in 32 studies, urine dipstick in 10 and both methods in 7. Definition of microhematuria was different among the included studies. Six studies included also patients presenting with macroscopic hematuria [14, 17, 19, 22, 26, 50]. Details on the microhematuria test are reported in Table 2.

3.3 Quality assessment

Overall quality assessment of the studies included in the systematic review according to QUADAS-2 tool is reported in Supplemental Figure 1.

3.4 Microhematuria prevalence and suspected acute renal colic

Primary outcome characteristics on microhematuria prevalence in patients with suspected acute renal colic are summarized in Table 2 and Figure 2.

Prevalence of microhematuria ranged from 35% to 94%, with a pooled estimate of 77% (95%CI: 73–80%) (Figure 2). The heterogeneity among the included studies was significant ($I^2 = 96\%$). A publication bias was detected by Egger's test ($p < 0.0001$).

Performing sub-group analyses taking into account different microhematuria tests, the pooled prevalence of microhematuria using urinalysis or urine dipstick was 74% (95%CI: 69–78%) and 80% (95%CI: 74–86%) respectively, without significant difference between two groups.

3.5 Microhematuria prevalence and confirmed urolithiasis

Secondary outcomes regarding main findings on microhematuria prevalence in patients with acute renal colic and confirmed urolithiasis are summarized in Table 3 and Figure 3.

Prevalence of microhematuria ranged from 44% to 100%, with a pooled estimate of 84% (95%CI: 80–87%) (Figure 3). Heterogeneity among the included studies was significant ($I^2 = 93\%$). A publication

bias was detected by Egger's test ($p = 0.0008$).

Performing sub-group analyses taking into account different microhematuria tests, the pooled prevalence of microhematuria using urinalysis or urine dipstick was 78% (95%CI: 74–82%) and 90% (95%CI: 83–95%), respectively.

4. Discussion

Many studies have evaluated the prevalence of microhematuria in patients with suspected acute renal colic (summarized in Table 1); this meta-analysis pooled data reported in the published studies to derive a more precise assessment. Overall, this systematic review and meta-analysis revealed a high prevalence of microhematuria in patients with acute renal colic (77%), including those with confirmed urolithiasis (84%). However, intending this prevalence as sensitivity, we reached moderate values, which make microhematuria alone a poor diagnostic test for acute renal colic, respectively for urolithiasis. In our meta-analysis heterogeneity was high; indeed, we found a poor definition regarding urine analysis across studies (see positive microhematuria definition in Table 2), with different cells count on microscopy, but also with various dipstick brands. Argyropoulos *et al.* [8] carried out a microscopic urinalysis when the dipstick was in doubt or with blood traces; microhematuria was confirmed in all of these cases. Thus, the authors concluded that urinary dipstick test is not inferior to microscopy. Bataille *et al.* [59] compared the sensitivity of urinary dipstick with microscopy and flow cytometry on *in vitro* contaminated human urine with human blood of volunteers at different concentrations. Urinary dipstick reached the best sensitivity, probably due to the ability to detect red blood cells after lysis, and was suggested as preferred test for screening of hematuria. Same results were previously reported by Kobayashi *et al.* [40] and Press *et al.* [52]. *De facto* we detected a trend toward a higher pooled prevalence of microhematuria by using urine dipstick compared to microscopic urinalysis. In addition Kobayashi *et al.* [40] found a relation between hematuria and pain onset, with the highest incidence of negative hematuria on day 3 and 4. Mefford *et al.* [18] showed an increased prevalence of hydronephrosis in patients with urolithiasis and negative microhematuria. As hydronephrosis is easy to screen with ultrasonography, Daniel *et al.* [60] developed the STONE PLUS Score with addition of point-of-care ultrasound of the kidney to the

original STONE Score. Presence of hydronephrosis improved the specificity up to 98% and helped to identify patients requiring urological intervention, without remarkably increasing risk stratification. Considering the moderate sensitivity of microhematuria in patients with renal colic, Xafis *et al.* [31] suggested to perform a MDCT without urinalysis as a prerequisite. This approach seems to show the best diagnostic accuracy; however, it would increase the number of MDCT with more costs and radiation exposure. Therefore, the focus should be placed in complicated urolithiasis (e.g., obstructive pyelonephritis) or dangerous alternative diagnosis. Rucker *et al.* [61] reported numerous diseases mimicking urolithiasis. Moore *et al.* [6] found a lower likelihood of a dangerous alternative diagnosis (< 2%) by using high STONE scores and suggested for this group the possibility to initially avoid compute tomography because till 90% of stones < 7 mm will pass through spontaneously [62]. With the same approach the American College of Emergency Physicians (ACEP) suggests in the Choosing Wisely group to avoid ordering computed tomography of the abdomen and pelvis in young except healthy emergency department patients (age <50) with known histories of kidney stones, or ureterolithiasis, presenting with symptoms consistent with uncomplicated renal colic [63]. Finally, alternative diagnosis mimicking renal colic could be found by ultrasonography at least in one study with the same accuracy as MDCT [64].

Some limitations and biases of our meta-analysis should be taken into account. Heterogeneity among studies may represent a potential source of bias in a meta-analysis. This heterogeneity is likely to arise through baseline differences among patients in the included studies (Table 1), or diversity in methodological aspects between different studies (Table 2). Unfortunately, we detected a significant heterogeneity in our meta-analysis. We believe that, beyond the various microhematuria tests (urinalysis vs dipstick), the most important source of heterogeneity could be the different definitions of microhematuria (Table 2).

In conclusion, microhematuria searched with urine dipstick showed higher diagnostic sensitivity and should be used in this setting as a “gold standard”; it is needed to calculate the STONE score, which can help to identify patients with decreased likelihood of a differential diagnosis, reducing costs and radiation exposure of MDCT. Finally, the concomitant use of ultrasound could increase the specificity

till 98% by hydronephrosis, identify patients requiring urological intervention and help to find alternative diagnosis in each risk group. Especially for searching differential diagnosis with ultrasound in patients with suspected renal colic, further studies should be undertaken. Larger prospective multicenter validation study of the STONE score could provide more definitive evidence.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Not applicable

Competing interests

No competing interests to declare

Funding

None

Authors' contribution

Concept: BM, AS. Literature search: MP, GT. Data interpretation: MP, GT, BM, AS. Manuscript writing: MP, GT, BM, AS. Critical review and final approval: all authors.

Acknowledgments

This work was carried out in collaboration with the Clinical Trial Unit of Ente Ospedaliero Cantonale (Ticino, Switzerland).

References

- [1] Bultitude M, Rees J. Management of renal colic. *Bmj*. 2012;345:e5499.
- [2] Tasian GE, Ross ME, Song L, Sas DJ, Keren R, Denburg MR, et al. Annual Incidence of Nephrolithiasis among Children and Adults in South Carolina from 1997 to 2012. *Clinical journal of the American Society of Nephrology: CJASN*. 2016;11:488-96.
- [3] Scales CD, Jr., Smith AC, Hanley JM, Saigal CS. Prevalence of kidney stones in the United States. *European urology*. 2012;62:160-5.
- [4] Curhan G. 47 - Nephrolithiasis. In: Gilbert SJ, Weiner DE, editors. *National Kidney Foundation Primer on Kidney Diseases (Sixth Edition)*. Philadelphia: W. B. Saunders; 2014. p. 405-11.

- [5] Romero V, Akpınar H, Assimos DG. Kidney stones: a global picture of prevalence, incidence, and associated risk factors. *Reviews in urology*. 2010;12:e86–96.
- [6] Moore CL, Bomann S, Daniels B, Luty S, Molinaro A, Singh D, et al. Derivation and validation of a clinical prediction rule for uncomplicated ureteral stone—the STONE score: retrospective and prospective observational cohort studies. *Bmj*. 2014;348:g2191.
- [7] Inci MF, Ozkan F, Bozkurt S, Sucakli MH, Altunoluk B, Okumus M. Correlation of volume, position of stone, and hydronephrosis with microhematuria in patients with solitary urolithiasis. *Medical science monitor: international medical journal of experimental and clinical research*. 2013;19:295–9.
- [8] Argyropoulos A, Farmakis A, Doulas K, Lykourinas M. The presence of microscopic hematuria detected by urine dipstick test in the evaluation of patients with renal colic. *Urological research*. 2004;32:294–7.
- [9] Ooi SB, Kour NW, Mahadev A. Haematuria in the diagnosis of urinary calculi. *Annals of the Academy of Medicine, Singapore*. 1998;27:210–4.
- [10] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6:e1000097.
- [11] Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in medicine*. 2002;21:1539–58.
- [12] Harbord RM, Egger M, Sterne JA. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Statistics in medicine*. 2006;25:3443–57.
- [13] Kim TH, Oh SH, Park KN, Kim HJ, Youn CS, Kim SH, et al. Factors associated with absent microhematuria in symptomatic urinary stone patients. *The American journal of emergency medicine*. 2018.
- [14] Desai V, Cox M, Deshmukh S, Roth CG. Contrast-enhanced or noncontrast CT for renal colic: utilizing urinalysis and patient history of urolithiasis to decide. *Emergency radiology*. 2018;25:455–60.
- [15] Turk H, Un S. Predictive factors for stone disease in patients with renal colic. *Archivio italiano di urologia, andrologia: organo ufficiale [di] Societa italiana di ecografia urologica e nefrologica*. 2017;89:143–5.

- [16] Shrestha R, Bista Y, Khan A. Current Diagnostic Approach and Initial Treatment Patterns for Renal Colic in Emergency Department. *Journal of Nepal Health Research Council*. 2017;15:38-43.
- [17] Odoemene CA, Okere P, Ugonabo MC. Ureterolithiasis: Management in an environment with limited facilities. *Nigerian journal of clinical practice*. 2017;20:622-8.
- [18] Mefford JM, Tungate RM, Amini L, Suh D, Anderson CL, Rudkin SE, et al. A Comparison of Urolithiasis in the Presence and Absence of Microscopic Hematuria in the Emergency Department. *The western journal of emergency medicine*. 2017;18:775-9.
- [19] Rapp DE, Wood NL, Bassignani M, Gergoudis L, Caulkins S, Kramolowsky EV. Clinical variables and stone detection in patients with flank pain. *The Canadian journal of urology*. 2016;23:8441-5.
- [20] Park YH, Jung RB, Lee YG, Hong CK, Ahn JH, Shin TY, et al. Does the use of bedside ultrasonography reduce emergency department length of stay for patients with renal colic?: a pilot study. *Clinical and experimental emergency medicine*. 2016;3:197-203.
- [21] Hernandez N, Song Y, Noble VE, Eisner BH. Predicting ureteral stones in emergency department patients with flank pain: an external validation of the STONE score. *World journal of urology*. 2016;34:1443-6.
- [22] Fukuhara H, Ichiyanagi O, Kakizaki H, Naito S, Tsuchiya N. Clinical relevance of seasonal changes in the prevalence of ureterolithiasis in the diagnosis of renal colic. *Urolithiasis*. 2016;44:529-37.
- [23] Dorfman M, Chan SB, Hayek K, Hill C. Pyuria and Urine Cultures in Patients with Acute Renal Colic. *The Journal of emergency medicine*. 2016;51:358-64.
- [24] Yan JW, McLeod SL, Edmonds ML, Sedran RJ, Theakston KD. Risk Factors Associated with Urologic Intervention in Emergency Department Patients with Suspected Renal Colic. *The Journal of emergency medicine*. 2015;49:130-5.
- [25] Lee DH, Chang IH, Kim JW, Chi BH, Park SB. Usefulness of Nonenhanced Computed Tomography for Diagnosing Urolithiasis without Pyuria in the Emergency Department. *Biomed Res Int*. 2015;2015:810971.
- [26] Hall TC, Stephenson JA, Rangaraj A, Mulcahy K, Rajesh A. Imaging protocol for suspected ureteric calculi in patients presenting to the emergency department. *Clinical radiology*. 2015;70:243-7.

- [27] Zwank MD, Ho BM, Gresback D, Stuck LH, Salzman JG, Woster WR. Does computed tomographic scan affect diagnosis and management of patients with suspected renal colic? *The American journal of emergency medicine*. 2014;32:367-70.
- [28] Abdel-Gawad M, Kadasne R, Anjekar C, Elsobky E. Value of Color Doppler ultrasound, kub and urinalysis in diagnosis of renal colic due to ureteral stones. *International braz j urol: official journal of the Brazilian Society of Urology*. 2014;40:513-9.
- [29] Lallas CD, Liu XS, Chiura AN, Das AK, Bagley DH. Urolithiasis location and size and the association with microhematuria and stone-related symptoms. *Journal of endourology*. 2011;25:1909-13.
- [30] Perez JA, Palmes Mde L, Ferrer JF, Urdangarain OO, Nunez AB. Renal colic at emergency departments. Epidemiologic, diagnostic and etiopathogenic study. *Archivos espanoles de urologia*. 2010;63:173-87.
- [31] Xafis K, Thalmann G, Benneker LM, Stoupis C, Buggy DJ, Zimmermann H, et al. Forget the blood, not the stone! Microhaematuria in acute urolithiasis and the role of early CT scanning. *Emergency medicine journal: EMJ*. 2008;25:640-4.
- [32] Serinken M, Karcioğlu O, Turkcuer I, Ozkan HI, Keysan MK, Bukiran A. Analysis of clinical and demographic characteristics of patients presenting with renal colic in the emergency department. *BMC research notes*. 2008;1:79.
- [33] Cupisti A, Pasquali E, Lusso S, Carlino F, Orsitto E, Melandri R. Renal colic in Pisa emergency department: epidemiology, diagnostics and treatment patterns. *Internal and Emergency Medicine*. 2008;3:241-4.
- [34] Matani YS, Al-Ghazo MA. Role of helical nonenhanced computed tomography in the evaluation of acute flank pain. *Asian journal of surgery*. 2007;30:45-51.
- [35] Kartal M, Eray O, Erdogru T, Yilmaz S. Prospective validation of a current algorithm including bedside US performed by emergency physicians for patients with acute flank pain suspected for renal colic. *Emergency medicine journal: EMJ*. 2006;23:341-4.
- [36] Kirpalani A, Khalili K, Lee S, Haider MA. Renal colic: comparison of use and outcomes of

- unenanced helical CT for emergency investigation in 1998 and 2002. *Radiology*. 2005;236:554-8.
- [37] Gaspari R], Horst K. Emergency ultrasound and urinalysis in the evaluation of flank pain. *Academic emergency medicine: official journal of the Society for Academic Emergency Medicine*. 2005;12:1180-4.
- [38] Unal D, Yeni E, Karaoglanoglu M, Verit A, Karatas OF. Can conventional examinations contribute to the diagnostic power of unenhanced helical computed tomography in urolithiasis? *Urologia internationalis*. 2003;70:31-5.
- [39] Tack D, Sourtzis S, Delpierre I, de Maertelaer V, Gevenois PA. Low-dose unenhanced multidetector CT of patients with suspected renal colic. *AJR American journal of roentgenology*. 2003;180:305-11.
- [40] Kobayashi T, Nishizawa K, Mitsumori K, Ogura K. Impact of date of onset on the absence of hematuria in patients with acute renal colic. *The Journal of urology*. 2003;170:1093-6.
- [41] Eray O, Cubuk MS, Oktay C, Yilmaz S, Cete Y, Ersoy FF. The efficacy of urinalysis, plain films, and spiral CT in ED patients with suspected renal colic. *The American journal of emergency medicine*. 2003;21:152-4.
- [42] Luchs JS, Katz DS, Lane MJ, Mellinger BC, Lumerman JH, Stillman CA, et al. Utility of hematuria testing in patients with suspected renal colic: correlation with unenhanced helical CT results. *Urology*. 2002;59:839-42.
- [43] Hamm M, Knopfle E, Wartenberg S, Wawroschek F, Weckermann D, Harzmann R. Low dose unenhanced helical computerized tomography for the evaluation of acute flank pain. *The Journal of urology*. 2002;167:1687-91.
- [44] Li J, Kennedy D, Levine M, Kumar A, Mullen J. Absent hematuria and expensive computerized tomography: case characteristics of emergency urolithiasis. *The Journal of urology*. 2001;165:782-4.
- [45] Hamm M, Wawroschek F, Weckermann D, Knopfle E, Hackel T, Hauser H, et al. Unenhanced helical computed tomography in the evaluation of acute flank pain. *European urology*. 2001;39:460-5.
- [46] Richards JR, Christman CA. Intravenous urography in the emergency department: when do we need it? *European journal of emergency medicine: official journal of the European Society for*

Emergency Medicine. 1999;6:129-33.

[47] Bove P, Kaplan D, Dalrymple N, Rosenfield AT, Verga M, Anderson K, et al. Reexamining the value of hematuria testing in patients with acute flank pain. *The Journal of urology*. 1999;162:685-7.

[48] Ghali AM, Elmalik EM, Ibrahim AI, Abdulhameed E, el Tahir MI. Cost-effective emergency diagnosis plan for urinary stone patients presenting with ureteric colic. *European urology*. 1998;33:529-37.

[49] Eskelinen M, Ikonen J, Lipponen P. Usefulness of history-taking, physical examination and diagnostic scoring in acute renal colic. *European urology*. 1998;34:467-73.

[50] Gimondo P, Mastropasqua G, Cremona A, Mucciaccio C. [Early and systematic use of ultrasonography in emergency patients with renal colic: analysis of the actual diagnostic efficacy]. *La Radiologia medica*. 1996;92:421-4.

[51] Boyd R, Gray AJ. Role of the plain radiograph and urinalysis in acute ureteric colic. *Journal of accident & emergency medicine*. 1996;13:390-1.

[52] Press SM, Smith AD. Incidence of negative hematuria in patients with acute urinary lithiasis presenting to the emergency room with flank pain. *Urology*. 1995;45:753-7.

[53] Chia SJ, Lau W, Tan PK, Consigliere D, Li MK, Low CH. Ureteric colic: value of initial investigations and the outcome. *Annals of the Academy of Medicine, Singapore*. 1995;24:366-9.

[54] Elton TJ, Roth CS, Berquist TH, Silverstein MD. A clinical prediction rule for the diagnosis of ureteral calculi in emergency departments. *J Gen Intern Med*. 1993;8:57-62.

[55] Stewart DP, Kowalski R, Wong P, Krome R. Microscopic hematuria and calculus-related ureteral obstruction. *The Journal of emergency medicine*. 1990;8:693-5.

[56] Freeland P. No haematuria—no IVU. *Irish journal of medical science*. 1987;156:270-1.

[57] Dunn PM, Keller RT, Jones SR. The absence of hematuria in patients with symptomatic urinary tract stones. *The Western journal of medicine*. 1985;142:717-9.

[58] Bishop NL. The influence of emergency urography and haematuria on the diagnosis of ureteric colic. *Clinical radiology*. 1980;31:605-10.

[59] Bataille A, Wetzstein M, Hertig A, Vimont S, Rondeau E, Galichon P. Evidence of dipstick

superiority over urine microscopy analysis for detection of hematuria. BMC research notes.

2016;9:435.

[60] Daniels B, Gross CP, Molinaro A, Singh D, Luty S, Jessey R, et al. STONE PLUS: Evaluation of Emergency Department Patients With Suspected Renal Colic, Using a Clinical Prediction Tool Combined With Point-of-Care Limited Ultrasonography. *Annals of emergency medicine*. 2016;67:439-48.

[61] Rucker CM, Menias CO, Bhalla S. Mimics of renal colic: alternative diagnoses at unenhanced helical CT. *Radiographics: a review publication of the Radiological Society of North America, Inc*. 2004;24 Suppl 1:S11-28; discussion S-33.

[62] Tao RZ, Qin ZQ, Liu FD, Lv JL. Efficacy and Safety of Tamsulosin in the Medical Expulsion Therapy for Distal Ureteral Calculi: A Systematic Review and Meta-Analysis of Placebo-Controlled Trials. *Urol J*. 2019;16:224-31.

[63] Ten Things Physicians and Patients Should Question American College of Emergency Physicians 2013-2014.

[64] Catalano O, Nunziata A, Sandomenico F, Siani A. Acute flank pain: comparison of unenhanced helical CT and ultrasonography in detecting causes other than ureterolithiasis. *Emergency radiology*. 2002;9:146-54.

Tables

Table 1: Basic study and patient characteristics. Patients presenting with acute renal colic at the emergency department.

Authors	Year	Country	Study design	No. of patients	% Male	Mean age \pm SD (years)
Kim <i>et al.</i> [13]	2018	South Korea	Retrospective, observational	798	68.6	48.2 \pm 13.3
Desai <i>et al.</i> [14]	2018	USA	Retrospective, observational	350	NR	NR
Türk and Ün [15]*	2017	Turkey	Prospective, observational	516	60.5	37 \pm 20.3
Shrestha <i>et al.</i> [16]*	2017	Nepal	Retrospective, observational	201	55.2	29 \pm 13.5
Odoemene <i>et al.</i> [17]*	2017	Nigeria	Prospective, observational	69	76.8	40.4 \pm 2.9
Mefford <i>et al.</i> [18]	2017	USA	Retrospective, observational	393	69	Median 43 (IQR 32-54)
Rapp <i>et al.</i> [19]	2016	USA	Retrospective, observational	613	47	49 \pm 0.6

Park <i>et al.</i> [20]	2016	South Korea	Prospective, RCT	103	66	45.6 ± 12.55
Hernandez <i>et al.</i> [21]	2016	USA	Retrospective, observational	536	56	45.9 ± 16.3
Fukuhara <i>et al.</i> [22]*	2016	Japan	Retrospective, observational	491	70.5	51.8 ± 15
Dorfman <i>et al.</i> [23]	2016	USA	Retrospective, observational	339	55.5	46.8 ± 16.5
Yan <i>et al.</i> [24]	2015	Canada	Prospective cohort study	565	62.8	46.6 ± 14.4
Lee <i>et al.</i> [25]	2015	South Korea	Retrospective, observational	2,218	71	43.3 ± 14.2
Hall <i>et al.</i> [26]*	2015	UK	Retrospective, observational	513	57.1	45 ± 23.3
Zwank <i>et al.</i> [27]	2014	USA	Prospective, observational	93	NR	39 ± NR
Abdel-Gawad <i>et al.</i> [28]*	2014	UAE	Retrospective, observational	939	87.9	37.9 ± 11
Inci <i>et al.</i> [7]	2013	Turkey	Retrospective, observational	83	42.2	42.1 ± 14.4
Lallas <i>et al.</i> [29]	2011	USA	Prospective, observational	32	NR	NR
Perez <i>et al.</i> [30]*	2010	Spain	Prospective, multicentre, cross-sectional case-control	146	57.53	51.34 ± NR
Xafis <i>et al.</i> [31]*	2008	Switzerland	Retrospective, observational	638	NR	44.3 ± 14.6
Serinken <i>et al.</i> [32]*	2008	Turkey	Retrospective, observational	235	75.7	31.1 ± 7
Čupisti <i>et al.</i> [33]	2008	Italy	Retrospective, observational	696	54	NR
Matani and Al-Ghazo [34]*	2007	Saudi Arabia / Jordan	Retrospective, observational	75	61.3	42.2 ± NR
Kartal <i>et al.</i> [35]*	2006	Turkey	Prospective, observational	227	64.8	38.4 ± 14
Kirpalani <i>et al.</i> [36]	2005	Canada	Retrospective, observational	299	NR	NR
Gaspari and Horst [37]	2005	USA	Prospective, observational	110	NR	NR
Argyropoulos <i>et al.</i> [8]	2004	Greece	Retrospective, observational	609	63.2	49.2 ± 15.9
Unal <i>et al.</i> [38]*	2003	Turkey	Prospective, observational	137	55	38 ± NR
Tack <i>et al.</i> [39]*	2003	Belgium	Prospective, observational	106	50	45 ± NR
Kobayashi <i>et al.</i> [40]	2003	Japan	Retrospective, observational	537	78	46.6 ± 14
Eray <i>et al.</i> [41]	2003	Turkey	Prospective, observational	65	60	38.8 ± 13.5
Lucks <i>et al.</i> [42]	2002	USA	Retrospective, observational	587	NR	NR
Hamm <i>et al.</i> [43]	2002	Germany	Prospective, observational	109	69.7	49 ± NR
Li <i>et al.</i> [44]*	2001	USA	Retrospective, observational	397	73	47 ± 15
Hamm <i>et al.</i> [45]	2001	Germany	Prospective, observational	125	72	55 ± 17
Richards and Christman [46]	1999	USA	Retrospective, observational	185	NR	NR
Bove <i>et al.</i> [47]	1999	USA	Retrospective, observational	195	NR	NR
Ooi <i>et al.</i> [9]*	1998	Singapore	Prospective, observational	122	93	39.7 ± NR
Ghali <i>et al.</i> [48]*	1998	Saudi Arabia	Prospective, observational	125	80	39.2 ± NR
Eskelinen <i>et al.</i> [49]	1998	Finland	Prospective,	57	NR	NR

Gimondo <i>et al.</i> [50]*	1996	Italy	observational Retrospective,	76	60.5	47 ± NR
Boyd and Gray [51]	1996	UK	observational Prospective,	52	NR	NR
Press and Smith [52]	1995	USA	observational Retrospective,	109	NR	NR
Chia <i>et al.</i> [53]	1995	Singapore	observational Prospective,	294	72.5	43.5 ± NR
Elton <i>et al.</i> [54]*	1993	USA	observational Retrospective / prospective,	275	71.2	46.2 ± 15.7
Stewart <i>et al.</i> [55]	1990	USA	observational Retrospective,	160	76.9	NR
Freeland [56]	1987	Northern Ireland	observational Retrospective,	134	NR	NR
Dunn <i>et al.</i> [57]	1985	USA	observational Retrospective,	76	NR	42.7 ± NR
Bishop [58]	1980	UK	observational Prospective,	50	NR	NR

Abbreviations (alphabetical order): IQR = interquartile range; NR = not reported; RCT = Randomized control study; SD = standard deviation; UAE = United Arab Emirates; UK = United Kingdom; USA = United States of America.

Note: *Enrolled also children.

Table 2: Data on microhematuria in patients presenting with suspected acute renal colic at the emergency department.

Authors	Microhematuria test	Type of hematuria	Positive microhematuria definition
Kim <i>et al.</i> [13]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Desai <i>et al.</i> [14]	Urinalysis	Microscopic or macroscopic	Positive urinalysis for RBCs or for k
Türk and Ün [15]	Urinalysis	Microscopic	NR
Shrestha <i>et al.</i> [16]	Urinalysis	Microscopic	Presence of 3 or more RBCs
Odoemene <i>et al.</i> [17]	Urinalysis	Microscopic or macroscopic	NR
Mefford <i>et al.</i> [18]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Rapp <i>et al.</i> [19]	Urinalysis	Microscopic or macroscopic	Presence of 4 or more RBCs/HPF
Park <i>et al.</i> [20]	Urinalysis	Microscopic	NR

Hernandez <i>et al.</i> [21]	Urine dipstick	Microscopic	Hematuria on urine dipstick
Fukuhara <i>et al.</i> [22]	Urinalysis or urine dipstick	Microscopic or macroscopic	Occult blood in urine
Dorfman <i>et al.</i> [23]	Urinalysis	Microscopic	Presence of 5 or more RBCs/HPF
Yan <i>et al.</i> [24]	Urinalysis	Microscopic	NR
Lee <i>et al.</i> [25]	Urinalysis	Microscopic	NR
Hall <i>et al.</i> [26]	Urine dipstick	Microscopic or macroscopic	Scores of 1+ to 3+ on urine dipstick hematuria
Zwank <i>et al.</i> [27]	Urinalysis	Microscopic	RBCs present
Abdel-Gawad <i>et al.</i> [28]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Inci <i>et al.</i> [7]	Urinalysis	Microscopic	Presence of 5 or more RBCs/HPF
Lallas <i>et al.</i> [29]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
	Urine dipstick	Microscopic	Trace or scores of 1+ to 4+ on urine dipstick
Perez <i>et al.</i> [30]	Urine dipstick	Microscopic	NR
Xafis <i>et al.</i> [31]	Urinalysis	Microscopic	Presence of 5 or more RBCs/HPF
Serinken <i>et al.</i> [32]	Urinalysis	Microscopic	Presence of 5 or more RBCs/HPF
Cupisti <i>et al.</i> [33]	Urine dipstick	Microscopic	NR
Matani and Al-Ghazo [34]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Kartal <i>et al.</i> [35]	Urinalysis	Microscopic	Presence of 10 or more RBCs/HPF
Kirpalani <i>et al.</i> [36]	Urine dipstick	Microscopic	Positive urine dipstick
Gaspari and Horst [37]	Urinalysis	Microscopic	Presence of 5 or more RBCs/HPF
Argyropoulos <i>et al.</i> [8]	Urine dipstick	Microscopic	Scores of 1+ to 3+ on urine dipstick hematuria
Unal <i>et al.</i> [38]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF

Tack <i>et al.</i> [39]	Urinalysis or Urine dipstick	Microscopic	Presence of 2 or more RBCs/HPF o
Kobayashi <i>et al.</i> [40]	Urine dipstick	Microscopic	Scores of 1+ to 3+ on urine dipstick
	Urinalysis	Microscopic	Presence of 5 or more RBCs/HPF
Eray <i>et al.</i> [41]	Urinalysis	Microscopic	Presence of 6 or more RBCs/HPF
Luchs <i>et al.</i> [42]	Urinalysis	Microscopic	Presence of 10 or more RBCs/HPF
Hamm <i>et al.</i> [45]	Urinalysis	Microscopic	Presence of more than 20 mg/dl h
Li <i>et al.</i> [44]	Urinalysis or Urine dipstick	Microscopic	Presence of any number of RBCs/H on urine dipstick
Hamm <i>et al.</i> [45]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Richards and Christman [46]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Bove <i>et al.</i> [47]	Urine dipstick	Microscopic	Positive urine dipstick
	Urinalysis	Microscopic	Presence of 6 or more RBCs/HPF
	Urinalysis or Urine dipstick	Microscopic	Presence of 2 or more RBCs/HPF c
Ooi <i>et al.</i> [9]	Urine dipstick	Microscopic	Scores of 1+ or more on urine dips
	Urinalysis	Microscopic	Presence of 6 or more RBCs/HPF ir RBCs/HPF in females
Ghali <i>et al.</i> [48]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Eskelinen <i>et al.</i> [49]	Urinalysis	Microscopic	Presence of 11 or more RBCs/HPF
Gimondo <i>et al.</i> [50]	Urine dipstick	Microscopic or macroscopic	Positive urine dipstick
Boyd and Gray [51]	Urine dipstick	Microscopic	Positive urine dipstick
Press and Smith [52]	Urinalysis	Microscopic	Presence of 1 or more RBCs/HPF
Chia <i>et al.</i> [53]	Urinalysis	Microscopic	Presence of 6 or more RBCs/HPF ir RBCs/HPF in females

Elton <i>et al.</i> [54]	Urinalysis	Microscopic	Presence of 4 or more RBCs/HPF
Stewart <i>et al.</i> [55]	Urinalysis	Microscopic	Presence of 3 or more RBCs/HPF
Freeland [56]	Urine dipstick	Microscopic	Trace or scores of 1+ to 3+ on urinalysis
Dunn <i>et al.</i> [57]	Urinalysis	Microscopic	Presence of 3 or more RBCs/HPF
Bishop [58]	Urine dipstick	Microscopic	Positive urine dipstick

Abbreviations (alphabetical order): NR = not reported; HPF = High power Field; RBC = Red Blood Cell.

Table 3: Data on microhematuria in patients presenting with confirmed urolithiasis at the emergency department.

Authors	Microhematuria test	No. patients with microhematuria	Microhematuria prevalence	Diagnostic modality
Kim <i>et al.</i> [13]	Urinalysis	750	750/798 (94%)	Unenhanced CT
Desai <i>et al.</i> [14]	Urinalysis	231	231/282 (81.9%)	Non-contrast CT
Türk <i>et al.</i> [15]	Urinalysis	344	344/388 (88.7%)	Non-contrast CT
Shrestha <i>et al.</i> [16]	Urinalysis	27	27/61 (44.3%)	Renal US
Odoemene <i>et al.</i> [17]*	Urinalysis	62	62/69 (89.9%)	Abdominal US
Mefford <i>et al.</i> [18]	Urinalysis	321	321/393 (81.7%)	Non-contrast CT
Rapp <i>et al.</i> [19]*	Urinalysis	177	177/222 (79.7%)	Non-contrast CT
Fukuhara <i>et al.</i> [22]*	Urinalysis or urine dipstick	323	323/358 (90.2%)	Plain abdominal X-ray, non-contrast CT
Dorfman <i>et al.</i> [23]	Urinalysis	254	245/339 (74.9%)	Abdominal X-ray
Hall <i>et al.</i> [26]*	Urine dipstick	193	193/233 (82.8%)	Non-enhanced CT
Zwank <i>et al.</i> [27]	Urinalysis	52	52/62 (83.9%)	CT
Abdel-Gawad <i>et al.</i> [28]	Urinalysis	835	835/939 (88.9%)	Color Doppler US

Inci <i>et al.</i> [7]	Urinalysis	46	46/83 (55.4)	helical CT Unenhanc
Lallas <i>et al.</i> [29]	Urinalysis	18	18/32 (56.3)	US, Abdom
	Urine dipstick	21	21/32 (65.6)	
Xafis <i>et al.</i> [31]	Urinalysis	341	341/507 (67.3)	Unenhanc
Kartal <i>et al.</i> [35]	Urinalysis	121	121/176 (68.8)	IVU, US, s
Gaspari and Horst [37]	Urinalysis	54	54/58 (93.1)	US, CT
Argyropoulos <i>et al.</i> [8]	Urine dipstick	539	539/564 (95.6)	Abdomen
Unal <i>et al.</i> [38]	Urinalysis	92	92/114 (80.7)	US, excret
Tack <i>et al.</i> [39]	Urinalysis or Urine dipstick	37	37/38 (97.4)	Excretory
Kobayashi <i>et al.</i> [40]	Urine dipstick	346	346/452 (76.5)	Abdomen
	Urinalysis	317	317/452 (70.1)	
Eray <i>et al.</i> [41]	Urinalysis	37	37/54 (68.5)	Abdomen
Luchs <i>et al.</i> 42[]	Urinalysis	492	492/587 (83.8)	CT, stone
Hamm <i>et al.</i> [43]	Urinalysis	53	53/80 (66.3)	Unenhanc
Li <i>et al.</i> [44]	Urinalysis or Urine dipstick	360	360/397 (90.7)	CT, IVP
Hamm <i>et al.</i> [45]	Urinalysis	76	76/91 (83.5)	Helical CT
Richards and Christman [46]	Urinalysis	88	88/98 (89.8)	IVU
Bove <i>et al.</i> [47]	Urine dipstick	70	70/87 (80.5)	CT
	Urinalysis	77	77/95 (81.1)	
	Urinalysis or Urine dipstick	82	82/95 (86.3)	
Ooi <i>et al.</i> [9]	Urine dipstick	62	62/65 (95.4)	Abdomen

	Urinalysis	46	46/65 (70.8)	
Ghali <i>et al.</i> [48]	Urinalysis	64	64/82 (78)	Abdomen
Gimondo <i>et al.</i> [50]*	Urine dipstick	29	29/29 (100)	US
Boyd and Gray [51]	Urine dipstick	29	29/29 (100)	Abdomen
Press and Smith [52]	Urinalysis	78	78/109 (71.6)	IVU
Stewart <i>et al.</i> [55]	Urinalysis	132	132/160 (82.5)	IVP
Freeland [56]	Urine dipstick	72	72/76 (94.7)	IVU or sto
Dunn <i>et al.</i> [57]	Urinalysis	62	62/76 (81.6)	IVU or sto
Bishop [58]	Urine dipstick	33	33/35 (94.3)	IVU

Abbreviations (alphabetical order): CT = computed tomography; HFU = High-power field; IVU = Intravenous Urography; MDCT = multidetector CT; NR = not reported; RBC = Red Blood Cell; SD = standard deviation; US = ultrasound

* This study included also patients with gross hematuria

Figures




Figure 1

PRISMA flow chart of the retrieved, excluded and analyzed studies.




Figure 2

Plots of individual studies and pooled prevalence of microhematuria in patients with acute renal colic, including 95% confidence intervals (95%CI).




Figure 3

Plots of individual studies and pooled prevalence of microhematuria in patients with confirmed urolithiasis, including 95% confidence intervals (95%CI).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

[Appendix 1.docx](#)

[Supplemental Figure 1.tif](#)