

Table. 1 Primer sequences for RT-qPCR

Gene	Forward primer (5'–3')	Reverse primer (5'–3')
<i>IL-17A</i>	GGCTGGAGAAGATACTGGTGTC	AGGCTGTCTTTGAAGGATGAGG
<i>RORγt</i>	ATGCTTTGGCGATGAGTCTTG	CGAGGATGAGATTGCCCTCTA
<i>IL-21</i>	CCTGGCAACATGGAGAGGATTGTC	TGCTGACTTTAGTTGGGCCTTCTG
<i>IL-6R</i>	CCTGACGACAAAGGCTGTGCTCT	GCTGAACTTGCTCCCGACACTACTG
<i>GAPDH</i>	GGCACAGTCAAGGCTGAGAATG	ATGGTCATGCAAGACGCCAGTA
<i>hsa-piR-15254</i>	GGTGCTGTGTAGTGCGCTATG	CAGTTGGTCAGCAGTTCTTGGTAT
<i>hsa-piR-5660</i>	TCACATTGCCAGGGATTGTCG	GGTCCGAGGTATTTCGCACT
<i>hsa-piR-6426</i>	TCGTATCCAGTGCGAATACCT	TCGTATCCAGTGCAGGGTCC
<i>U6</i>	CGCTTCGGCAGCACATATAC	TTCACGAATTTGCGTGTGCATC

IL-17A interleukin 17A, *ROR γ t* retinoic acid receptor-related orphan receptors γ t, *IL-21* interleukin 21, *IL-6R* interleukin 6 receptor, *GAPDH* glyceraldehyde-3-phosphate dehydrogenase, *RT-qPCR* reverse transcription-quantitative polymerase chain reaction