

# Homologies between SARS-CoV-2 and allergen proteins may direct T cell-mediated heterologous immune responses

## Kathrin Balz

Institute of Laboratory Medicine, Universities of Giessen and Marburg Lung Center (UGMLC), Philipps University Marburg, German Center for Lung Research (DZL), Marburg, Germany

## Meng Chen

Sean N. Parker Center for Allergy and Asthma Research at Stanford University and Division of Pulmonary, Allergy & Critical Care Medicine, Stanford, CA, USA

## Abhinav Kaushik

Sean N. Parker Center for Allergy and Asthma Research at Stanford University and Division of Pulmonary, Allergy & Critical Care Medicine, Stanford, CA, USA

## Franz Cemic

TH Mittelhessen, Department of Computer Science, University of Applied Sciences Gießen, Hessen, Deutschland

## Vanessa Heger

TH Mittelhessen, Department of Computer Science, University of Applied Sciences Gießen, Hessen, Deutschland

## Harald Renz

Institute of Laboratory Medicine, Universities of Giessen and Marburg Lung Center (UGMLC), Philipps University Marburg, German Center for Lung Research (DZL), Marburg, Germany

## Kari Nadeau

Sean N. Parker Center for Allergy and Asthma Research at Stanford University and Division of Pulmonary, Allergy & Critical Care Medicine, Stanford, CA, USA

## Chrysanthi Skevaki (✉ [skevakioffice@gmail.com](mailto:skevakioffice@gmail.com))

Institute of Laboratory Medicine, Universities of Giessen and Marburg Lung Center (UGMLC), Philipps University Marburg, German Center for Lung Research (DZL), Marburg, Germany <https://orcid.org/0000-0001-5194-5635>

---

## Research Article

**Keywords:** bioinformatics, COVID-19, allergy, T cell epitope, heterologous immunity, cross-reactivity

**Posted Date:** October 6th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-86873/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

1 **Homologies between SARS-CoV-2 and allergen proteins may direct T cell-**  
2 **mediated heterologous immune responses**

3

4 Kathrin Balz<sup>1\*</sup>, MSc, Meng Chen<sup>2\*</sup>, MD, Abhinav Kaushik<sup>2\*</sup>, PhD, Franz Cemic<sup>3</sup>, PhD,  
5 Vanessa Heger<sup>3</sup>, BSc, Harald Renz<sup>1</sup>, MD, Kari Nadeau<sup>2#</sup>, MD, PhD, Chrysanthi  
6 Skevaki<sup>1#</sup>, MD

7

8 <sup>1</sup> Institute of Laboratory Medicine, Universities of Giessen and Marburg Lung Center  
9 (UGMLC), Philipps University Marburg, German Center for Lung Research (DZL),  
10 Marburg, Germany

11

12 <sup>2</sup> Sean N. Parker Center for Allergy and Asthma Research at Stanford University and  
13 Division of Pulmonary, Allergy & Critical Care Medicine, Stanford, CA, USA.

14

15 <sup>3</sup> TH Mittelhessen, Department of Computer Science, University of Applied Sciences  
16 Gießen, Hessen, Deutschland

17 \* equal contribution

18 # equal contribution

19

20 **MATERIALS AND CORRESPONDENCE**

21 Chrysanthi Skevaki, PD Dr.

22 Institute of Laboratory Medicine, Philipps University Marburg

23 Baldingerstrasse, 35043 Marburg, Germany

24 Phone +49 6421 586 6235

25 Email: chrysanthi.skevaki@uk-gm.de

26

27 **ACKNOWLEDGEMENTS**

28 CS is supported by the Universities Giessen and Marburg Lung Center (UGMLC), the

29 German Center for Lung Research (DZL), University Hospital Giessen and Marburg

30 (UKGM) research funding according to article 2, section 3 cooperation agreement,

31 and the Deutsche Forschungsgemeinschaft (DFG)-funded SFB 1021 (C04), KFO 309

32 (P10), and SK 317/1-1 (Project number 428518790) as well as by the Foundation for  
33 Pathobiochemistry and Molecular Diagnostics.

34 KCN is supported by NIH grants U01AI140498-03 and R01AI140134-01, The  
35 Sunshine Foundation, and the Sean N. Paker Center for Allergy and Asthma Research  
36 at Stanford University.

37

### 38 **COMPETING INTERESTS**

39 For CS: Consultancy and research funding, Hycor Biomedical, Bencard Allergie and  
40 Thermo Fisher Scientific; Research Funding, Mead Johnson Nutrition (MJN).

41 For KCN: Dr. Nadeau reports grants from National Institute of Allergy and Infectious  
42 Diseases (NIAID), Food Allergy Research & Education (FARE), End Allergies Together  
43 (EAT), Allergenics, and Ukko Pharma; Grant awardee at NIAID, National Institute of  
44 Environmental Health Sciences (NIEHS), National Heart, Lung, and Blood Institute  
45 (NHLBI), and the Environmental Protection Agency (EPA); is involved in Clinical trials  
46 with Regeneron, Genentech, Alimmune Therapeutics, DBV Technologies, AnaptysBio,  
47 Adare Pharmaceuticals, and Stallergenes-Greer; Research Sponsorship by Novartis,  
48 Sanofi, Astellas, Nestle; Data and Safety Monitoring Board member at Novartis and  
49 NHLBI; Cofounded Before Brands, Alladapt, ForTra, and Iggenix; Chief Intellectual  
50 Office at FARE, Director of the World Allergy Organization (WAO) Center of Excellence  
51 at Stanford, Personal fees from Regeneron, Astrazeneca, ImmuneWorks, and Cour  
52 Pharmaceuticals; Consultant and Advisory Board Member at European Academy of  
53 Allergy and Clinical Immunology (EAACI) Research and Outreach Committee, Ukko,  
54 Before Brands, Alladapt, IgGenix, Probio, Vedanta, Centecor, Seed, Novartis, NHBLI,  
55 EPA, National Scientific Committee of Immune Tolerance Network (ITN) and NIH  
56 Programs; US patents for basophil testing, multifood immunotherapy and prevention,  
57 monoclonal antibody from plasmoblasts, and device for diagnostics.

58 All other authors have no competing interests to declare.

59

## 60 **AUTHOR CONTRIBUTIONS**

61 Conception or design of the work: CS, KCN; Data collection: KB, AK, FC, VH; Data  
62 analysis and interpretation: CS, KCN, KB, AK; Drafting the article: CS, KCN, KB, MC,  
63 AK; Critical revision of the article: CS, KCN, KB, MC, AK, HR; Final approval of the  
64 version to be published: All

65

## 66 **LIST OF ABBREVIATIONS**

67

68	ACE2	Angiotensin Converting Enzyme 2
69	ARDS	Acute Respiratory Distress Syndrome
70	BLAST	Basic Local Alignment Search Tool
71	HLA	Human Leukocyte Antigen
72	IEDB	Immune Epitope Database
73	MHC	Major Histocompatibility Complex
74	NCBI	National Center for Biotechnology Information
75	RNA	ribonucleic acid
76	+ssRNA	positive-sense single-stranded RNA
77	Th 1	t-helper 1
78	Th 2	t-helper 2

79

80 **ABSTRACT**

81 The outbreak of the new Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-  
82 CoV-2) is a public health emergency. Asthma does not represent a risk factor for  
83 COVID-19 in several published cohorts. We hypothesized that the SARS-CoV-2  
84 proteome contains T cell epitopes, which are potentially cross-reactive to allergen  
85 epitopes. We aimed at identifying homologous peptide sequences by means of two  
86 distinct complementary bioinformatics approaches. Pipeline 1 included prediction of  
87 MHC Class I and Class II epitopes contained in the SARS-CoV-2 proteome and  
88 allergens along with alignment and elaborate ranking approaches. Pipeline 2 involved  
89 alignment of SARS-CoV-2 overlapping peptides with known allergen-derived T cell  
90 epitopes. Our results indicate a large number of MHC Class I epitope pairs including  
91 known as well as *de novo* predicted allergen T cell epitopes with high probability for  
92 cross-reactivity. Allergen sources, such as *Aspergillus fumigatus*, *Phleum pratense*  
93 and *Dermatophagoides* species are of particular interest due to their association with  
94 multiple cross-reactive candidate peptides, independently of the applied bioinformatic  
95 approach. In contrast, peptides derived from food allergens, as well as MHC class II  
96 epitopes did not achieve high *in silico* ranking and were therefore not further  
97 investigated. Our findings warrant further experimental confirmation along with  
98 examination of the functional importance of such cross-reactive responses.

99

## 100 **Introduction**

101 The World Health Organization (WHO) has declared the outbreak of the new Severe  
102 Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2, ssRNA virus, associated  
103 with COVID-19) as a public health emergency. As per the WHO report of 20 September  
104 2020, more than 30 million cases and over 950000 deaths have been reported  
105 worldwide.<sup>1</sup> Human coronaviruses are positive-sense single-stranded RNA (+ssRNA)  
106 viruses, with SARS-CoV-2 and SARS-CoV belonging to the B-lineage of the  
107 Betacoronavirus genera and MERS-CoV to the C-lineage of the same genera.<sup>2,3</sup> The  
108 clinical features in patients affected with these respiratory viruses ranges from  
109 asymptomatic carriers to severe respiratory illness with pneumonia and acute  
110 respiratory distress syndrome (ARDS). In addition, a number of interesting vascular  
111 and inflammatory presentations have been noted, including a multisystem  
112 inflammatory syndrome in children.

113 We have previously reported on heterologous immune responses induced by  
114 influenza, another respiratory RNA virus, against allergens, which mediated protection  
115 from experimental allergic asthma.<sup>4</sup> Indeed, virus-induced T cell mediated  
116 heterologous immunity has been widely described in a variety of settings, which can  
117 confer protection or drive immunopathology against other antigens.<sup>5,6</sup> Given that the  
118 host immune response to SARS-CoV-2 and associated disease course can be so  
119 varied from patient to patient, this spectrum of presentations raises the question of  
120 what drives the differential host immune response. There is still little known about  
121 asthma phenotypes and severity of COVID-19. In general, asthma has not been shown  
122 to be a risk factor for COVID-19 in several published cohorts.<sup>7,8</sup> However, recent  
123 studies from the UK and the USA indicated higher numbers of asthmatics in COVID-  
124 19 patients.<sup>9</sup>

125 Interestingly, the UK Biobank recently reported that non-allergic patients had a higher  
126 risk of severe COVID-19, compared to patients with allergic asthma<sup>10</sup>. Moreover,  
127 increased numbers of activated T cells were found among asthmatic COVID-19  
128 patients who showed a less severe disease, suggesting that activated T cells have a  
129 positive impact on severity of SARS-CoV-2 infection<sup>11</sup>. These preliminary clinical  
130 observations along with our prior experimental evidence involving RNA viruses led us  
131 hypothesize that SARS-CoV-2 may share a degree of protein sequence homology to  
132 allergens, which may lead to the generation of cross-reactive T cell epitopes. Pre-  
133 existing T cells specific for such cross-reactive allergen-derived epitopes may have an  
134 impact on COVID-19 outcome via aberrant cytokine responses to the virus peptides.  
135 Indeed, these cytokines could prevent an overshooting T1 inflammatory reaction, both  
136 locally (as in the case of preexisting pulmonary CD4<sup>+</sup> T cells specific to inhalant  
137 allergens) and/or systemically. Therefore, we sought to predict potentially cross-  
138 reactive allergen- and SARS-CoV-2-derived MHC Class I and Class II T cell epitopes,  
139 which can be presented by the most prevalent HLA alleles.

140

## 141 **Methods and Results**

142 In order to examine our working hypothesis, we applied two distinct independent,  
143 complementary and systematic bioinformatics approaches (**Figure 1**): a) Pipeline 1-  
144 prediction of MHC Class I and Class II epitopes contained in the SARS-CoV-2  
145 proteome and a comprehensive set of allergen protein sequences combined with  
146 alignment strategies and ranking of results based on clinical and sequence  
147 conservation criteria and b) Pipeline 2- alignment of SARS-CoV-2 overlapping peptides  
148 with known allergen-derived T cell epitopes.<sup>12</sup>

### 149 **Pipeline 1**



150 More than >2500 allergen protein sequences were downloaded (dates of access  
151 10.09.2017) from Allergome<sup>13-16</sup> (**Supplementary Table S1**), and protein sequences  
152 for SARS-CoV-2 from UniProt<sup>17</sup> (**Supplementary Table S2**). Viral T cell epitope  
153 prediction was performed using smm<sup>18</sup>, ann<sup>19</sup> and consensus<sup>20</sup> for MHC Class I (IC50  
154 threshold  $\leq 5000$ ), and netMHCII<sup>21</sup> for MHC Class II (affinity score threshold for strong  
155 binders: 0.500; for weak binders: 2.000) (**Supplementary Table S3**). Epitopes  
156 predicted by all methods were aligned against all allergen proteins with NCBI protein  
157 blast platform<sup>22</sup>. Allergen proteins associated with an alignment e-value  $< 10$  were  
158 further processed for T cell epitope prediction using netMHC<sup>23</sup> and netMHCpan<sup>24</sup> for  
159 MHC Class I, and netMHCII and netMHCIIpan<sup>25</sup> for MHC Class II prediction (affinity  
160 score threshold for strong binders: 0.500; for weak binders: 2.000). Viral and allergen  
161 epitopes were pairwise aligned with Biopython module pairwise 2<sup>26</sup> and for pairs with  
162 a score  $> 8$ , a final pair combined score (pcs) was calculated (**Supplementary**  
163 **Methods**). Duplicates among the resulting candidate epitope pairs were removed  
164 before further processing. Therefore, possible sequence repetition due to isoforms and  
165 isoallergens (**Supplementary Table S1**) do not influence further analyses. In total, we  
166 obtained more than 5000 candidate pairs for each, MHC Class I and Class II. The top  
167 30 candidate epitope pairs, as per pair combined score, are listed for aero- and food  
168 allergens, MHC Class I and Class II presentation background in **Supplementary**  
169 **Tables S4-7**, respectively. The top 30 MHC Class II restricted predicted virus-allergen  
170 pairs achieved relatively low pcs (24-657) as compared to Class I epitope pairs (1036-  
171 10816). Among our top 30 MHC Class I potentially cross-reactive allergen derived  
172 epitopes, we identified more than 20 distinct protein families (Allfam database). In  
173 addition to MHC binding affinity and homology between peptide sequences, also other  
174 factors (e.g. conservation, association with clinical reactions) are important for the  
175 clinical relevance of peptides predicted to be cross-reactive at the T cell level. In order

176 to capture this information level in our ranking, all allergen peptides and associated  
177 sources listed among the top 30 candidate epitope pairs were evaluated further with a  
178 scoring system (**Supplementary Fig. S1 and Supplementary Methods**). We found  
179 that the top 5 Class I aeroallergens were on average associated with higher pcs as  
180 compared to the top 5 potentially cross-reactive food allergens (**Table 1** for MHC Class  
181 I and **Table 2** for MHC Class II peptide pairs).

## 182 **Pipeline 2**

183 We obtained all known allergen-derived linear T cell epitope peptides from the IEDB,  
184 containing peptides known to bind MHC molecules with at least one published  
185 experimental evidence (e.g. based on the results of a T cell assay) (**Supplementary**  
186 **Table S8**). A total of 8,207 antigenic peptides from 142 antigens were selected for  
187 evaluation, among which, peptides with ambiguous amino acids (e.g. with unknown  
188 amino acid 'X' or any special character) were removed from the subsequent analysis.  
189 Therefore, all included peptides could be defined in full. Next, SARS-CoV-2 protein  
190 sequences were analyzed for the potential antigenic regions by splitting each of the  
191 sequence into sequential *k-mers* (length=15), and homology with allergen antigenic  
192 peptides was then profiled. Within a given threshold range, we found 43 unique SARS-  
193 CoV-2 peptides that belong to replicase poly protein and spike glycoprotein  
194 (**Supplementary Table S9**). These peptides demonstrate homology with antigenic  
195 peptides of 6 different allergens, all of which are known to be respiratory allergens (e.g.  
196 aeroallergens; **Figure 1**). However, despite the homology, it is likely that some of the  
197 peptides may not have strong MHC Class I binding affinity, and thus be less likely to  
198 be presented as antigens by HLA molecules. Therefore, we assessed the binding  
199 affinity of these peptides with human MHC Class I molecules, across a broad range of  
200 alleles that are known to bind viral proteins (52 most common HLA-A and HLA-B  
201 alleles). We observed that some of these peptides (n=79) were predicted to have MHC

202 Class I binding epitope regions associated with at least one of the Class I HLA alleles  
203 with IC50 < 500nm (**Supplementary Table S10**). These antigenic peptides were  
204 predicted to bind with 20 most frequently occurring HLA Class I alleles, in which  
205 HLA\*02:03 and HLA\*02:06 were predicted to present the highest number of epitope  
206 residues. To further investigate if these peptides are specific to the coronavirus family,  
207 we performed the BLAST comparison with 2807 known viral antigenic peptides of  
208 bacteria, influenza-and corona- virus family (non-SARS CoV-2) from IEDB (with at  
209 least one T cell assay evidence) and filtered out matching peptides (Blast e-value < 1  
210 & identity > 70%). Finally, we present 48 high-affinity HLA-binding peptides which are  
211 unique to the SARS-CoV-2 proteome, not common to bacteria, influenza and corona  
212 virus family antigenic peptides within a given threshold range (**Supplementary Table**  
213 **S11**) with 14 high confidence HLA Class I binding peptides with IC50 < 50nm (**Table**  
214 **3**).

215

## 216 **Discussion**

217 We have applied two independent, complementary and systematic bioinformatic  
218 approaches in order to identify potentially cross-reactive allergen- and SARS-CoV-2-  
219 T cell epitopes. Our *in silico* analysis revealed numerous candidate epitope pairs,  
220 including previously published and predicted peptides, while both applied pipelines  
221 highlighted an important role of MHC class I inhalant allergens. Although the frequency  
222 of allergen-specific CD8<sup>+</sup> T cells is likely to be low, rare cell subsets have been quite  
223 often shown to play an important pathophysiological role<sup>27</sup>, and new technologies and  
224 bioinformatic approaches for identification of such populations are steadily emerging<sup>28</sup>.  
225 Quite importantly, the SARS-CoV-2 Nsp6<sub>141-149</sub>, which was identified among our top  
226 potentially cross-reactive epitope pairs, has been recently described by an  
227 independent group.<sup>29</sup> To our knowledge, this is the first report on *in silico* predicted T

228 cell epitope cross-reactivity between SARS-CoV-2 and allergens. While a limitation of  
229 our study is the *in silico* nature of the work, the sequence homology between SARS-  
230 CoV-2 and clinically relevant respiratory allergens is along the lines of previously  
231 reported cross-reactivity between RNA virus- and allergen-derived peptides at the level  
232 of T memory cells.<sup>4</sup> Moreover, our current findings generate further hypotheses in how  
233 the adaptive immune system responds differentially with respect to the atopy status of  
234 the host. Our present study warrants an immediate investigation of these predicted T  
235 cell epitopes to link their possible role in driving the immune response against the  
236 SARS-CoV-2 and eventually shape COVID-19 outcome.

237 There are several different avenues through which the similarities may influence the  
238 host immune response. For instance, in hosts sensitized to one of the predicted  
239 aeroallergens, the identified similarities with the SARS-CoV-2 proteome may be  
240 protective if they prevent an overwhelming Th1 response and the accompanying  
241 cytokine storm. Furthermore, allergen-specific T cells may develop a memory  
242 response against heterologous SARS-CoV-2 epitopes, which is faster and more  
243 efficient. Conversely, such heterologous immune responses could have an adverse  
244 outcome by attenuating the antiviral response. T2 immune bias could potentially lead  
245 to inadequate virus clearance due to attenuated CD8<sup>+</sup> responses. Indeed, there is  
246 evidence of a reciprocal relationship between atopy and production of type I and III  
247 Interferons in response to viral infections<sup>30</sup>. Given that underlying atopic conditions  
248 have not been identified as a significant risk factor for severe clinical courses in those  
249 infected with SARS-CoV-2, the epitope homology most likely plays a protective role<sup>7,8</sup>.  
250 Interestingly, Jackson et al recently reported that nasal epithelial cells from children  
251 with atopic asthma express significantly lower levels of ACE2 receptor as compared to  
252 cells from children without asthma or with non-atopic asthma<sup>31</sup>. Similarly, another study  
253 using adult bronchial brush samples showed an inverse correlation between ACE2

254 gene expression and a Th2 dependent gene expression signature<sup>32</sup>. Differential  
255 expression of ACE2 receptors among atopic individuals could represent a distinct and  
256 unrelated mechanism of action in this context. Our *in silico* data provide ground to  
257 investigate the role of cellular immune responses in regards to the interaction between  
258 atopy/asthma and COVID-19. Indeed, the role of SARS-CoV-2-specific T cells in  
259 exposed and non-exposed individuals, thereby underlining the importance of  
260 heterologous immunity, has been very recently described<sup>33,34</sup>. Further experimental  
261 studies are needed to explore the involved pathogenetic mechanisms and potential  
262 clinical implications of underlying aeroallergen sensitization on the immune response  
263 to SARS-CoV-2.

264

#### 265 **Data availability**

266 The data used and analyzed in the present study are available from the  
267 corresponding author on reasonable request.

268

#### 269 **References**

- 270 1. Coronavirus Disease (COVID-19) Situation Reports; 22.09.2020. [Cited 2020  
271 September 22.] Available from [https://www.who.int/emergencies/diseases/novel-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/)  
272 [coronavirus-2019/situation-reports/](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/).
- 273 2. Cascella, M, Rajnik, M, Cuomo, A, Dulebohn, SC, Di Napoli, R. StatPearls.  
274 Features, Evaluation, and Treatment of Coronavirus (COVID-19). *Treasure Island*  
275 (FL); (2020).
- 276 3. Nextstrain / groups / blab / sars-like-cov; 21.09.2020. [Cited 2020 September 22.]  
277 Available from <https://nextstrain.org/groups/blab/sars-like-cov>.

- 278 4. Skevaki, C. *et al.* Influenza-derived peptides cross-react with allergens and  
279 provide asthma protection. *The Journal of allergy and clinical immunology*. **142**,  
280 804–14 (2018).
- 281 5. Pusch, E, Renz, H, Skevaki, C. Respiratory virus-induced heterologous immunity:  
282 Part of the problem or part of the solution? *Allergo Journal: interdisziplinäre*  
283 *Zeitschrift für Allergologie und Umweltmedizin* : Organ der Deutschen  
284 Gesellschaft für Allergie- und Immunitätsforschung 2018. **27**, 28–45.
- 285 6. Balz, K, Trassl, L, Härtel, V, Nelson, PP, Skevaki, C. Virus-Induced T Cell-  
286 Mediated Heterologous Immunity and Vaccine Development. *Frontiers in*  
287 *immunology*. **11**, 513 (2020).
- 288 7. Liu, S, Zhi, Y, Ying, S. COVID-19 and Asthma: Reflection During the Pandemic.  
289 *Clinical reviews in allergy & immunology*. **59**, 78–88 (2020).
- 290 8. Carli, G, Cecchi, L, Stebbing, J, Parronchi, P, Farsi, A. Is asthma protective  
291 against COVID-19? *Allergy* (2020).
- 292 9. Asthma Prevalence; 22.09.2020. [Cited 2020 September 22.] Available from  
293 <https://www.cdc.gov/asthma/data-visualizations/prevalence.htm>.
- 294 10. Zhu, Z. *et al.* Association of asthma and its genetic predisposition with the risk of  
295 severe COVID-19. *The Journal of allergy and clinical immunology*. **146**, 327-  
296 329.e4 (2020).
- 297 11. Shi, W. *et al.* Clinical characteristics of COVID-19 patients combined with allergy.  
298 *Allergy*. **75**, 2405–08 (2020).
- 299 12. Vita, R. *et al.* The Immune Epitope Database (IEDB): 2018 update. *Nucleic acids*  
300 *research*. **47**, D339-D343 (2019).
- 301 13. Mari, A, Scala, E, Palazzo, P, Ridolfi, S, Zennaro, D, Carabella, G. Bioinformatics  
302 applied to allergy: allergen databases, from collecting sequence information to

303 data integration. The Allergome platform as a model. *Cellular immunology*. **244**,  
304 97–100 (2006).

305 14.Cui, J. *et al.* Computer prediction of allergen proteins from sequence-derived  
306 protein structural and physicochemical properties. *Molecular immunology*. **44**,  
307 514–20 (2007).

308 15.Radauer, C, Bublin, M, Wagner, S, Mari, A, Breiteneder, H. Allergens are  
309 distributed into few protein families and possess a restricted number of  
310 biochemical functions. *The Journal of allergy and clinical immunology*. **121**, 847-  
311 52.e7 (2008).

312 16.Mari, A, Rasi, C, Palazzo, P, Scala, E. Allergen databases: current status and  
313 perspectives. *Current allergy and asthma reports*. **9**, 376–83 (2009).

314 17.The universal protein resource (UniProt). *Nucleic acids research*. **36**, D190-5  
315 (2008).

316 18.Peters, B, Sette, A. Generating quantitative models describing the sequence  
317 specificity of biological processes with the stabilized matrix method. *BMC*  
318 *bioinformatics*. **6**, 132 (2005).

319 19.Buus, S. *et al.* Sensitive quantitative predictions of peptide-MHC binding by a  
320 'Query by Committee' artificial neural network approach. *Tissue antigens*. **62**,  
321 378–84 (2003).

322 20.Moutaftsi, M. *et al.* A consensus epitope prediction approach identifies the breadth  
323 of murine T(CD8+)-cell responses to vaccinia virus. *Nature biotechnology*. **24**,  
324 817–19 (2006).

325 21.Nielsen, M, Lund, O. NN-align. An artificial neural network-based alignment  
326 algorithm for MHC class II peptide binding prediction. *BMC bioinformatics*. **10**, 296  
327 (2009).

- 328 22. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC bioinformatics*.  
329 **10**, 421 (2009).
- 330 23. Nielsen, M. *et al.* Reliable prediction of T-cell epitopes using neural networks with  
331 novel sequence representations. *Protein science : a publication of the Protein*  
332 *Society*. **12**, 1007–17 (2003).
- 333 24. Hoof, I. *et al.* NetMHCpan, a method for MHC class I binding prediction beyond  
334 humans. *Immunogenetics*. **61**, 1–13 (2009).
- 335 25. Andreatta, M. *et al.* Accurate pan-specific prediction of peptide-MHC class II  
336 binding affinity with improved binding core identification. *Immunogenetics*. **67**,  
337 641–50 (2015).
- 338 26. Cock, PJA. *et al.* Biopython: freely available Python tools for computational  
339 molecular biology and bioinformatics. *Bioinformatics* (Oxford, England). **25**, 1422–  
340 23 (2009).
- 341 27. Murray, SE, Toren, KG, Parker, DC. Peripheral CD4(+) T-cell tolerance is induced  
342 in vivo by rare antigen-bearing B cells in follicular, marginal zone, and B-1  
343 subsets. *European journal of immunology*. **43**, 1818–27 (2013).
- 344 28. Arvaniti, E, Claassen, M. Sensitive detection of rare disease-associated cell  
345 subsets via representation learning. *Nature communications*. **8**, 14825 (2017).
- 346 29. Grifoni, A. *et al.* Sequence Homology and Bioinformatic Approach Can Predict  
347 Candidate Targets for Immune Responses to SARS-CoV-2. *Cell host & microbe*.  
348 **27**, 671-680.e2 (20202).
- 349 30. Edwards, MR. *et al.*, Viral infections in allergy and immunology: How allergic  
350 inflammation influences viral infections and illness. *The Journal of allergy and*  
351 *clinical immunology*. **140**, 909–20 (2017).



- 352 31. Jackson, DJ. *et al.* Association of respiratory allergy, asthma, and expression of  
353 the SARS-CoV-2 receptor ACE2. *The Journal of allergy and clinical immunology*.  
354 **146**, 203-206.e3 (2020).
- 355 32. Bradding, P. *et al.* ACE2, TMPRSS2, and furin gene expression in the airways of  
356 people with asthma-implications for COVID-19. *The Journal of allergy and clinical*  
357 *immunology*. **146**, 208–11 (2020).
- 358 33. Braun, J. *et al.* SARS-CoV-2-reactive T cells in healthy donors and patients with  
359 COVID-19. *Nature* (2020).
- 360 34. Grifoni, A. *et al.* Targets of T Cell Responses to SARS-CoV-2 Coronavirus in  
361 Humans with COVID-19 Disease and Unexposed Individuals. *Cell*. **181**, 1489-  
362 1501.e15 (2020).

363 **Figure legends**

364 **Figure 1: Schematic overview of the bioinformatics approaches**

365 A: Pipeline 1; SARS-CoV-2 proteins were aligned against >2500 allergen protein  
366 sequences (see methods) and MHC class I-and II- restricted potentially cross-reactive  
367 T cell epitope pairs were identified for the most frequent human HLA alleles. B: Pipeline  
368 2; In an independent framework, we performed the comparative analysis of sequential  
369 *kmers* from SARS-CoV-2 protein sequences with known IEDB allergen peptides to  
370 predict the cross-reactive viral peptide pool.

371

372

373

374

375

376

377

378

379

380

381

382


383

384

385

386

### Linear T cell epitopes pool



390  
391  
392  
393  
394

**Table 1. The Top 5 candidate human HLA class I T cell potentially cross-reactive epitope pairs between SARS-CoV-2 and aero-and food-allergens based on pair combined score and application of additional clinical and conservation related criteria (see Suppl. Figure 1) (pipeline 1). Pcs=pair combined score**

		Top 30 pair combined score					
		candidate	allergen epitope	protein family	viral epitope	MHC allele	pcs
<b>MHC I</b> <b>AERO</b>	Nr.1	Mus m 1	GSNTFTILK	Lipocalin	TSNSFDVLK	HLA-A_11_01	8188
	Nr. 2	Asp f 5	MLYEVLWNL	Fungalysin metalloprotease	YLYALVYFL	HLA-A_02_01	5974
	Nr. 3	Aln g 1	SGVSPVSYQK	Bet v 1 family	ATSRTLSEYK	HLA-A_11_01	2413
	Nr.4	Phl p 5	KYKTFVATF	Group 5/6 grass pollen allergen	MFDAYVNTF	HLA-A_24_02	1910
	Nr. 5	Mus m 1	GSNTFTILK	Lipocalin	VTNNTFTLK	HLA-A_11_01	3286
<b>FOOD</b>	Nr. 1	Gal d 5	FLGHFIYSV	Serum albumin	TMADLVYAL	HLA-A_02_01	1408
	Nr. 2	Gal d 6	YLLDLLPAA	Lipoprotein	TLMNVLTLV	HLA-A_02_01	4185
	Nr. 3	Gal d 6	RPAYRRYLL	Lipoprotein	RPPLNRNYV	HLA-B_07_02	2274
	Nr. 4	Cor a 1	APHGGGSIL	Bet v 1 family	VPGLPGTIL	HLA-B_07_02	2571
	Nr. 5	Gal d 6	KVFRFSMFK	Lipoprotein	LVASIKNFK	HLA-A_11_01	1194

395  
396

397  
398  
399  
400  
401  
402

**Table 2. The Top 5 candidate human HLA class II T cell potentially cross-reactive epitope pairs between SARS-CoV-2 and aero-and food-allergens based on pair combined score and application of additional clinical and conservation related criteria (see Fig.1) (pipeline 1)**  
**Pcs=pair combined score**


		Top 30 pair combined score					
<b>MHC II</b>		candidate	allergen epitope	protein family	viral epitope	MHC allele	pcs
<b>AERO</b>	Nr.1	Phl p 5	FVATFGAAS	Group 5/6 grass pollen allergen	FSSTFNVPM	HLA-DRB1_04_01	87
	Nr. 2	Asp f 4	LTALAAGSA	Unclassified	VTALRANSA	HLA-DRB1_01_01	479
	Nr. 3	Phl p 5	FVATFGAAS	Group 5/6 grass pollen allergen	FSSTFNVPM	HLA-DRB1_04_01	53
	Nr.4	Phl p 5	FVATFGPAS	Group 5/6 grass pollen allergen	FSSTFNVPM	HLA-DRB1_04_01	29
	Nr. 5	Phl p 5	FKVAATAAN	Group 5/6 grass pollen allergen	FSSTFNVPM	HLA-DRB1_04_01	38
<b>FOOD</b>	Nr. 1	Gal d 5	FLYAPAILS	Serum albumin	FYILPSIIS	HLA-DRB1_01_01	299
	Nr. 2	Gal d 6	ILVDAVLKE	Lipoprotein	VVADAVIKT	HLA-DRB1_03_01	113
	Nr. 3	Gal d 6	VYSDVPIEK	Lipoprotein	VVADAVIKT	HLA-DRB1_03_01	29
	Nr. 4	Ara h 1	FIMPAAHPV	Cupin	FVMMSAPPA	HLA-DRB1_01_01	258
	Nr. 5	Gal d 5	FLYAPAILS	Serum albumin	FLYENAFLP	HLA-DRB1_01_01	81

403

**Table 3. HLA-I binding high confidence (IC50 < 50nm) SARS-CoV-2 antigenic peptides (pipeline 2)**

Allele	HLA-I-Binding Peptide	IC50	SARS-CoV-2 Protein name
HLA-A*68:01	NIFGTVYEK	6	R1AB_SARS2_Replicase_polyprotein
HLA-A*02:06	YTVELGTEV	9.4	R1A_SARS2_Replicase_polyprotein
HLA-A*68:02	YTVELGTEV	10.8	R1A_SARS2_Replicase_polyprotein
HLA-B*15:03	LASHMYCSF	10.8	R1A_SARS2_Replicase_polyprotein
HLA-B*40:02	HEGKTFYVL	11	SPIKE_SARS2_Spike_glycoprotein
HLA-B*40:01	GETLPTEVL	11.9	R1AB_SARS2_Replicase_polyprotein
HLA-A*02:06	TVYEKLKPV	13.4	R1AB_SARS2_Replicase_polyprotein
HLA-A*30:02	ASHMYCSFY	13.9	R1A_SARS2_Replicase_polyprotein
HLA-B*40:01	HEGKTFYVL	13.9	SPIKE_SARS2_Spike_glycoprotein
HLA-A*11:01	NIFGTVYEK	24	R1AB_SARS2_Replicase_polyprotein
HLA-B*35:01	LASHMYCSF	24.5	R1A_SARS2_Replicase_polyprotein
HLA-A*68:02	TVYEKLKPV	26	R1AB_SARS2_Replicase_polyprotein
HLA-A*02:01	WLTNIFGTV	34.1	R1AB_SARS2_Replicase_polyprotein
HLA-A*02:06	WLTNIFGTV	34.7	R1AB_SARS2_Replicase_polyprotein
HLA-B*15:03	LTNIFGTVY	35.7	R1AB_SARS2_Replicase_polyprotein
HLA-B*15:25	LASHMYCSF	39.1	R1A_SARS2_Replicase_polyprotein
HLA-B*15:25	LTNIFGTVY	39.8	R1AB_SARS2_Replicase_polyprotein
HLA-A*02:01	TVYEKLKPV	47.8	R1AB_SARS2_Replicase_polyprotein
HLA-B*15:01	LASHMYCSF	48.1	R1A_SARS2_Replicase_polyprotein
HLA-B*15:03	ASHMYCSFY	49	R1A_SARS2_Replicase_polyprotein

# Figures



**Figure 1**

Schematic overview of the bioinformatics approaches A: Pipeline 1; SARS-CoV-2 proteins were aligned against >2500 allergen protein sequences (see methods) and MHC class I-and II- restricted potentially cross-reactive T cell epitope pairs were identified for the most frequent human HLA alleles. B: Pipeline 2; In an independent framework, we performed the comparative analysis of sequential kmers from SARS-CoV-2 protein sequences with known IEDB allergen peptides to predict the cross-reactive viral peptide pool.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementsHomologiesbetweenSARSCoV2Balzetal.pdf](#)
- [SupplementaryTableS1.xlsx](#)
- [SupplementaryTableS2.xlsx](#)
- [SupplementaryTables47.xlsx](#)
- [SupplementaryTableS8.xlsx](#)
- [SupplementaryTableS9.xlsx](#)

- [SupplementaryTableS10.xls](#)
- [SupplementaryTableS11.xlsx](#)