

Quantum Networks and Symmetries

Kiara Hansenne

University of Siegen <https://orcid.org/0000-0002-5914-002X>

Zhen-Peng xu (✉ zhen-peng.xu@uni-siegen.de)

<https://orcid.org/0000-0001-8514-2603>

Tristan Kraft

University of Innsbruck

Otfried Gühne

University of Siegen <https://orcid.org/0000-0002-6033-0867>

Article

Keywords:

Posted Date: September 16th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-885489/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Quantum Networks and Symmetries

Kiara Hansenne,^{1,*} Zhen-Peng Xu,^{1,*} Tristan Kraft,^{2,1} and Otfried Gühne¹

¹*Naturwissenschaftlich-Technische Fakultät, Universität Siegen, Walter-Flex-Straße 3, 57068 Siegen, Germany*

²*Institute for Theoretical Physics, University of Innsbruck, Technikerstraße 21A, 6020 Innsbruck, Austria*

(Dated: September 7, 2021)

Quantum networks are promising tools for the implementation of long-range quantum communication. The characterization of quantum correlations in networks and their usefulness for information processing is therefore central for the progress of the field, but so far only results for small basic network structures or pure quantum states are known. Here we show that symmetries provide a versatile tool for the analysis of correlations in quantum networks. We provide an analytical approach to characterize correlations in large network structures with arbitrary topologies. As examples, we show that entangled quantum states with a bosonic or fermionic symmetry can not be generated in networks; moreover, cluster and graph states are not accessible. Our methods can be used to design certification methods for the functionality of specific links in a network and have implications for the design of future network structures.

INTRODUCTION

A central paradigm for quantum information processing is the notion of quantum networks [1–4]. In an abstract sense, a quantum network consists of quantum systems as nodes on specific locations, where some of the nodes are connected via links. These links correspond to quantum channels, which may be used to send quantum information (e.g., a polarized photon) or where entanglement may be distributed. Crucial building blocks for the links, such as photonic quantum channels between a satellite and ground stations [5–8] or the high-rate distribution of entanglement between nodes [9, 10] have recently been experimentally demonstrated. Clearly, such real physical implementations are always noisy and may only work probabilistically, but there are various theoretical approaches to deal with this [11–14].

For the further progress of the field, it is essential to design methods for the certification and benchmarking of a given network structure or a single specific link within it. Moreover, it is important to analyze the capability of a given network structure to generate global quantum correlations. In principle, any multiparticle state can be generated in a connected network, but this may require complicated protocols using long-lived quantum memories, assisted by coherent operations and classical communication. So, in view of current experimental limitations, the question arises which states can be prepared in the network with moderate effort, e.g., with simple local operations.

This question has attracted some attention, with several lines of research emerging. First, the problem has been considered in the classical setting, such as the analysis of causal structures [15–17] or in the study of hidden variable models, where the hidden variables are not

equally distributed between every party [18–24]. Concerning quantum correlations, several initial works appeared in the last year, suggesting slightly different definitions of network entanglement [25–28]. These have been further investigated [29–31] and methods from the classical realm have been extended to the quantum scenario [32]. Still, the present results are limited to simple networks like the triangle network, noise-free networks or networks build from specific quantum states, or bounded to small dimensions due to numerical limitations.

In this paper we present an analytical approach to characterize quantum correlations in arbitrary network topologies. Our approach is based on symmetries, which may occur as permutation symmetries or invariance under certain local unitary transformations. Symmetries play an outstanding role in various field of physics [33] and they have already turned out to be useful for various other problems in quantum information theory [34–40]. On a technical level, we combine the inflation technique for quantum networks [17, 27] with estimates known from the study of entropic uncertainty relations [41–43].

Based on our approach, we derive simply testable inequalities in terms of expectation values, which can be used to decide whether a given state may be prepared in a network or not. With this we can prove that large classes of states cannot be prepared in networks using simple communication, for instance all multiparticle graph states with up to twelve vertices with noise, as well as all mixed entangled permutationally symmetric states. This delivers various methods for benchmarking: First, the observation of such states in a network certifies the implementation of advanced network protocols. Second, our results allow to design simple tests for the proper working of a specific link in a given network.

RESULTS

Network entanglement.— To start, let us define the

* K.H. and Z.P.X. contributed equally. Correspondence should be addressed to Z.P.X.: zhen-peng.xu@uni-siegen.de




FIG. 1. *Triangle quantum network.* Three sources ϱ_a , ϱ_b and ϱ_c distribute parties to three nodes, Alice, Bob and Charlie (A , B and C). Alice, Bob and Charlie each end up with a bipartite system $X = X_1X_2$ on which they perform a local channel $\mathcal{E}_X^{(\lambda)}$ ($X = A, B, C$) depending on a classical random variable λ .

types of correlations that can be prepared in a network. In the simplest scenario Alice, Bob and Charlie aim to prepare a tripartite quantum state using three bipartite source states ϱ_a , ϱ_b and ϱ_c , see Fig. 1. Parties belonging to a same source state are sent to different parties of the network, i.e. A , B or C , such that the global state reads $\varrho_{ABC} = \varrho_c \otimes \varrho_b \otimes \varrho_a$. Note that here the order of the parties on both sides of the equation is different. After receiving the states, each party may still apply a local operation \mathcal{E}_X (for $X = A, B, C$), in addition these operations may be coordinated by shared randomness. This leads to a global state of the form

$$\varrho = \sum_{\lambda} p_{\lambda} \mathcal{E}_A^{(\lambda)} \otimes \mathcal{E}_B^{(\lambda)} \otimes \mathcal{E}_C^{(\lambda)} [\varrho_{ABC}], \quad (1)$$

and the question arises, which three-party states can be written in this form and which cannot?

Some remarks are in order: First, the definition of network states in Eq. (1) can directly be extended to more parties or more advanced sources, e.g. one can consider the case of five parties A, B, C, D, E , where some sources distribute four-party states between some of the parties. Second, the scenario considered uses local operations and shared randomness (LOSR) as allowed operations, which is a smaller set than local operations and classical communication (LOCC). In fact, LOCC are much more difficult to implement, but using LOCC and teleportation *any* tripartite state can be prepared from bipartite sources. On the other hand, the set LOSR is strictly larger than, e.g., the unitary operations considered in [25, 26]. Finally, the discerning reader may have noticed that in Eq. (1) the state ϱ_{ABC} does not depend on the shared random variable λ , but since the dimension of the source states ϱ_i is not bounded one can always remove a dependency on λ in the ϱ_i by enlarging the dimension [27]. Equivalently, one may remove the depen-

dency of the maps \mathcal{E}_X on λ and the shared randomness may be carried by the source states only.

Symmetries.— Symmetry groups can act on quantum states in different ways. First, the elements of a unitary symmetry group may act transitively on the density matrix ϱ . That is, ϱ is invariant under transformations like

$$\varrho \mapsto U \varrho U^{\dagger} = \varrho. \quad (2)$$

If $\varrho = |\psi\rangle\langle\psi|$ is pure, this implies $U|\psi\rangle = e^{i\phi}|\psi\rangle$ and $|\psi\rangle$ is, up to some phase, an eigenstate of some operator. Second, for pure states one can also identify directly a certain subspace of the entire Hilbert space that is equipped with a certain symmetry, e.g. symmetry under exchange of two particles. Denoting by Π the projector onto this subspace, the symmetric pure states are defined via

$$\Pi|\psi\rangle = |\psi\rangle \quad (3)$$

and for mixed states one has $\varrho = \Pi\varrho\Pi$. Note that if $\varrho = \sum_k p_k |\phi_k\rangle\langle\phi_k|$ has some decomposition into pure states, then each $|\phi_k\rangle = \Pi|\phi_k\rangle$ has to be symmetric, too.

In the following, we consider mainly two types of symmetries. First, we consider multi-qubit states obeying a symmetry as in Eq. (2) where the symmetry operations consist of an Abelian group of tensor products of Pauli matrices. These groups are usually referred to as stabilizers in quantum information theory [44], and they play a central role in the construction of quantum error correcting codes. Pure states obeying such symmetries are also called stabilizer states, or, equivalently, graph states [45]. Second, we consider states with a permutational (or bosonic) symmetry [36, 39], obeying relations as in Eq. (3) with Π being the projector onto the symmetric subspace.

GHZ states.— As a warming-up exercise we discuss the Greenberger-Horne-Zeilinger (GHZ) state of three qubits,

$$|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle) \quad (4)$$

in the triangle scenario. This simple case was already the main example in previous works on network correlations [25–27], but it allows us to introduce our concepts and ideas in a simple setting, such that their full generalization is later conceivable.

The GHZ state is an eigenstate of the observables

$$g_1 = X_A X_B X_C, \quad g_2 = 1_A Z_B Z_C, \quad g_3 = Z_A Z_B 1_C. \quad (5)$$

Here and in the following we use the shorthand notation $1_A X_B Y_C = \mathbb{1} \otimes \sigma_x \otimes \sigma_y$ for tensor products of Pauli matrices. Indeed, these g_k commute and generate the stabilizer $\mathcal{S} = \{\mathbb{1}, g_1, g_2, g_3, g_1 g_2, g_1 g_3, g_2 g_3, g_1 g_2 g_3\}$. Clearly, for any $S_i \in \mathcal{S}$ we have $S_i |GHZ\rangle = |GHZ\rangle$ and so $\langle GHZ | S_i | GHZ \rangle = 1$.

As a tool for studying network entanglement, we use the inflation technique [17, 32]. The basic idea is depicted




FIG. 2. *Triangle network and two of its inflations.* The first figure represents the triangle network of Fig. 1, with global state ϱ . Using the same source states (represented by lines of same colour) and same local channels, one can build the so-called inflated state τ , which is separable with respect to the $ABC|A'B'C'$ partition. The state γ is build similarly, but with a rewiring of the sources, leading to an inflated state that is in general not separable and different from τ . Still, this procedure imposes that several marginals of ϱ , τ and γ are equal, e.g. $\varrho_{AC} = \tau_{A'C'} = \gamma_{A'C'}$.

in Fig. (2). If a state ϱ can be prepared in the network scenario, then one can also consider a scenario where each source state is sent two-times to multiple copies of the parties. In this multicopy scenario, the source states may, however, also be wired in a different manner. In the simplest case of doubled sources, this may lead to two different states, τ and γ . Although ϱ , τ and γ are different states, some of their marginals are identical, see Fig. (2) and Supplementary Note 1. If one can prove that states τ and γ with the marginal conditions do not exist, then ϱ cannot be prepared in the network.

Let us start by considering the correlation $\langle Z_A Z_B \rangle$ in ϱ , τ and γ . The values are equal in all three states, $\langle Z_A Z_B \rangle_{\varrho} = \langle Z_A Z_B \rangle_{\tau} = \langle Z_A Z_B \rangle_{\gamma}$, and the same holds for the correlation $\langle Z_B Z_C \rangle$. Note that these should be large, if ϱ is close to a GHZ state, as $Z_A Z_B$ is an element of the stabilizer. Using the general relation $\langle Z_A Z_C \rangle \geq \langle Z_A Z_B \rangle + \langle Z_B Z_C \rangle - 1$ [27] we can use this to estimate $\langle Z_A Z_C \rangle$ in γ . Due to the marginal conditions, we have $\langle Z_A Z_C \rangle_{\gamma} = \langle Z_{A'} Z_C \rangle_{\tau}$, implying that this correlation in τ must be large, if ϱ is close to a GHZ state. On the other hand, the correlation $\langle X_A X_B X_C \rangle$ corresponds also to a stabilizer element and should be large in the state ϱ as well as in τ .

The key observation is that the observables $X_A X_B X_C$ and $Z_{A'} Z_C$ anticommute; moreover, they have only eigenvalues ± 1 . For this situation, strong constraints on the expectation values are known: If M_i are pairwise anticommuting observables with eigenvalues ± 1 , then $\sum_i \langle M_i \rangle^2 \leq 1$ [43]. This fact has already been used to derive entropic uncertainty relations [41, 42] or monogamy relations [46, 47]. For our situation, it directly implies that for τ the correlations $\langle X_A X_B X_C \rangle$ and $\langle Z_{A'} Z_C \rangle$ cannot both be large. Or, expressing everything in terms of the original state ϱ , if $\langle Z_A Z_B \rangle + \langle Z_B Z_C \rangle - 1 \geq 0$ then a condition for preparability of a state in the network is

$$\langle X_A X_B X_C \rangle^2 + (\langle Z_A Z_B \rangle + \langle Z_B Z_C \rangle - 1)^2 \leq 1. \quad (6)$$




FIG. 3. *Square network and two of its inflations.* Similar to the triangle network of Fig. 2, the state τ is generated using two copies of the sources and channels used to generate ϱ . Then, one goes to a higher-order inflation by using three copies of the sources and the local channels. By rewiring one obtains the inflated state ξ . Again, one has several equalities between the marginals of ϱ , τ and ξ .

This is clearly violated by the GHZ state. In fact, if one considers a GHZ state mixed with white noise, $\varrho = p|GHZ\rangle\langle GHZ| + (1-p)\mathbb{1}/8$, then these states are detected already for $p \geq 4/5$. Note that using the other observables of the stabilizer and permutations of the particles, also other conditions like $\langle Y_A Y_B X_C \rangle^2 + (\langle Z_A Z_B \rangle + \langle Z_A Z_C \rangle - 1)^2 \leq 1$ can be derived.

Using these techniques as well as concepts based on covariance matrices [28, 29] and classical networks [23], one can also derive bounds on the maximal GHZ fidelity achievable by network states. In fact, for network states

$$F_{GHZ} = \langle GHZ | \varrho | GHZ \rangle \leq \frac{1}{\sqrt{2}} \approx 0.7071 \quad (7)$$

holds, as explained in Supplementary Note 2. This is a clear improvement on previous analytical bounds, although it does not improve a fidelity bound obtained by numerical convex optimization [27].

Cluster and graph states.— The core advantage of our approach is the fact that it can directly be generalized to more particles and complicated networks, while the existing numerical and analytical approaches are mostly restricted to the triangle scenario.

Let us start the discussion with the four-qubit cluster state $|C_4\rangle$. This may be defined as the unique common +1-eigenstate of

$$\begin{aligned} g_1 &= X_A Z_B 1_C Z_D, & g_2 &= Z_A X_B Z_C 1_D, \\ g_3 &= 1_A Z_B X_C Z_D, & g_4 &= Z_A 1_B Z_C X_D. \end{aligned} \quad (8)$$

For later generalization, it is useful to note that the choice of these observables is motivated by a graphical analogy. For the square graph in Fig. 3 one can associate to any vertex a stabilizing operator in the following manner: One takes X on the vertex i , and Z on its neighbours, i.e. the vertices connected to i . This delivers the observables in Eq. (8), but it may also be applied to general graphs, leading to the notion of graph states [45].

If a quantum state can be prepared in the square network, then we can consider the third order inflated state ξ shown in Fig. 3. In the inflation ξ , the three observables $X_{B'}X_D$, $Z_{B'}X_CZ_D$, and $X_A Y_B Y_D$ anticommute. These observables act on marginals which are identical to those in ϱ . Consequently, for any state that can be prepared in the square network, the relation

$$\langle X_{B'}X_D \rangle^2 + \langle Z_{B'}X_CZ_D \rangle^2 + \langle X_A Y_B Y_D \rangle^2 \leq 1 \quad (9)$$

holds. All these observables are also within the stabilizer of the cluster state, so the cluster state violates this inequality with an lhs equal to three. This proves that cluster states mixed with white noise cannot be prepared in the square network for $p > 1/\sqrt{3} \approx 0.577$. Again, with the same strategy different nonlinear witnesses like $\langle X_A X_C \rangle^2 + \langle Y_A Y_B Z_C Z_D \rangle^2 \leq 1$ for network entanglement in the square network can be derived. This follows from the second inflation in Fig. 3, see also the detailed discussion below. Furthermore, it can be shown that states with a cluster state fidelity of $F_{C_4} > 0.7377$ cannot be prepared in a network. The full discussion is given in Supplementary Note 3.

This approach can be generalized to graph states. As already mentioned, starting from a general graph one can define a graph state by stabilizing operators in analogy to Eq. (8). The resulting states play an eminent role in quantum information processing. For instance, the so-called cluster states, which correspond to graphs of quadratic and cubic square lattices, are resource states for measurement-based quantum computation [48] and topological error correction [49, 50].

Applying the presented ideas to general graphs results in the following: If a graph contains a triangle, then under simple and weak conditions an inequality similar to Eq. (6) can be derived. This excludes then the preparability of noisy graph states in *any* network with bipartite sources only. Note that this is a stronger statement than proving network entanglement only for the network corresponding to the graph, as was considered above for the cluster state.

At first sight, the identification of a specific triangle in the graph may seem a weak condition, but here the entanglement theory of graph states helps: It is well known that certain transformations of the graph, so-called local complementations, change the graph state only by a local unitary transformation [51, 52], so one may apply these to generate the triangle with the required properties. Indeed, this works for all cases we considered (e.g. the full classification up to 12 qubits from Refs. [53, 54]) and we can summarize¹:

¹ We note that similar statements on genuine multiparticle states or symmetric states have been formulated in Refs. [26], but these are restricted to pure states only and cannot be applied in the presence of noise.

Observation 1. (a) No graph state with up to 12 vertices can be prepared in a bipartite network. (b) If a graph contains a vertex with degree $d \leq 3$, then it cannot be prepared in any network with bipartite sources. (c) The two- and three-dimensional cluster states cannot be prepared in any network.

In all the cases, it follows that graph states mixed with white noise, $\varrho = p|G\rangle\langle G| + (1-p)\mathbb{1}/2^N$ are network entangled for $p > 4/5$, independently of the number of qubits. A detailed discussion is given in Supplementary Note 3.

Permutational symmetry.— Now we consider multiparticle quantum states of arbitrary dimension that obey a permutational or bosonic symmetry. Mathematically, these states act on the symmetric subspace only, meaning that $\varrho = \Pi^+ \varrho \Pi^+$, where Π^+ is the projector on the symmetric subspace. For example, in the case of three qubits this space is four-dimensional, and spanned by the Dicke states $|D_0\rangle = |000\rangle$, $|D_1\rangle = (|001\rangle + |010\rangle + |100\rangle)/\sqrt{3}$, $|D_2\rangle = (|011\rangle + |101\rangle + |110\rangle)/\sqrt{3}$, and $|D_3\rangle = |111\rangle$.

The symmetry has several consequences [36, 39]. First, if one has a decomposition $\varrho = \sum_k p_k |\psi_k\rangle\langle\psi_k|$ into pure states, then all $|\psi_k\rangle$ have to come from the symmetric space, too. Since pure symmetric states can be either fully separable (like $|D_0\rangle$) or genuine multiparticle entangled (like $|D_1\rangle$), this implies that mixed symmetric states have also only these two possibilities. That is, if a mixed symmetric state is separable for one bipartition, it must be fully separable.

Second, permutational invariance can also be characterized by the flip operator $F_{XY} = \sum_{ij} |ij\rangle_{XY} \langle ji|_{XY}$ on the particles X and Y . Symmetric multiparticle states obey $\varrho = F_{XY}\varrho = \varrho F_{XY}$ for any pair of particles, where the second equality directly follows from hermiticity. Conversely, concluding full permutational symmetry from two-particle properties, does only require this relation for pairs such that the F_{XY} generate the full permutation group. Finally, it is easy to check that if the marginal ϱ_{XY} of a multiparticle state ϱ obeys $\varrho_{XY} = F_{XY}\varrho_{XY}$, then the full state ϱ obeys the same relation, too.

Armed with these insights, we can explain the idea for our main result. Consider a three-particle state with bosonic symmetry which can be prepared in a triangle network, the inflation γ from Fig. (2) and the reduced state γ_{ABC} in this inflation. This obeys $\gamma_{ABC} = F_{XY}\gamma_{ABC}$ for XY equal to AB or BC . Since $F_{AB}F_{BC}F_{AB} = F_{AC}$, this implies that the reduced state $\tau_{AC'}$ obeys $\tau_{AC'} = F_{AC'}\tau_{AC'}$ and hence also the six-particle state τ . Moreover, τ also obeys similar constraints for other pairs of particles (like AB , BC , $A'B'$ and $B'C'$) and it is easy to see that jointly with AC' these generate the full permutation group. So, τ must be fully symmetric. But τ is separable with respect to the $ABC|A'B'C'$ bipartition, so τ and hence $\varrho = \tau_{ABC}$ must be fully separable.

The same argument can easily be extended to more

complex networks, which are not restricted to use bipartite sources. We can summarize¹:

Observation 2. *Let ρ be a permutationally symmetric multipartite state of N particles. Then, ρ can be generated by a network with $(N - 1)$ -partite sources, if and only if it is fully separable.*

We add that this Observation can also be extended to the case of fermionic antisymmetry, a detailed discussion is given in the Supplementary Note 4.

Certifying network links.— For the technological implementation of quantum networks, it is of utmost importance to design certification methods to test and benchmark different realizations. One of the basic questions is, whether a predefined quantum link works or not. Interestingly, Consider a network where the link between two particles is absent or not properly working. For definiteness, we may consider the square network on the lhs of Fig. 3 and the parties A and C . In the second inflation τ we have for the marginals $\tau_{AC} = \tau_{A'C}$. This implies that the observables $X_A X_C$ and $Z_A Z_C$ on the original state ρ correspond to anticommuting observables on τ , so we have $\langle X_A X_C \rangle^2 + \langle Z_A Z_C \rangle^2 \leq 1$. Using higher-order inflations, one can extend and formulate it for general networks: If a state can be prepared in a network with bipartite sources but without the link AC , then

$$\langle X_A X_C P_{R_1} \rangle^2 + \langle Y_A Y_C P_{R_2} \rangle^2 + \langle Z_A Z_C P_{R_3} \rangle^2 \leq 1. \quad (10)$$

Here the P_{R_i} are arbitrary observables on disjoint subsets of the other particles, $R_i \cap R_j = \emptyset$. If a state was indeed prepared in a real quantum network then violation of this inequality proves that the link AC is working and distributing entanglement. In Supplementary Note 5, details are discussed and examples are given, where this test allows to certify the functionality of a link even if the reduced state ρ_{AC} is separable.

DISCUSSION

We have provided an analytical method to analyze correlations arising in quantum networks from few measurements. With this, we have shown that large classes of states with symmetries, namely noisy graph states and permutationally symmetric states cannot be prepared in networks. Moreover, our approach allows to design simple tests for the functionality of a specified link in a network.

Our results open several research lines of interest. First, they are of direct use to analyze quantum correlations in experiments and to show that multipartite entanglement is needed to generate observed quantum correlations. Second, they are useful for the design of networks in the realistic setting: For instance, we have shown that the generation of graph states from bipartite sources necessarily requires at least some communication

between the parties, which may be of relevance for quantum repeater schemes based on graph states that have been designed [55]. Moreover, it has been shown that GHZ states provide an advantage for multipartite conference key agreement over bipartite sources [56], which may be directly connected to the fact that their symmetric entanglement is inaccessible in networks. Finally, our result open the door for further studies of entanglement in networks, e.g. using limited communication (first results on this have recently been reported [57]) or restricted quantum memories, which is central for future realizations of a quantum internet.

ACKNOWLEDGMENTS

We thank Xiao-Dong Yu for discussions. We thank Carlos de Gois for pointing out a typo in a figure. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, project numbers 447948357 and 440958198), the Sino-German Center for Research Promotion (Project M-0294), and the ERC (Consolidator Grant 683107/TempoQ). K.H. acknowledges support from the House of Young Talents of the University of Siegen. Z.P.X. acknowledges support from the Humboldt foundation.

-
- [1] H. J. Kimble. The quantum internet. *Nature* **453**, 1023 (2008).
 - [2] C. Simon. Towards a global quantum network. *Nat. Phot.* **11**, 678 (2017).
 - [3] S. Wehner, D. Elkouss, and R. Hanson. Quantum internet: A vision for the road ahead. *Science* **362**, 9288 (2018).
 - [4] J. Biamonte, M. Faccin, and M. De Domenico. Complex networks from classical to quantum. *Commun. Phys.* **2**, 53 (2019).
 - [5] J. Yin, Y. Cao, Y.-H. Li, et al. Satellite-based entanglement distribution over 1200 kilometers. *Science* **356**, 1140 (2017).
 - [6] S.-K. Liao, W.-Q. Cai, J. Handsteiner, et al. Satellite-Relayed Intercontinental Quantum Network. *Phys. Rev. Lett.* **120**, 030501 (2017).
 - [7] S.-K. Liao, W.-Q. Cai, W.-Y. Liu, et al. Satellite-to-ground quantum key distribution. *Nature* **549**, 43 (2018).
 - [8] J. Yin, Y.-H. Li, S.-K. Liao, M. Yang, Y. Cao, L. Zhang, J.-G. Ren, W.-Q. Cai, W.-Y. Liu, S.-L. Li, et al. Entanglement-based secure quantum cryptography over 1,120 kilometres. *Nature* **582**, 501 (2020).
 - [9] P. C. Humphreys, N. Kalb, J. P. Morits, R. N. Schouten, R. F. Vermeulen, D. J. Twitchen, M. Markham, and R. Hanson. Deterministic delivery of remote entanglement on a quantum network. *Nature* **558**, 268 (2018).
 - [10] L. J. Stephenson, D. P. Nadlinger, B. C. Nichol, S. An, P. Drmota, T. G. Ballance, K. Thirumalai, J. F. Goodwin, D. M. Lucas, and C. J. Ballance. High-Rate, High-

- Fidelity Entanglement of Qubits Across an Elementary Quantum Network. *Phys. Rev. Lett.* **124**, 110501 (2020).
- [11] E. Shchukin, F. Schmidt, and P. van Loock. Waiting time in quantum repeaters with probabilistic entanglement swapping. *Phys. Rev. A* **100**, 032322 (2019).
- [12] S. Khatiri, C. T. Matyas, A. U. Siddiqui, J. P. Dowling. Practical figures of merit and thresholds for entanglement distribution in quantum networks. *Phys. Rev. Research* **1**, 023032 (2019).
- [13] B. C. Coutinho, W. J. Munro, K. Nemoto, Y. Omar. Robustness of Noisy Quantum Networks. [arXiv:2103.03266](https://arxiv.org/abs/2103.03266).
- [14] L. Bugalho, B. C. Coutinho, Y. Omar. Distributing Multipartite Entanglement over Noisy Quantum Networks. [arXiv:2103.14759](https://arxiv.org/abs/2103.14759).
- [15] R. Chaves, L. Luft, T. O. Maciel, D. Gross, D. Janzing, and B. Schölkopf, Proceedings of the 30th Conference on Uncertainty in Artificial Intelligence (2014), p. 112.
- [16] R. Chaves, C. Majenz, and D. Gross. Information-theoretic implications of quantum causal structures. *Nature Commun.* **6**, 5766 (2015).
- [17] E. Wolfe, R. W. Spekkens, and T. Fritz. The Inflation Technique for Causal Inference with Latent Variables. *J. Causal Inference* **7**, 20170020 (2019).
- [18] C. Branciard, N. Gisin, and S. Pironio. Characterizing the Nonlocal Correlations Created via Entanglement Swapping. *Phys. Rev. Lett.* **104**, 170401 (2010).
- [19] C. Branciard, D. Rosset, N. Gisin, and S. Pironio. Bilocal versus nonbilocal correlations in entanglement-swapping experiments. *Phys. Rev. A* **85**, 032119 (2012).
- [20] T. Fritz. Beyond Bell's theorem: correlation scenarios. *New J. Phys.* **14**, 103001 (2012).
- [21] D. Rosset, C. Branciard, T. J. Barnea, G. Pütz, N. Brunner, and N. Gisin. Nonlinear Bell Inequalities Tailored for Quantum Networks. *Phys. Rev. Lett.* **116**, 010403 (2016).
- [22] M.-O. Renou, E. Bäumer, S. Boreiri, N. Brunner, N. Gisin, and S. Beigi. Genuine Quantum Nonlocality in the Triangle Network. *Phys. Rev. Lett.* **123**, 140401 (2019).
- [23] N. Gisin, J. D. Bancal, Y. Cai, et al. Constraints on nonlocality in networks from no-signaling and independence. *Nat. Commun.* **11**, 2378 (2020).
- [24] P. Contreras-Tejada, C. Palazuelos, J. I. de Vicente. Genuine multipartite entanglement in noisy quantum networks highly depends on the topology. [arXiv:2106.04634](https://arxiv.org/abs/2106.04634) (2021).
- [25] T. Kraft, S. Designolle, C. Ritz, N. Brunner, O. Gühne, and M. Huber. Quantum entanglement in the triangle network. *Phys. Rev. A* **103**, L060401 (2021).
- [26] M.-X. Luo. New Genuinely Multipartite Entanglement. *Adv. Quantum Technol.*, **4**, 2000123 (2021).
- [27] M. Navascués, E. Wolfe, D. Rosset, and A. Pozas-Kerstjens. Genuine Network Multipartite Entanglement. *Phys. Rev. Lett.* **125**, 240505 (2020).
- [28] J. Åberg, R. Nery, C. Duarte, and R. Chaves. Semidefinite Tests for Quantum Network Topologies. *Phys. Rev. Lett.* **125**, 110505 (2020).
- [29] T. Kraft, C. Spee, X.-D. Yu, and O. Gühne. Characterizing quantum networks: Insights from coherence theory. *Phys. Rev. A* **103**, 052405 (2021).
- [30] M.-X. Luo. Network configuration theory for all networks. [arXiv:2107.05846](https://arxiv.org/abs/2107.05846) (2021).
- [31] P. Contreras-Tejada, C. Palazuelos, J. I. de Vicente. Genuine Multipartite Nonlocality Is Intrinsic to Quantum Networks. *Phys. Rev. Lett.* **126**, 040501 (2021).
- [32] E. Wolfe, A. Pozas-Kerstjens, M. Grinberg, D. Rosset, A. Acín, and M. Navascués. Quantum Inflation: A General Approach to Quantum Causal Compatibility. *Phys. Rev. X* **11**, 021043 (2021).
- [33] S. Coleman, Aspects of Symmetry, Cambridge University Press, Massachusetts (1985).
- [34] R. F. Werner. Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model. *Phys. Rev. A* **40**, 4277 (1989).
- [35] K. G. H. Vollbrecht and R. F. Werner. Entanglement measures under symmetry. *Phys. Rev. A* **64**, 062307 (2001).
- [36] K. Eckert, J. Schliemann, D. Bruß, and M. Lewenstein. Quantum Correlations in Systems of Indistinguishable Particles. *Ann. Phys.* **299**, 88 (2002).
- [37] C. M. Caves, C. A. Fuchs, and R. Schack. Unknown quantum states: The quantum de Finetti representation. *J. Math. Phys.* **43**, 4537 (2002).
- [38] A. C. Doherty, P. A. Parrilo, and F. M. Spedalieri. Distinguishing Separable and Entangled States. *Phys. Rev. Lett.* **88**, 187904 (2002).
- [39] G. Tóth and O. Gühne. Entanglement and Permutational Symmetry. *Phys. Rev. Lett.* **102**, 170503 (2009).
- [40] C. Eltschka and J. Siewert. Entanglement of Three-Qubit Greenberger-Horne-Zeilinger-Symmetric States. *Phys. Rev. Lett.* **108**, 020502 (2012).
- [41] S. Wehner and A. Winter. Higher entropic uncertainty relations for anti-commuting observables. *J. Math. Phys.* **49**, 062105 (2008).
- [42] S. Niekamp, M. Kleinmann, and O. Gühne. Entropic uncertainty relations and the stabilizer formalism. *J. Math. Phys.* **53**, 012202 (2012).
- [43] G. Tóth and O. Gühne. Entanglement detection in the stabilizer formalism. *Phys. Rev. A* **72**, 022340 (2005).
- [44] D. Gottesman. Stabilizer Codes and Quantum Error Correction. [PhD Thesis, Caltech](https://arxiv.org/abs/quant-ph/9705052) (1997).
- [45] M. Hein, W. Dür, J. Eisert, R. Raussendorf, M. Van den Nest, and H.-J. Briegel, in *Quantum Computers, Algorithms and Chaos*, Proceedings of the International School of Physics "Enrico Fermi," Vol. 162, Varenna, 2005, edited by G. Casati, D. L. Shepelyansky, P. Zoller, and G. Benenti (IOS Press, Amsterdam, 2006). See also [arXiv:quant-ph/0602096](https://arxiv.org/abs/quant-ph/0602096).
- [46] M. C. Tran, R. Ramanathan, M. McKague, D. Kaszlikowski, and T. Paterek. Bell monogamy relations in arbitrary qubit networks. *Phys. Rev. A* **98**, 052325 (2018).
- [47] P. Kurzyński, T. Paterek, R. Ramanathan, W. Laskowski, and D. Kaszlikowski. Correlation Complementarity Yields Bell Monogamy Relations. *Phys. Rev. Lett.* **106**, 180402 (2011).
- [48] R. Raussendorf and H. J. Briegel. A One-Way Quantum Computer. *Phys. Rev. Lett.* **86**, 5188 (2001).
- [49] A. Kitaev. Fault-tolerant quantum computation by anyons. *Annals Phys.* **303**, 2 (2003).
- [50] X.-C. Yao, T.-X. Wang, H.-Z. Chen, et al. Experimental demonstration of topological error correction. *Nature* **482**, 489–494 (2012).
- [51] M. Van den Nest, J. Dehaene, and B. De Moor. Graphical description of the action of local Clifford transformations on graph states. *Phys. Rev. A* **69**, 022316 (2004).
- [52] M. Hein, J. Eisert, and H. J. Briegel. Multiparty entanglement in graph states. *Phys. Rev. A* **69**, 062311 (2004).
- [53] A. Cabello, L. E. Danielsen, A. J. López-Tarrida, and J. R. Portillo. Optimal preparation of graph states. *Phys.*

- Rev. A **83**, 042314 (2011).
- [54] L. E. Danielsen, [Database of Entanglement in Graph States \(2011\)](#), see also <http://www.iu.uib.no/~larsed/entanglement/>.
- [55] M. Epping, H. Kampermann, and D. Bruß. Large-scale quantum networks based on graphs. *New J. Phys.* **18**, 053036 (2016).
- [56] M. Epping, H. Kampermann, C. Macchiavello, and D. Bruß. Multi-partite entanglement can speed up quantum key distribution in networks. *New J. Phys.* **19**, 093012 (2017).
- [57] C. Spee and T. Kraft. Transformations in quantum networks via local operations assisted by finitely many rounds of classical communication. [arXiv:2105.01090 \(2021\)](#).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalInformation.pdf](#)