

Relationship between Internet addiction and sleep disturbance in high school students: a cross-sectional study

Mikiko Tokiya

Oita Daigaku

Osamu Itani

Nihon Daigaku Byoin

Yuichiro Otsuka

Nihon Daigaku Byoin

Yoshitaka Kaneita (✉ nusmpublichealth@gmail.com)

Nihon Daigaku Byoin <https://orcid.org/0000-0003-1287-7590>

Research article

Keywords: Pittsburgh Sleep Quality Index, Young Diagnostic Questionnaire, Baseline survey, Multiple logistic regression analysis, Epidemiological study

Posted Date: April 15th, 2020

DOI: <https://doi.org/10.21203/rs.2.18949/v3>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on August 11th, 2020. See the published version at <https://doi.org/10.1186/s12887-020-02275-7>.

Abstract

Background: Worldwide, the increase in the number of Internet users has increased Internet dependence. In adolescents, this dependence interferes with sleep, which is important for the development of psychophysiological capabilities. However, few large-scale surveys have descriptively examined the relationship between Internet addiction (IA) and sleep disturbance using standardized questionnaires. We conducted this study to determine the relationship between sleep disturbance in adolescents and IA based on the categories of the Young Diagnostic Questionnaire (YDQ) through a complete survey of one prefecture in Japan.

Methods: In 2016, a self-report questionnaire was used to survey high school students (n=10,405, age range: 15–16 years) in all 54 day-boarding high schools in the selected prefecture. We defined “sleep disturbance” by scores greater than 5.5 points on the Japanese version of the Pittsburgh Sleep Quality Index. IA was evaluated using the YDQ: “IA,” when 5 of the 8 YDQ items were present; “at-risk,” when either 3 or 4 YDQ items were reported; and “non-IA,” when less than 2 YDQ items were positive. Multiple logistic regression analysis was undertaken with sleep disturbance as the dependent variable, IA as the explanatory variable, and adjustments for 8 items.

Results: High YDQ scores were associated with high prevalence rates of sleep disturbance in both male and female participants. These findings persisted after controlling for other factors in the multiple regression model.

Conclusions: Among Japanese adolescents, there exists a significant independent relationship between IA and sleep disturbances.

Background

The Internet is a network that connects information devices across the world to provide convenient information and communication technology that enables various activities, from exchanging electronic mail and information to shopping. In 2016, when this study was conducted, 48% of all people worldwide used the Internet [1]. In this context, a large survey of Japanese youth found that 6.2% of boys and 9.8% of girls had problematic Internet use [2].

Internet addiction (IA) has been defined as “an impulse-control disorder that does not involve an intoxicant” [3]. This survey examined the concept of IA. Generalized Internet Addiction (GIA) is a concept that was initially introduced by Davis et al. [4] based on a cognitive-behavioral model. More recently, a meta-analysis of 89,281 individuals in 31 countries from 1996 to 2012 reported an IA prevalence rate of 6%, with a median age of 18.42 years (standard deviation [SD], 5.02; range, 12–41) [5]; individuals between 15 and 24 years accounted for approximately 25% of Internet users worldwide [1]. Moreover, this age range includes adolescents, which means that policies pertaining to the issue of IA must address this population as well.

A previous study on IA among adolescents reported a significant relationship between this addiction and psychiatric disturbances, including “interpersonal sensitivity,” “depression,” “anxiety,” “hostility,” and “psychoticism.” [6]. Furthermore, adolescent IA has been reported to be a risk factor for problematic alcohol use in adulthood [7, 8]. Recent studies using functional magnetic resonance imaging have reported that IA is related to structural and functional damage in the prefrontal cortex [9]. With such severe negative impacts on life, the gravity of this problem has been increasingly recognized, and several epidemiological studies have been conducted to determine factors related to IA. For example, a study of 100,000 Japanese youth reported that IA was related to frequency and amount of alcohol consumption [10]. A study of 2,620 Chinese high school students reported a relationship between IA and emotional anxieties and lack of empathy [11]. Excessive smartphone use may be associated with musculoskeletal discomfort and mental health problems [12, 13].

For adolescents, sleep behavior is a daily, routine, lifestyle component that has a major impact on physical and mental health [14]. Moreover, adolescent sleep is important because of its significant effects on the development of key psychophysiological functions, including behavior, emotions, and attention [15-21]. Therefore, it is necessary to investigate any relationships that exist between sleep and IA. Some studies have reported the relationship between IA and depression and sleep disturbance [22, 23], nighttime sleep duration and subjective insomnia [24], poor sleeping habits [25], smartphone dependence [26], and sleep quality [27]. Sleeping habits are associated with other lifestyle habits, such as extracurricular activities and skipped meals [28, 29]. However, the relationship between IA and sleep disturbance in adolescents has not yet been fully analyzed or elucidated, because few large-scale surveys have been undertaken with standard indicators, such as the Pittsburgh Sleep Quality Index (PSQI) [30].

We hypothesized that sleep disorders in puberty are associated with a general degree of Internet dependence but that the association is also attributable, in part, to other lifestyle habits. It is important to take lifestyle habits into account, because they can weaken the relationship between sleep disorders and Internet dependence. Therefore, we conducted an epidemiological study to determine the relationship between IA and sleep disturbance in Japanese high school students by using standardized sleep disorder questionnaires and representative Internet-dependence questionnaires.

Materials And Methods

Study population and design

With the consent of the President of the Association of High School Principals and the prefectural Education Bureau of a certain prefecture in Japan, we sent requests for participation to the principals of all 54 day-boarding high schools within the above prefecture and sent the following documents via the postal service to each principal: (1) letter requesting cooperation, (2) planning document containing the study purpose and method, and (3) the survey form to be used in the study. We stated that a self-administered questionnaire form would be used in the survey, with assured protection of respondent privacy. A total of 10,405 students were registered at the 54 day-boarding high schools.

The survey procedure was as follows: (1) the teachers distributed the following three items: an explanatory document, a self-administered questionnaire form, and an envelope; (2) after filling in their responses in the questionnaire form, the surveyed students placed the completed questionnaire form in the provided collection envelope and sealed the envelope; (3) the teachers collected the sealed envelopes; and (4) the envelopes containing the self-administered questionnaire forms were first unsealed and opened when they were to be used for data input at the assigned research facility. The survey period was from June to December 2016.

Measurement

The survey forms collected information on participant demographic characteristics, sleep disturbance, and IA.

Demographic characteristics

Information was collected on the name of the school, grade, and name and sex of the student. By school names, students were further grouped on the basis of whether they were attending a public school or a private school. Questions on daily-life habits included school-commute time, time spent engaging in school sports (or clubs), time spent on study outside school hours, television-viewing time, and skipped meals. These questions were similar to those used in previous studies among adolescents [10, 31-33] (Appendix A). The questions on emotions and perceptions were measured by depressed mood (mental health) and school-life satisfaction.

We adopted the measure of depressed moods (mental health) from previous studies [31, 33]. The question was “Over the past 30 days, did you have feelings of heaviness or depression more than usual?” We measured school life satisfaction using a 2013 survey conducted by the Cabinet Office on the attitudes of young people in Japan and other countries [34]. The question was “Are you satisfied with your school life? Or are you dissatisfied with it?”

Measurement of sleep disturbance: Japanese version of the Pittsburgh Sleep Inventory

Sleep disturbance was evaluated by the Japanese version of the Pittsburgh Sleep Quality Index (J-PSQI) [35-37]. On the basis of previous studies, scores ≥ 5.5 points on the J-PSQI were considered indicative of sleep disturbance [35-37].

Measurement of IA: Japanese version of the Young Diagnostic Questionnaire

IA was evaluated using the Young Diagnostic Questionnaire (YDQ) [3, 38-44]. For the present study, we used the Japanese version of the YDQ (J-YDQ) that has been used in previous studies [31]. The J-YDQ is an evaluation tool composed of 8 questions, which are rated 1 point for “yes” and 0 point for “no,” with the total score ranging from 0 to 8 points. The participants were then grouped into three categories: “IA,” when 5 of the 8 YDQ items were present; “at-risk,” when either 3 or 4 YDQ items were reported; and “non-IA,” when less than 2 YDQ items were positive. [25, 33, 39, 43, 45, 46].

Ethical considerations

The participation of students in the present study was voluntary. As our cohort included 15- to 16-year-old adolescents, we obtained written informed consent either directly from the students, if their supervising teacher confirmed that their judgment was acceptable, or from their parents, if the supervising teacher thought that the parents' consent was necessary. The following statements were included in the consent document distributed to students and their families: (1) the survey was part of an epidemiological study and involved neither an evaluation for school grading nor any type of punishment; (2) students were free to cooperate in the survey, and failure to cooperate would not incur any disadvantage; (3) the school teachers would not view the responses provided; and (4) respondent privacy would be strictly protected. The study forms were stored securely, and data were entered into a password-protected database. Data were anonymized prior to the analysis by deleting all personal identifiers. The Faculty of Medicine, Oita University Ethics Committee approved the study (approval no. 932).

Statistical analysis

Students who did not completely fill out the J-PSQI and the YDQ were excluded from the analysis. First, distribution by sex (boys or girls) was plotted for the J-PSQI and YDQ scores. Second, prevalence rates for different categories of sleep disturbance and IA were calculated separately for each sex by using the chi-square test. Furthermore, for each of the three YDQ categories, prevalence rates for sleep disturbance were calculated for each sex separately. Finally, multiple logistic regression analysis was conducted to investigate the relationship between IA (explanatory variable) and sleep disturbance (dependent variable). The type of school, school-commute time, sports and club time, outside-class study time, television-viewing time, skipped meals, depressed mood (mental health), and school-life satisfaction rates were used as adjusted variables. The Statistical Package for Social Sciences version 22 (SPSS, IBM Corp. NY, USA) for Windows was used for all statistical analyses. A p -value <0.05 was considered statistically significant.

Results

A flowchart of the participant-selection process is shown in Figure 1. Of the 54 schools ($n=10,405$) that were requested to participate, 40 schools responded positively. At the time of the study, there were 7,186 first-year (of the 3-year program) high school students, of whom 6,950 provided responses (response rate: 96.7%). Of these, 5,264 students (2,635 boys and 2,629 girls) provided informed consent (effective response rate: 73.3%).

Figure 2 shows the distribution of J-PSQI and YDQ scores. The J-PSQI scores for both boys and girls were symmetrically distributed around a cutoff point value of 5.5 points. The mean and SD of the total J-PSQI score was 5.51 ± 2.63 (range: 0–17) and 5.98 ± 2.62 (range: 0–18) for boys and girls, respectively. With regard to the YDQ scores, 0 point was the most frequent score for both boys and girls, with higher scores among a small number of participants.

Table 1 shows the percentages of students with sleep disturbance and the number of participants included within the three YDQ categories. Students defined as having sleep disturbance comprised 50.5% of all participants. In both boy and girl participants, a higher percentage of sleep disturbance was observed in the following groups: private high school ($p < 0.05$), students with a high frequency of skipped meals within a 1-week period ($p < 0.001$), and proportions of students reporting higher scores on variables of depressed mood (mental health; $p < 0.001$). Overall, based on the YDQ scores, 9.4% of boys and 14.8% of girls were classified with IA, whereas 23.1% of boys and 28.1% of girls were graded as at high risk of IA. In both boys and girls with IA, we observed the proportions of students who reported higher scores for skipped meals ($p < 0.001$). In the study cohort of both boys and girls, approximately 60% of those who answered yes or often for depressed mood (mental health) and 60–70% of those who answered dissatisfied for school-life satisfaction were found to have sleep disorders ($p < 0.001$ for both). Furthermore, boys with IA spent less time engaging in school sports or in clubs, whereas girls with IA spent less time on outside-school study ($p < 0.001$ for both). For boys and girls with high YDQ scores, we observed that a higher proportion of students answered “yes” or “often” about having a depressive mood (mental health) and ‘dissatisfied’ about school-life satisfaction. ($p < 0.001$ for both). Moreover, girls at risk of IA did not study much outside of school hours ($p < 0.001$).

Table 2 presents the rates of students with sleep disturbance for each of the three categories of YDQ. For the overall study population, a higher YDQ score in any category was associated with a higher rate of students with sleep disturbance.

Table 3 shows the results of the multiple logistic regression analysis of the relationship between YDQ categories and sleep disturbance. In the entire study population, a higher YDQ score in any category was associated with higher unadjusted odds ratios (ORs) for sleep disturbance ($p < 0.001$). The association between the YDQ scores and sleep disturbance remained significant ($p < 0.001$), even after adjusting for lifestyle variables.

Discussion

The present study aimed to clarify the relationship between sleep disturbance in adolescents and IA. We found a relationship between adolescent sleep disturbance and IA in one prefecture in Japan and found that sleep disturbances are more prevalent in categories with higher YDQ scores. Furthermore, the results of the multivariate analysis revealed significantly higher adjusted ORs between categories with high YDQ and sleep disorders, which suggests a significant relationship between higher YDQ scores (3–4 or ≥ 5) and sleep disorders.

Despite the importance of adolescent sleep, sleep disturbance was present in more than half of the study participants. Furthermore, the increase in sleep disturbance among the young was similar to recent studies using the PSQI [47, 48], suggesting that this is not limited to this study area. In this study, sleep disturbance was more frequent in boys who participated in school sports (or clubs), skipped meals overall, and had depressive moods (mental health). These results are similar to those of previous studies

linking regular sleep habits to psychological and physical health [14]. In addition, the prevalence of sleep disturbance was higher in private schools and for students with longer commutes. In general, private schools often have longer commute distances; so, it may be associated with reduced sleep time as shown previously [49]. Previous studies have suggested that sleep quality is related to health and emotions among the youth [14, 50]. Future longitudinal studies should examine changes over time for these and other variables to better understand the relationship between sleep disorders and Internet dependence.

In addition,

The association between YDQ scores and sleep disorders is similar to that reported by Bakken et al. [51] from a study among Norwegian adults, age 16 or older, where individuals with high YDQ scores had significantly higher prevalence rates of sleep disturbance than non-problematic Internet users. Furthermore, the present study found differences between sexes, with a higher sleep disturbance adjusted OR in girls with IA than in boys. This finding is similar to the results of Durkee et al. [25], who confirmed a significant relationship between insufficient sleep and IA in girl participants, which is in line with the findings of a previous study [25].

There are several possible mechanisms for the relationship between sleep disturbance and IA. Our findings suggest that IA itself results in sleep disturbance, which is in line with a previous study, which reported that problematic Internet use was a significant predictor of sleep disturbance [22]. Moreover, Chen et al. reported that IA predicted a “disturbed circadian rhythm” as leading to sleep disturbance [30], which is one explanation for the relationship between IA and sleep disturbance.

The second possible mechanism is the converse of the one discussed above, whereby sleep disturbance might lead to the development of IA. In a longitudinal study, Chen et al. reported that difficulty in falling asleep and nocturnal awakening were predictors of IA [30].

A third possible mechanism is that both conditions contribute to each other. In other words, sleep disturbance may contribute to IA, and IA may contribute to sleep disturbance. Several studies on adults have confirmed that, in brain imaging, both sleep disturbance and IA cause changes in the gray matter [52, 53]. A study of retired military personnel showed that, regardless of any coincidental psychiatric state, individuals with a high PSQI score presented with reduced volume of the entire cortex and frontal lobes [53]. Another study that did not control for sleep disturbance reported that individuals with IA reportedly had reduced gray matter density [52]. These findings suggest the possibility that IA may cause organic (structural) changes in sleep-related neural pathways.

The following three points can be considered as the strengths of this study. First, the sample size was adequate to ensure statistical power. Second, to investigate the relationship between sleep disturbance and IA, we used the PSQI and YDQ, which have been frequently used as standard indices in several epidemiological surveys [5, 10, 25, 39, 41, 43, 51, 52, 54-59]. Third, in our analysis, we evaluated the relationships between sleep disturbance and IA for each of the three categories of the YDQ, including

at-risk Internet use; this category has not been sufficiently investigated in previous epidemiological studies.

Nonetheless, the present study has a number of limitations. First, the present study deals with the results of a cross-sectional survey, which implies we cannot formulate any conclusion about the direction of causality. Second, statistical analysis should have taken into account schools as cluster units. Third, we did not adjust ORs for all the items that may be related to IA. For example, we did not ask questions about other psychiatric disorders such as attention deficit and hyperactivity disturbance (ADHD), which is reportedly associated with IA [60-62] and sleep disturbance [62] in adolescents. In the present study, all participants were enrolled as daytime high school students and regularly attended school. In such a scenario, the number of students with ADHD is considered low. Fourth, as we conducted the survey within each of the schools, non-attending students could not participate, and future surveys that will enable participation of non-attending students should be undertaken. Fifth, our survey population was limited to students in a single prefecture in Japan; thus, there was certainly a sampling bias. Last, we did not investigate specific Internet-use disorders [4, 63-69], which need to be investigated in detail for preventive measures to be developed.

Conclusions

In summary, we observed that high Internet dependence was related to sleep disturbance in high school students within a prefecture in Japan. Studies have suggested that both sleep disturbances and IA affect the gray matter in the brain. Future research should undertake longitudinal surveys that investigate the factors related to the occurrence of IA and sleep disturbance.

Abbreviations

IA: Internet addiction

YDQ: Young Diagnostic Questionnaire

J-PSQI: The Japanese version of the Pittsburgh Sleep Quality Index

J-YDQ: The Japanese version of the Young Diagnostic Questionnaire

Declarations

Ethics approval and consent to participate

The participation of students in the present study was voluntary. Our cohort included 15- to 16-year-old adolescents, and all students eligible to participate in this study had completed junior high school courses. The consent form distributed to the students or their families stated that: (1) the survey was part of an epidemiological study and involved neither an evaluation for school grading, nor any type of

punishment; (2) the students were free to choose whether to participate in the survey and failure to participate would not incur any disadvantage; (3) the school teachers would not view the responses provided; and (4) the privacy of students would be strictly protected. This survey was conducted among students who provided written voluntary informed consent and had parental consent for study participation. The present study was approved (approval no. 932) by the Oita University Ethics Committee.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available due to the sensitive nature of the raw data; however, all pertinent study datasets are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no actual or potential competing financial interests.

Funding

This study was supported by a Grant-in-Aid for Scientific Research (grant no. JP 17K09117) conferred by the JSPS KAKENHI. The funding bodies had no role in the design of the study and collection, analysis, and interpretation of data, and in writing the manuscript.

Authors' contributions

MT, YK, OI, and YO designed the survey questionnaire, and MT, YK, OI, and YO conducted the survey. MT wrote the initial manuscript draft and was responsible for submission for publication. YK, OI, and YO undertook a critical revision of the manuscript. MT conducted the initial analyses. YK, OI, and YO provided important feedback on aspects to improve the study conduct, and YK shared critical insights and suggestions to optimize the study conduct. All authors have read and approved the final manuscript.

Acknowledgements

The authors express their deep gratitude to all of the high school student participants and to their teachers for their cooperation of this study. The authors also express heartfelt thanks to Yukiko Abe for her cooperation in the collection of survey forms and data analysis. The authors thank Editage (www.editage.com) for English language editing.

References

1. ICT Facts and Figures 2017. <https://www.itu.int/en/ITU-D/Statistics/Documents/facts/ICTFactsFigures2017.pdf>. Accessed on 01 Sep 2019.
2. Mihara S, Osaki Y, Nakayama H, Sakuma H, Ikeda M, Itani O, et al. Internet use and problematic Internet use among adolescents in Japan: a nationwide representative survey. *Addict Behav Rep.* 2016;4:58-64.
3. Young KS. Internet addiction: the emergence of a new clinical disorder. *Cyberpsychol Behav.* 1998;1(3):237-44.
4. Davis RA. A cognitive-behavioral model of pathological Internet use. *Comput Human Behav.* 2001;17(2):187-95.
5. Cheng C, Li AY. Internet addiction prevalence and quality of (real) life: a meta-analysis of 31 nations across seven world regions. *Cyberpsychol Behav Soc Netw.* 2014;17(12):755-60.
6. Dong G, Lu Q, Zhou H, Zhao X. Precursor or sequela: pathological disorders in people with Internet addiction disorder. *PloS One.* 2011;6(2):e14703.
7. Gamez-Guadix M, Calvete E, Orue I, Las Hayas C. Problematic Internet use and problematic alcohol use from the cognitive-behavioral model: a longitudinal study among adolescents. *Addict Behav.* 2015;40:109-14.
8. Lee BH, Lee HK. Longitudinal study shows that addictive Internet use during adolescence was associated with heavy drinking and smoking cigarettes in early adulthood. *Acta Paed.* 2017;106(3):497-502.
9. Park B, Han DH, Roh S. Neurobiological findings related to Internet use disorders. *Psych Clin Neurosci.* 2017;71(7):467-78.
10. Morioka H, Itani O, Osaki Y, Higuchi S, Jike M, Kaneita Y, et al. The association between alcohol use and problematic Internet use: a large-scale nationwide cross-sectional study of adolescents in Japan. *J Epidemiol.* 2017;27(3):107-11.
11. Cao F, Su L. Internet addiction among Chinese adolescents: prevalence and psychological features. *Child.* 2007;33(3):275-81

12. Yang SY, Lin CY, Huang YC, Chang JH. Gender differences in the association of smartphone use with the vitality and mental health of adolescent students. *J Am Coll Health*. 2018;66(7):693-701.
13. Yang SY, Chen MD, Huang YC, Lin CY, Chang JH. Association between smartphone use and musculoskeletal discomfort in adolescent students. *J Com Health*. 2017;42(3):423-30.
14. Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, et al: Recommended amount of sleep for pediatric populations: a consensus statement of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2016;12(6):785-6.
15. Beebe DW, Rose D, Amin R. Attention, learning, and arousal of experimentally sleep-restricted adolescents in a simulated classroom. *J Adolesc Health*. 2010;47(5):523-5.
16. Brand S, Kirov R. Sleep and its importance in adolescence and in common adolescent somatic and psychiatric conditions. *Int J Gen Med*. 2011;4:425-42.
17. Carskadon MA, Acebo C, Jenni OG. Regulation of adolescent sleep: implications for behavior. *Ann New York Acad Sci*. 2004;1021:276-91.
18. Casement MD, Keenan KE, Hipwell AE, Guyer AE, Forbes EE. Neural reward processing mediates the relationship between insomnia symptoms and depression in adolescence. *Sleep*. 2016;39(2):439-47.
19. Dahl RE, Lewin DS. Pathways to adolescent health sleep regulation and behavior. *J Adolesc Health*. 2002;31(6 Suppl):175-84.
20. Fallone G, Owens JA, Deane J. Sleepiness in children and adolescents: clinical implications. *Sleep Med Rev*. 2002;6(4):287-306.
21. Wolfson AR, Carskadon MA. Sleep schedules and daytime functioning in adolescents. *Child Dev*. 1998;69(4):875-87.
22. Tan Y, Chen Y, Lu Y, Li L. Exploring associations between problematic internet use, depressive symptoms and sleep disturbance among southern Chinese adolescents. *Int J Environ Res Pub Health*. 2016;13(3).
23. Alimoradi Z, Lin C-Y, Broström A, Bülow PH, Bajalan Z, Griffiths MD, et al. Internet addiction and sleep problems: a systematic review and meta-analysis. *Sleep Med Rev*. 2019;47:51-61.
24. Yen CF, Ko CH, Yen JY, Cheng CP. The multidimensional correlates associated with short nocturnal sleep duration and subjective insomnia among Taiwanese adolescents. *Sleep*. 2008;31(11):1515-25.
25. Durkee T, Carli V, Floderus B, Wasserman C, Sarchiapone M, Apter A, et al. Pathological internet use and risk-behaviors among European adolescents. *Int J Environ Res Pub Health*. 2016;13(3).
26. Yang SY, Chen KL, Lin PH, Wang PY. Relationships among health-related behaviors, smartphone dependence, and sleep duration in female junior college students. *Soc Health Behav*. 2019;2:26-31.
27. Wang PY, Chen KL, Yang SY, Lin PH. Relationship of sleep quality, smartphone dependence, and health-related behaviors in female junior college students. *PloS One*. 2019;14(4):e0214769.
28. Bartel KA, Gradisar M, Williamson P. Protective and risk factors for adolescent sleep: a meta-analytic review. *Sleep Med Rev*. 2015;21:72-85.

29. Kaneita Y, Ohida T, Osaki Y, Tanihata T, Minowa M, Suzuki K, et al. Insomnia among Japanese adolescents: a nationwide representative survey. *Sleep*. 2006;29(12):1543-50.
30. Chen YL, Gau SS. Sleep problems and Internet addiction among children and adolescents: a longitudinal study. *J Sleep Res*. 2016;25(4):458-65.
31. Munezawa T, Kaneita Y, Osaki Y, Kanda H, Minowa M, Suzuki K, et al. The association between use of mobile phones after lights out and sleep disturbances among Japanese adolescents: a nationwide cross-sectional survey. *Sleep*. 2011;34(8):1013-20.
32. Itani O, Kaneita Y, Ikeda M, Kondo S, Yamamoto R, Osaki Y, et al. Disorders of arousal and sleep-related bruxism among Japanese adolescents: a nationwide representative survey. *Sleep Med*. 2013;14(6):532-41.
33. Itani O, Kaneita Y, Munezawa T, Ikeda M, Osaki Y, Higuchi S, et al. Anger and impulsivity among Japanese adolescents: a nationwide representative survey. *J Clin Psych*. 2016;77(7):e860-6.
34. Office of the Director General for Policy Planning for Policies on Cohesive Society CO. International Survey of Youth Attitude 2013. https://www8.cao.go.jp/youth/english/survey/2013/pdf_index.html. Accessed on 01 Sep 2019.
35. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psych Res*. 1989;28(2):193-213.
36. Doi Y, Minowa M, Okawa M, Uchiyama M. Development of the Japanese version of the Pittsburgh Sleep Quality Index. *Jap J Psych Treat*. 1998;13(3):755-63 (in Japanese).
37. Doi Y, Minowa M, Uchiyama M, Okawa M, Kim K, Shibui K, et al. Psychometric assessment of subjective sleep quality using the Japanese version of the Pittsburgh Sleep Quality Index (PSQI-J) in psychiatric disordered and control subjects. *Psych Res*. 2000;97(2-3):165-72.
38. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 4th ed. Arlington: American Psychiatric Publishing; 1994.
39. Siomos KE, Dafouli ED, Braimiotis DA, Mouzas OD, Angelopoulos NV. Internet addiction among Greek adolescent students. *Cyberpsychol Behav*. 2008;11(6):653-7.
40. Shek DT, Tang VM, Lo CY. Internet addiction in Chinese adolescents in Hong Kong: assessment, profiles, and psychosocial correlates. *ScientificWorldJournal*. 2008;8:776-87.
41. Osada H. Internet addiction in Japanese college students: is Japanese version of Internet Addiction Test (JIAT) useful as a screening tool? *Bull Senshu Univ Sch Hum Sci Psychol*. 2013;3(1):71-80.
42. Laconi S, Rodgers RF, Chabrol H. The measurement of Internet addiction: A critical review of existing scales and their psychometric properties. *Comput Human Behav*. 2014;41:190-202.
43. Johansson A, Gotestam KG. Internet addiction: characteristics of a questionnaire and prevalence in Norwegian youth (12-18 years). *Scand J Psychol*. 2004;45(3):223-9.
44. Aboujaoude E. Problematic Internet use: an overview. *World Psych*. 2010;9(2):85-90.
45. Christos CF, Constantinos CF, Apostolos PK. Internet addiction among Greek university students: demographic associations with the phenomenon, using the Greek version of Young's Internet

- Addiction Test. *Int J Econ Sci App Res.* 2010;3(1):49-74.
46. Dowling NA, Quirk KL. Screening for Internet dependence: do the proposed diagnostic criteria differentiate normal from dependent Internet use? *Cyberpsychol Behav.* 2009;12(1):21-7.
 47. Alotaibi AD, Alosaimi FM, Alajlan AA, Bin Abdulrahman KA. The relationship between sleep quality, stress, and academic performance among medical students. *J Family Community Med.* 2020;27(1):23-8.
 48. Becker SP, Jarrett MA, Luebbe AM, Garner AA, Burns GL, Kofler MJ. Sleep in a large, multi-university sample of college students: sleep problem prevalence, sex differences, and mental health correlates. *Sleep Health.* 2018;4(2):174-81.
 49. Hall WA, Nethery E. What does sleep hygiene have to offer children's sleep problems? *Paediatr Respir Rev.* 2019;31:64-74.
 50. Hosker DK, Elkins RM, Potter MP. Promoting mental health and wellness in youth through physical activity, nutrition, and sleep. *Child Adolesc Psychiatr Clin N Am.* 2019;28(2):171-93.
 51. Bakken IJ, Wenzel HG, Gotestam KG, Johansson A, Oren A. Internet addiction among Norwegian adults: a stratified probability sample study. *Scand J Psychol.* 2009;50(2):121-7.
 52. Zhou Y, Lin FC, Du YS, Qin LD, Zhao ZM, Xu JR, et al. Gray matter abnormalities in Internet addiction: a voxel-based morphometry study. *Eur J Radiol.* 2011;79(1):92-5.
 53. Chao LL, Mohlenhoff BS, Weiner MW, Neylan TC. Associations between subjective sleep quality and brain volume in Gulf War veterans. *Sleep.* 2014;37(3):445-52.
 54. Lund HG, Reider BD, Whiting AB, Prichard JR. Sleep patterns and predictors of disturbed sleep in a large population of college students. *J Adolesc Health.* 2010;46(2):124-32.
 55. Cheung LM, Wong WS. The effects of insomnia and Internet addiction on depression in Hong Kong Chinese adolescents: an exploratory cross-sectional analysis. *J Sleep Res.* 2011;20(2):311-7.
 56. Smagula SF, Stone KL, Fabio A, Cauley JA. Risk factors for sleep disturbances in older adults: evidence from prospective studies. *Sleep Med Rev.* 2016;25:21-30.
 57. Kuss DJ, Griffiths MD, Karila L, Billieux J. Internet addiction: a systematic review of epidemiological research for the last decade. *Curr Pharm Des.* 2014;20(25):4026-52.
 58. Durkee T, Kaess M, Carli V, Parzer P, Wasserman C, Floderus B, et al. Prevalence of pathological Internet use among adolescents in Europe: demographic and social factors. *Addiction.* 2012;107(12):2210-22.
 59. Shek DT, Yu L. Internet addiction phenomenon in early adolescents in Hong Kong. *ScientificWorldJournal.* 2012;2012:104304.
 60. Yen JY, Ko CH, Yen CF, Wu HY, Yang MJ. The comorbid psychiatric symptoms of Internet addiction: attention deficit and hyperactivity disorder (ADHD), depression, social phobia, and hostility. *J Adolesc Health.* 2007;41(1):93-8.
 61. Ko CH, Yen JY, Chen CS, Yeh YC, Yen CF. Predictive values of psychiatric symptoms for Internet addiction in adolescents: a 2-year prospective study. *Arch Ped Adol Med.* 2009;163(10):937-43.

62. Weinstein A, Yaacov Y, Manning M, Danon P, Weizman A. Internet addiction and attention deficit hyperactivity disorder among schoolchildren. *Isr Med Assoc J.* 2015;17(12):731-4.
63. Brand M, Young KS, Laier C, Wolfling K, Potenza MN. Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: an interaction of Person-Affect-Cognition-Execution (I-PACE) model. *Neurosci Biobehav Rev.* 2016;71:252-66.
64. Wu TY, Lin CY, Arestedt K, Griffiths MD, Brostrom A, Pakpour AH. Psychometric validation of the Persian nine-item Internet Gaming Disorder Scale - Short Form: Does gender and hours spent online gaming affect the interpretations of item descriptions? *J Behav Addict.* 2017;6(2):256-63.
65. Lin CY, Brostrom A, Nilsen P, Griffiths MD, Pakpour AH. Psychometric validation of the Persian Bergen Social Media Addiction Scale using classic test theory and Rasch models. *J Behav Addict.* 2017;6(4):620-9.
66. Lin CY, Imani V, Brostrom A, Arestedt K, Pakpour AH, Griffiths MD. Evaluating the psychometric properties of the 7-Item Persian Game Addiction Scale for Iranian adolescents. *Front Psychol.* 2019;10:149.
67. Lin C-Y, Imani V, Broström A, Nilsen P, Fung XCC, Griffiths MD, et al. Smartphone application-based addiction among Iranian adolescents: a psychometric study. *Intl J Ment Health Ad.* 2019;17(4):765-80.
68. Yam CW, Pakpour AH, Griffiths MD, Yau WY, Lo CM, Ng JMT, et al. Psychometric testing of three Chinese online-related addictive behavior instruments among Hong Kong University students. *Psych Quarterly.* 2019;90(1):117-28.
69. Lin C-Y, Lin C-K, Imani V, Griffiths MD, Pakpour AH. Evaluation of the Selfitis Behavior Scale across two Persian-speaking countries, Iran and Afghanistan: advanced psychometric testing in a large-scale sample. *Int J Ment Health Ad.* 2019. doi.org/10.1007/s11469-019-00124-y.

Tables

Due to technical limitations, tables are only available as a download in the supplemental files section.

Figures




Figure 1

Flowchart of the participant-selection process.

Figure. Distribution of J-PSQI and YDQ Internet addiction Score




Figure 2

Distribution of J-PSQI and YDQ Internet Addiction Scores.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [renamedf5832sup.docx](#)
- [Tables.docx](#)
- [Appendix.docx](#)